

7.1 Wave Nature of Matter

De Broglie Wavelength
 Diffraction of electrons
 Uncertainty Principle
 Wave Function
 Tunneling

Wave properties of matter

Material particles behave as waves with a wavelength given by the De Broglie wavelength (Planck's constant/momentum)

$$\lambda = \frac{h}{p}$$

The particles are diffracted by passing through an aperture in a similar manner as light waves.

The wave properties of particles mean that when you confine it in a small space its momentum (and kinetic energy) must increase. (uncertainty principle) This is responsible for the size of the atom.

De Broglie Wavelength



Lois De Boglie

Momentum of a photon - inverse to wavelength.

$$p = \frac{E}{c} \quad \text{Einstein's special relativity theory}$$

since $E = \frac{hc}{\lambda}$


$$p = \frac{h}{\lambda}$$

Wavelength of a particle- inverse to momentum.

$$\lambda = \frac{h}{p}$$

De Broglie proposed that this wavelength applied to material particles as well as for photons. (1924)

Wave properties



Diffraction of waves
 Increases as the ratio λ/a

de Broglie wavelength of a baseball




$m = 0.15 \text{ kg}$
 $v = 45 \text{ m/s}$

A baseball with a mass of 0.15 kg is pitched at 45 m/s
 What is its De Broglie wavelength?

$$\lambda = \frac{h}{p} = \frac{h}{mv} = \frac{6.6 \times 10^{-34} \text{ Js}}{(0.15 \text{ kg})(45 \text{ m/s})} = 9.8 \times 10^{-35} \text{ m}$$

Diffraction effects of a baseball are negligible

de Broglie wavelength of an electron



Find the de Broglie wavelength of a 1000 eV electron.

$$KE = \frac{1}{2}mv^2 = \frac{m^2v^2}{2m} = \frac{p^2}{2m} = eV$$

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2meV}}$$

$V = 1000 \text{ V}$

atomic dimensions


$$\lambda = \frac{6.6 \times 10^{-34} \text{ Js}}{\sqrt{2(9.1 \times 10^{-31} \text{ kg})(1.6 \times 10^{-19} \text{ C})(1000 \text{ J/C})}} = 4.0 \times 10^{-11} \text{ m}$$

Proof of the wave nature of electrons by Electron Diffraction

- Davisson – Germer Experiment
- Thompson Experiment

Showed the wave nature of light by diffraction of electrons by crystals.

Diffraction of light from crystals




Crystals act as a three-dimensional diffraction grating
Light with wavelength close to the inter-atomic spacing (x-rays) is diffracted.

Diffraction of electrons from crystals

Davisson-Germer Experiment (1927)

An electron beam is scattered from a crystal



The scattered beam shows a diffraction pattern expected for the crystal spacing.

Comparison between electron diffraction and x-ray diffraction


George Thomson (1927)

Aluminum powder Diffraction pattern

Either x-rays or electrons


The electron wavelength was adjusted to the same value as the x-ray by varying the voltage.

The diffraction pattern for x-rays and electrons are very similar.



Material particles have wave properties.


Electron microscopy




$$\lambda = \frac{h}{mv}$$

Shorter wavelength can be obtained by increasing v , the speed of the electron.

Electron microscopy




electron microscope pictures of ribosomes



model of ribosome

Electron microscopy can be used to image structures of molecules.

Diffraction of particles Probabilistic Interpretation of the wave amplitude.




The amplitude² is interpreted as the **probability** of the particle hitting the screen at a certain position
This is true for electrons as well as photons.

Wavefunction

In quantum mechanics the result of an experiment is given in terms of a wavefunction Ψ . The square of the wavefunction Ψ^2 is the probability of the particle being at a certain position.

The wavefunction can be calculated using the Schrödinger Equation. For instance for electrons in an atom.

Wave property of particles




Decreasing the slit reduces Δx
But increases the width of the diffraction, Δv_x

When Δx decreases, Δp_x increases.

Uncertainty Principle

Particle passing through a slit --
The uncertainty in position is Δx

The uncertainty in the x component of momentum is $\Delta p_x = m\Delta v_x$



The particle is diffracted

$$\Delta x \sin \theta = \lambda = \frac{h}{mv}$$

$$\sin \theta = \theta = \frac{\Delta v_x}{v}$$

$$\Delta x \frac{\Delta v_x}{v} = \frac{h}{mv}$$

Therefore $\Delta x \Delta p_x = h$


most often written as an inequality

$$\Delta x \Delta p_x \geq \frac{h}{4\pi}$$

The position and velocity cannot be known with unlimited certainty.

The size of an atom

What accounts for the size of the hydrogen atom?



$$PE = -\frac{k_0 e^2}{r}$$

PE \rightarrow - infinity as $r \rightarrow$ zero

Classical electrostatics predicts that the potential energy of the hydrogen atom should go to -infinity

The finite size of the atom is a quantum mechanical effect.

As the size of the atom r decreases - The uncertainty in the momentum increases until the uncertainty limits the momentum. Then **the kinetic energy must increase** due to the uncertainty principle.

Use linear momentum as a rough estimate.

$$\Delta x \Delta p_x \approx h$$

$$\Delta p_x \approx \frac{h}{\Delta x} = \frac{h}{r}$$




$$\Delta p_x = \frac{h}{2r} \approx p_x \quad p_x \text{ cannot be smaller than } \Delta p_x$$

$$KE = \frac{1}{2} m v_x^2 = \frac{p_x^2}{2m} \approx \frac{h^2}{8r^2 m}$$


KE increases as $1/r^2$

$E = KE + PE$ goes through a minimum as a function of r




Tunneling across a barrier

a macroscopic object impinging on a barrier the object cannot penetrate within the barrier.





a wave particle impinging on a barrier can penetrate within the barrier for distance. and go through the barrier if it is thin enough.



The probability of tunneling decreases exponentially with the width of the barrier.


Electron Tunneling Microscope




$$\text{Electron_Probability} \propto |\Psi|^2 \approx \Psi_0^2 e^{-2\alpha r}$$

$$\alpha \sim 10 \text{ nm}^{-1}$$

The tunneling probability falls off exponentially with r and is a sensitive function of distance



Tunneling in Photosynthesis



Bacterial Reaction Center

Electron transfer time vs distance

Electron transfer is due to tunneling. Transfer times are a sensitive function of distance.