

Physics 161: Black Holes: Lecture 11: 28 Jan 2011

Professor: Kim Griest

11 Shooting light rays into black holes, Inside a black hole

Let's calculate radial motion into black holes another way, a way that is often very useful, because it bypasses the geodesic equations. Let's calculate how long it takes to shoot a ray of light into black hole.

For light we know the invariant interval $ds^2 = 0$, that is the metric distance, aka proper time, is 0. So for light itself how long does it take to get into a black hole? It takes the same time it take light to go anywhere, zero! From light's point of view no time ever passes since $d\tau = 0$ always. OK, that means we set the metric equal to zero for light. This is the very useful trick. Considering radial infall, we can also set $d\phi = d\theta = 0$, and get

$$dt = \frac{dr}{(1 - r_S/r)}.$$

Integrating both sides from $t = 0$ at $r = r_0$ to $t = t$ at $r = r$, we find

$$t = r_0 - r + r_S \ln \left(\frac{r_0 - r_S}{r - r_S} \right).$$

From our previous work we expect the time for light go from $r = 30$ km to $r = r_S$ to be less than a millisecond. But plugging into the above equation we get a factor $\ln((30 - 8.85)/(8.85 - 8.85)) \rightarrow \ln(\infty) \rightarrow \infty$. There is a logarithmic divergence and again we see we never get into the black hole!

Of course, from the point of view of the light ray, or the falling guy, they get in and are crushed within a millisecond. It is just that time viewed from far away runs differently at the horizon of a black hole. The problem is with the use of this far away time. Also, while it is true that everything that ever falls into a black hole seems to "hang up" at the horizon, it is not that case that someone looking closely at a black hole sees all that junk. Remember the light coming to you from the falling objects redshifts to infinity and therefore those objects become invisible very quickly.

As an interesting aside consider what would happen if you tried to lower yourself slowly into a black hole on a very strong rope. I'm not going to do the calculation but the effective acceleration of gravity you would feel would increase without limit. The effective g , which is $GM/r^2 = 9.8$ m/s² here on Earth becomes

$$g = (GM/r^2)(1 - r_S/r)^{-1/2}.$$

Thus the force becomes infinitely strong at $r = r_S$, the rope will break and you will fall to your death.

11.1 Inside the Black Hole

The Schwarzschild metric is the solution to Einstein's GR equations in the vacuum around a spherically symmetric object. Thus we expect these to work even if the object is smaller than r_S . What happens at

and inside $r = r_S = 2GM/c^2$? The metric is bad at $r = r_S$, but actually it is OK inside.

$$ds^2 = - \left(1 - \frac{2GM}{rc^2}\right) dt^2 + \left(1 - \frac{2GM}{rc^2}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

First consider lightcones, the causally connected past and future. How did we find these in Special Relativity? Considering just one space and one time dimension: $ds^2 = -dt^2 + dx^2 = 0$, for light. This defines the null geodesics. Solving we find $dt = \pm dr$, or $dt/dr = \pm 1$, which implies the lightcones are lines of 45° in a spacetime diagram. Remember also that above the 45° lines is the timelike $ds^2 < 0$ future, while outside the lines is the $ds^2 > 0$ spacelike elsewhere. In Special Relativity all lightcones have these 45° lines.

How about in General Relativity. Again we set $ds^2 = 0$ to find the null geodesics, giving $-dt^2(1 - r_S/r) + dr^2/(1 - r_S/r) = 0$, or

$$\frac{dt}{dr} = \pm \frac{1}{1 - \frac{r_S}{r}}.$$

Thus the lightcones are not 45° lines. As $r \rightarrow \infty$, $r_S/r \rightarrow 0$, so $dt/dr \rightarrow \pm 1$, and the lightcones are at 45° , but closer to the spherical mass the angles are smaller, $dt/dr > \pm 1$. Thus the lightcones squeeze-up. Thus the path of light as it travels towards a spherical mass in a spacetime diagram is not on a 45° line, but on a curved line that gets more steep as it approaches the Schwarzschild radius.

Fig: Path of light as it approaches a black hole

We can calculate the angle of the lightcone at any value of r very simply. Just use

$$\frac{dt}{dr} = \tan \theta.$$

Thus at 30 km from a $3M_\odot$ mass hole the angle of the lightcone is $dt/dr = 1/(1 - 8.85/30) = 1.42$, or $\theta = 55^\circ$. This angle goes to 90° at $r = r_S$. So we see that right at $r = r_S$, $dt/dr \rightarrow \infty$, and no progress in r is possible! Not even light can make it into the black when far-away time is used as the coordinate. How does this jive with the fact that we calculated things can fall into black holes in just milliseconds? The point is that t is time measured by someone far away, not the traveler.

Now look what happens to the metric when $r < r_S$. Notice that since $r_S/r > 1$, the terms $(1 - r_S/r)$ become less than zero.

$$ds^2 = -dt^2(1 - r_S/r) + dr^2/(1 - r_S/r) + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2.$$

Thus we see that the t coordinate and the r coordinate terms switch signs! Remember that in GR r is not a distance, it is a coordinate. You have to use the metric to find distances. Same with t . The minus sign is how we recognize the time coordinate in GR, so this means that r becomes timelike and t becomes spacelike! If you try to find the lightcones inside the black hole you see they flip over sideways!

Fig: Lightcones around and inside a black hole

The same force that moves everyone towards the future now moves things towards $r = 0$, which is the future lightcone! This one way of understanding the reason that once inside the black hole you can't get out. There is no force in nature that can move you backward in time. Thus inside a hole there is no force in nature that can move something towards larger r ! Note that while proper time increases, t actually decreases. This is not a problem since t is no longer a time dimension. It is a space dimension; r is the time. We can see this explicitly by considering two points inside, say near $r = 2$ km and $r = 2.1$ km,

so $dr = 0.1$ km. Take $dt = 0$ and calculate the invariant interval: $ds^2 = +dr^2/(1 - 8.85/2) = -0.0029$ km < 0 , that is the interval is timelike, and thus these points can be causally connected! Similarly, two events with $dr = 0$ and $dt \neq 0$ have $ds^2 > 0$ and therefore are spacelike separated and cannot be causally connected. Moving straight up in the spacetime diagram is not allowed.

The angle of the lightcone just inside r_S is still 90° , but rather than being squeezed, it is now wide open. As you move in towards $r = 0$, the lightcone closes up, but continues to point toward $r = 0$. Inside $dr/dt = \pm(1 - r_S/r)$. As $r \rightarrow 0$, $dr/dt \rightarrow \pm\infty$, or $dt/dr \rightarrow 0$, and the cone squeezes in pointing towards the center of the hole. So while at first when you are inside the hole there some freedom of movement, in the end you are directed directly at the center.