


## Lecture XIII: Double Well Potential: Tunneling and Instantons

How can phenomena of QM tunneling be described by Feynman path integral?

No semi-classical expansion!

▷ E.g QM transition probability of particle in double well:  $G(a, -a; t) \equiv \langle a | e^{-i\hat{H}t/\hbar} | -a \rangle$



▷ Feynman Path Integral:

$$G(a, -a; t) = \int_{q(0)=-a}^{q(t)=a} Dq \exp \left[ \frac{i}{\hbar} \int_0^t dt' \left( \frac{m}{2} \dot{q}^2 - V(q) \right) \right]$$

Stationary phase analysis: classical e.o.m.  $m\ddot{q} = -\partial_q V$

↳ only singular (high energy) solutions *Switch to alternative formulation...*

▷ Imaginary (Euclidean) time Path Integral: Wick rotation  $t = -i\tau$

N.B. (relative) sign change! “ $V \rightarrow -V$ ”


$$G(a, -a; \tau) = \int_{q(0)=-a}^{q(\tau)=a} Dq \exp \left[ -\frac{1}{\hbar} \int_0^\tau d\tau' \left( \frac{m}{2} \dot{q}^2 + V(q) \right) \right]$$

Saddle-point analysis: classical e.o.m.  $m\ddot{q} = +V'(q)$  in inverted potential!

solutions depend on b.c.

- (1)  $G(a, a; \tau) \rightsquigarrow q_{\text{cl}}(\tau) = a$
- (2)  $G(-a, -a; \tau) \rightsquigarrow q_{\text{cl}}(\tau) = -a$
- (3)  $G(a, -a; \tau) \rightsquigarrow q_{\text{cl}} : \text{rolls from } -a \text{ to } a$

Combined with small fluctuations, (1) and (2) recover propagator for single well



(3) accounts for QM tunneling and is known as an “instanton” (or “kink”)

- ▷ Instanton: classically forbidden trajectory connecting two degenerate minima — i.e. topological, and therefore particle-like

For  $\tau$  large,  $\dot{q}_{cl} \simeq 0$  (*evident*), i.e. “first integral”  $m\dot{q}_{cl}^2/2 - V(q_{cl}) = \epsilon \xrightarrow{\tau \rightarrow \infty} 0$   
*precise value of  $\epsilon$  fixed by b.c. (i.e.  $\tau$ )*


Saddle-point action *(cf. WKB  $\int dq p(q)$ )*

$$S_{inst.} = \int_0^\tau d\tau' \left( \frac{m}{2} \dot{q}_{cl}^2 + V(q_{cl}) \right) \simeq \int_0^\tau d\tau' m \dot{q}_{cl}^2 = \int_{-a}^a dq_{cl} m \dot{q}_{cl} = \int_{-a}^a dq_{cl} (2mV(q_{cl}))^{1/2}$$

Structure of instanton: For  $q \simeq a$ ,  $V(q) = \frac{1}{2}m\omega^2(q - a)^2 + \dots$ , i.e.  $\dot{q}_{cl} \xrightarrow{\tau \rightarrow \infty} \omega(q_{cl} - a)$

$$q_{cl}(\tau) \xrightarrow{\tau \rightarrow \infty} a - e^{-\tau\omega}, \text{ i.e. temporal extension set by } \omega^{-1} \ll \tau$$

Implies existence of approximate saddle-point solutions involving many instantons (and anti-instantons): instanton gas



- ▷ Accounting for fluctuations around n-instanton configuration

$$G(a, \pm a; \tau) \simeq \sum_{n \text{ even/odd}} K^n \int_0^\tau d\tau_1 \int_0^{\tau_1} d\tau_2 \cdots \int_0^{\tau_{n-1}} d\tau_n \overbrace{A_n(\tau_1, \dots, \tau_n)}^{A_{n,cl.} A_{n,qu.}}$$

constant  $K$  set by normalisation

$A_{n,cl.} = e^{-nS_{inst.}/\hbar}$  — ‘classical’ contribution

$A_{n,qu.}$  — quantum fluctuations (imported from single well):  $G_{s.w.}(0, 0; t) \sim \frac{1}{\sqrt{\sin \omega t}}$

$$A_{n,qu.} \sim \prod_i^n \frac{1}{\sqrt{\sin(-i\omega(\tau_{i+1} - \tau_i))}} \sim \prod_i^n e^{-\omega(\tau_{i+1} - \tau_i)/2} \sim e^{-\omega\tau/2}$$

$$G(a, \pm a; \tau) \simeq \sum_{n \text{ even/odd}} K^n e^{-nS_{inst.}/\hbar} e^{-\omega\tau/2} \overbrace{\int_0^\tau d\tau_1 \int_0^{\tau_1} d\tau_2 \cdots \int_0^{\tau_{n-1}} d\tau_n}^{\tau^n/n!}$$

$$= \sum_{n \text{ even/odd}} e^{-\omega\tau/2} \frac{1}{n!} (\tau K e^{-S_{inst.}/\hbar})^n$$

Using  $e^x = \sum_{n=0}^\infty x^n/n!$ ,

N.B. non-perturbative in  $\hbar!$


$$G(a, a; \tau) \simeq C e^{-\omega\tau/2} \cosh(\tau K e^{-S_{inst.}/\hbar})$$

$$G(a, -a; \tau) \simeq C e^{-\omega\tau/2} \sinh(\tau K e^{-S_{inst.}/\hbar})$$

Consistency check: main contribution from

$$\bar{n} = \langle n \rangle \equiv \frac{\sum_n n X^n / n!}{\sum_n X^n / n!} = X = \tau K e^{-S_{\text{inst.}}/\hbar}$$

no. per unit time,  $\bar{n}/\tau$  exponentially small, and indep. of  $\tau$ , i.e. dilute gas



▷ Physical interpretation: For infinite barrier — two independent oscillators, coupling splits degeneracy — symmetric/antisymmetric

$$G(a, \pm a; \tau) \simeq \langle a|S\rangle e^{-\epsilon_S \tau/\hbar} \langle S|\pm a\rangle + \langle a|A\rangle e^{-\epsilon_A \tau/\hbar} \langle A|\pm a\rangle$$

$$|\langle a|S\rangle|^2 = \langle a|S\rangle \langle S|-a\rangle = \frac{C}{2}, \quad |\langle a|A\rangle|^2 = -\langle a|A\rangle \langle A|-a\rangle = \frac{C}{2}$$

Setting:  $\epsilon_{A/S} = \hbar\omega/2 \pm \Delta\epsilon/2$


$$G(a, \pm a; \tau) \simeq \frac{C}{2} \left( e^{-(\hbar\omega - \Delta\epsilon)\tau/2\hbar} \pm e^{-(\hbar\omega + \Delta\epsilon)\tau/2\hbar} \right) = C e^{-\omega\tau/2} \begin{cases} \cosh(\Delta\epsilon\tau/\hbar) \\ \sinh(\Delta\epsilon\tau/\hbar) \end{cases}.$$

▷ Remarks:

(i) Legitimacy? How do (neglected) terms  $O(\hbar^2)$  compare to  $\Delta\epsilon$ ?

In fact, such corrections are bigger but act equally on  $|S\rangle$  and  $|A\rangle$

i.e.  $\Delta\epsilon = \hbar K e^{-S_{\text{inst.}}/\hbar}$  is dominant contribution to splitting



(ii) Unstable States and Bounces: survival probability:  $G(0, 0; t)$ ? No even/odd effect:

$$G(0, 0; \tau) = C e^{-\omega\tau/2} \exp \left[ \tau K e^{-S_{\text{inst.}}/\hbar} \right] \stackrel{\tau=it}{=} C e^{-i\omega t/2} \exp \left[ -\frac{\Gamma}{2} t \right]$$

Decay rate:  $\Gamma \sim |K| e^{-S_{\text{inst.}}/\hbar}$  (i.e.  $K$  imaginary) *N.B. factor of 2*