

Chirikov, Chaos and

KAM \rightarrow an OV

\rightarrow small divisors \rightarrow Chirikov criterion

\rightarrow Development chaos, (Standard Map)

\rightarrow KAM Theorem

\rightarrow Aspects of Chaos.

Shirikov and Chaos and KAM \rightarrow An OV.

- Recall:

- defined action angle variables
- addressed "perturbative integrability"
- \rightarrow defined resonant surface
- \rightarrow noted island formation at resonant surface due resonant perturbations

Some key observations:

• resonant surface defined by $\underline{n} \cdot \underline{\omega} = 0$

- averaging/secular P.T. recovers island with Hess:


$$H = \underbrace{B}_{\sim \frac{\partial \omega}{\partial J}} \frac{AJ^2}{2} - \underbrace{F}_{\sim H_1} \cos \phi$$

- but secular P.T. really only works with one resonance / slow variable in region of averaging


What of multiple perturbed resonances?

c.e.

before:



now



do c/s ends overlap or not?

if no: can consider 2 isolated resonances with un-perturbed tone in between.

- if yes:
- orbits no longer are 'localized' to vicinity of a single resonance
 - integrity of surfaces between resonances is violated \Rightarrow pass 1 to other
 - orbit can pass from resonance-to-resonance \Rightarrow sample volume, not surface.


- \Rightarrow
- \rightarrow tori between 2 resonances are \odot destroyed
- \rightarrow motion fills volume, not surface

\Rightarrow enter chaos! \leftrightarrow {breakdown of integrability for Hamiltonian system.

So

Working criterion for onset of Hamiltonian chaos is Chirikov Island Overlap Criterion

n.b. critical amplitude for onset chaos. $W_I \sim \left(\frac{\partial H_1}{\partial J} \right)^{1/2}$



$\Delta\Omega \equiv$ spacing of resonances

$\Delta W_{1,2} \equiv$ $\frac{1}{2}$ widths of resonances
disturbance (i.e. ends)
at neighboring resonant
frequencies

\Rightarrow $\Delta W_1 + \Delta W_2 = \Delta\Omega$

Chirikov criterion for:

- overlap of i.e. ends at resonances
- destruction of surfaces, between

\Rightarrow
- onset chaos, mixing (in volume
set by resonant helicities)

i.e.:

\rightarrow global criterion (i.e. region not
point).


→ end state of resonance distortions:

C.E.

integrable


integrable

— — —
nested surfaces →




resonant
distortions

— — —
unperturbed
state ($H_1=0$)




integrability breaking down



overlapping
islands
→

non-integrable



destroyed
surfaces,
chaos
mixing

→ Why Believe this story?

- For numerical studies, convenient to work with maps instead ODE's.

∴ enter the Standard Map, c.e.
 (Taylor, Chirikov, early '60's)

unclass
 $\Theta_{n+1} = \Theta_n + \overset{\text{rate}}{p_n} \text{ mod } 2\pi$

$p_{n+1} = p_n + k \overset{\text{perturbation}}{\sin \Theta_{n+1}} \text{ mod } 2\pi$

perturbed winding rate . \uparrow strength. $\Theta \rightarrow$ position
 $p \rightarrow$ momentum

→ 2D, 2 degs freedom

→ phase space is (toroidal) surface
 (1 dim angle, 1 dim radius/action)


→ $\det \begin{vmatrix} \frac{\partial \Theta_{n+1}}{\partial \Theta_n} & \frac{\partial \Theta_{n+1}}{\partial p_n} \\ \frac{\partial p_{n+1}}{\partial \Theta_n} & \frac{\partial p_{n+1}}{\partial p_n} \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ k \cos \Theta_{n+1} & 1 + k \cos \Theta_{n+1} \end{vmatrix}$

area preserving ✓ = 1
 " " " "

→ physicist: kicked rotor

$H(p_0, \Theta, t) = \frac{p_0^2}{2I} + k \cos \Theta \sum_n \delta(t - nT)$

so



$\sigma = 0$
 vertical impulsive force, at T period

$$\frac{dP_0}{dt} = K \cos \theta \sum_n \delta(t - nT)$$

$$\frac{d\theta}{dt} = P_0 / I$$

integrating and $T/I = 1 \Rightarrow$ standard map,
 no lab animal for studies of stochasticity.

\Rightarrow so:

① $\Delta p_{max} = 2K^{1/2}$ (dyn. system)
 $(m=1)$
 \rightarrow island size $\sim \sqrt{K}$

$$\Delta_{res} = 2\pi$$

$\Rightarrow K_{crit} \approx 2.47$ \rightarrow destruction of surfaces for $m=1$ overlap

② if interaction bet. ~~period 1, period 2~~ ~~period 1, period 2~~ ~~period 1, period 2~~

$\Rightarrow K_{crit} \approx 1.46$ \rightarrow ~~overlap~~ ~~overlap~~ ~~overlap~~ overlap criterion.

\therefore if exame off fig 7.3, pg. 275 see!

a.) $K = .5$

- surfaces preserved, except for stochastic layer near separatrices.
- ~~period 1~~, ~~period 2~~ islands clearly preserved

b.) $K = 1$

- onset stochasticity
- surface between ^{1, 2} islands destroyed _{period}

c.) $K = 2.5$

- stochasticity, strong
- ~~period~~ unstable \rightarrow islands only residues

d.) $K = 4$

- ~~period 1~~ going unstable.

Point:

- Chirikov is slight under-estimate of stochasticity onset




Figure 7.13 Plots of p modulo 2π for four values of K : (a) $K = 0.5$; (b) $K = 1.0$; (c) $K = 2.5$; (d) $K = 4.0$. (This figure courtesy of Y. Du.)




Figure 7.14 Two KAM curves bounding an annular region.

KAM curves (as, for example, in the island structures surrounding elliptic orbits), these chaotic orbits are necessarily restricted to lie between the bounding KAM curves. (As we shall discuss later, this picture is fundamentally different for systems of higher dimensionality.)

- ignores secondary islands, stochastic layers etc

⇒ Resonance overlap leads to breakdown of integrability, destruction of tori, onset of chaos, and mixing.

⇒ Prototype of mechanism for onset of deterministic, Hamiltonian chaos.

Now:

- story presented is 'tip of very large iceberg'

→ see Lichtenberg & Leiberman, and literature for details

- non-Hamiltonian chaos is fundamentally different ⇒ attractors.

Some key Questions:

① - is there a theorem? ⇒ can we prove the story? → KAM theorem

② how characterize chaos? \rightarrow dynamical entropy?

③ how calculate in chaotic regime \rightarrow stat mech.

Very Abbreviated story:

\rightarrow KAM Theorem (Kolmogorov, Arnold, Moser)

how resolve the "small divisor" problem, rigorously?

i.e. can we integrate the system perturbatively?!

Thm:

pert.
 \downarrow

For $H = H_0 + \epsilon H_1$, if ϵ is small enough, then for almost all frequencies $\underline{\omega}^*$, there exists an invariant torus $T(\underline{\omega}^*)$ of perturbed system that $T(\underline{\omega}^*)$ is close to $T_0(\underline{\omega}^*)$.

$T_0 \rightarrow$ torus at invariant surface, unperturbed
 $T \rightarrow$ perturbed surface.

Translation :

non-resonant

" A sufficiently irrational torus can survive a sufficiently weak perturbation "

Some clarification :

↳ threshold for chaos

- What is an irrational or non-resonant torus ?

$$| \underline{m} \cdot \underline{\omega} | > K(\underline{\omega}) |m|^{-N+1}$$

N dots

$$|m| = |m_1| + |m_2| + \dots + |m_N|$$

$K(\underline{\omega})$ indep. \underline{m}

↳ set of N dim. vectors not satisfying above \Leftrightarrow set $\mu \rightarrow 0$

↳ irrational tori are 'common', rational tori are 'unusual'

- What does "survive" mean ?

torus of perturbed system has frequency

$$\underline{\omega}(\epsilon) = K(\epsilon) \underline{\omega}_0 \quad \text{s/t } K(\epsilon) \rightarrow 1 \text{ as } \epsilon \rightarrow 0$$

→ smooth distortion.

- KAM theorem says for small ϵ , the perturbed system's phase volume not occupied by surviving tori is small, and $\rightarrow \epsilon$ with ϵ .

Key Mathematical Points:

- rationals are set of $n=0$ in numbers.
- number theoretic arguments for 'sufficiently non-resonant'
- exponential decay of $H_1(m) \sim e^{-|m|}$

Implication:

\Rightarrow Resonant, rational tori are key to Hamiltonian chaos, integrability breakdown.

N.B: ^① Poincaré - Birkhoff theorem provides rigorous underpinning for resonant tori and their distortion by perturbations.


^② Can improve on Chirikov by Greene calculation of when KAM torus penetrated by perturbation.

→ Characterizing Chaos


① → how locally diagnose chaos?

② → how characterize strength of chaos.

① Chaos → exponential divergence of neighboring trajectories
 ⇒ 'instability' - sensitivity to i.c.
 - Butterfly effect



$$l \sim l_0 e^{ht}$$



$h \equiv$ Lyapunov exponent

stretching +
volume preservation

N.B.:

→ Lyapunov exponent \Leftrightarrow direction

→ chaos \Leftrightarrow stretching

→ volume preservation (Hamiltonian)

$$\sum_i h_i = 0$$

dir. must have negative h
 corresponding to direction
 of shrinking.

→ Lyapunov exponent is local, -pt./ball
Chirikov is global.


② Strength → Metric or K - β^k
(Kolmogorov-Sinai) Entropy

K - β^k Entropy: Rate of creation of information
as system evolves

→ phase space is partitioned
c.e. → resolution scale
→ coarse graining

→ rate of info creation ↔ rate at which
two sub-grid. sep. orbits become separated
by $>$ grid scale → need keep track of
these as distinct orbits.

c.e.



① }
②' } when
"info
created"

② has greater metric entropy.

c.e. } they pt. is info created when
separation exceeds the partition scale.

$$\Rightarrow \left[h(\alpha) = \sum_{h_i > 0} h_i \right], \quad \text{for Hamiltonian systems}$$

\downarrow
 K-S entropy dynamical entropy

∴ positive Lyapunov exponents characterize dynamical entropy of system.

⑤ How calculate in chaotic regime?

→ Stat. Mech. ↓ → i.e. pdf

→ in particular, mean field theory
i.e. coarse grained probability most practical.

$$\Theta_{n+1} = \Theta_n + \Delta\Theta_n \quad (\text{relax mode})$$

$$P_{n+1} = P_n + k \sin \Theta_{n+1}$$

if Θ random (chaos!) ⇒ $k \gg$ Kurt per stoch. assumed

$$P_{n+1} - P_n = \Delta P_n = k \sin \Theta_{n+1}$$

$$\langle \Delta P_n^2 \rangle = k^2 \langle \sin^2 \Theta_{n+1} \rangle \approx \frac{k^2}{2} \quad \langle \rangle = \text{avg over steps (time)}$$

i.e. $\langle \Delta P_n^2 \rangle$ diffusive \sim random walk

$$\langle p^2/2 \rangle \simeq Dn. \quad (7.44)$$

The quasilinear result (7.42) is valid for very large K . For moderately large, but not very large, values of K , neglected correlation effects can significantly alter the diffusion coefficient from the quasilinear value. These effects have been analytically calculated by Rechester and White (1980) (see also Rechester *et al.* (1981), Karney *et al.* (1981) and Carey *et al.* (1981)). Figure 7.17 shows a plot of the diffusion coefficient D normalized to D_{QL} as a function of K from the paper of Rechester and White. The solid curve is their theory, and the dots are obtained by numerically calculating the spreading of a cloud of points and obtaining D from Eq. (7.44). Note the decaying oscillations about the quasilinear value as K increases.⁷

7.3.4 Other examples

So far in this section we have dealt exclusively with the standard map. We now discuss some other examples, also reducible to two-dimensional maps, where similar phenomena are observed.

We first consider a time-independent two-degree-of-freedom system investigated by Schmidt and Chen (1994). This system, depicted in Figure 7.18, consists of two masses, a large mass M connected to a linearly behaving spring of spring constant k_s and a small mass m which elastically




Figure 7.17 D/D_{QL} versus K for the standard map (Rechester and White, 1980).

$$\frac{dx}{dt} = v$$

$$\frac{dv}{dt} = \epsilon \sin x \quad (*)$$

$$\frac{\sin[\pi(2N+1)]}{\sin \pi H}$$

Now, as θ random, φ diffuses
so!

$$\langle (\Delta p_n)^2 \rangle = 2 D n$$

\downarrow
diffusion coeff
in p (momentum)

\Rightarrow

$$D = \frac{1}{4} v^2$$

\rightarrow quasi-linear
diffusion coefficient
 \rightarrow QK works rather well \rightarrow
see off p. 281

For systematic: Fokker-Planck Theory

$f \rightarrow$ dist.

schematically

$$f(t+\tau, p) = \int d\Delta p f(t, p-\Delta p) T(\Delta p, \tau)$$

\rightarrow Chapman-Kolmogorov Eqn.

\rightarrow T is transition probability, for step
 Δp in \mathcal{X}

\rightarrow expansion \rightarrow Fokker-Planck Theory / Eqn.
and can calculate divergence.