

Lectures 17: Jackknife sampling error propagation for nonlinear functions of fit parameters

Jackknife sampling

Introduction

The bootstrap method is not always the best one. One main reason is that the bootstrap samples are generated from \hat{F} and not from F . **Can we find samples/resamples exactly generated from F ?**

- If we look for samples of size n , then the answer is **no!**
- If we look for samples of size m ($m < n$), then we can indeed find (re)samples of size m exactly generated from F simply by looking at different subsets of our original sample \mathbf{x} !

Looking at different subsets of our original sample amounts to sampling without replacement from observations x_1, \dots, x_n to get (re)samples (now called **subsamples**) of size m . This leads us to subsampling and the jackknife.

Jackknife sampling

Jackknife

- The jackknife has been proposed by Quenouille in mid 1950's.
- In fact, the jackknife predates the bootstrap.
- The jackknife (with $m = n - 1$) is less computer-intensive than the bootstrap.
- *Jackknife* describes a swiss penknife, easy to carry around. By analogy, Tukey (1958) coined the term in statistics as a general approach for testing hypotheses and calculating confidence intervals.

Jackknife sampling

Jackknife samples

Definition

The **Jackknife samples** are computed by leaving out one observation x_i from $\mathbf{x} = (x_1, x_2, \dots, x_n)$ at a time:

$$\mathbf{x}_{(i)} = (x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$$

- The dimension of the jackknife sample $\mathbf{x}_{(i)}$ is $m = n - 1$
- n different Jackknife samples : $\{\mathbf{x}_{(i)}\}_{i=1 \dots n}$.
- No sampling method needed to compute the n jackknife samples.

Jackknife sampling

Jackknife replications

Definition

The i th **jackknife replication** $\hat{\theta}_{(i)}$ of the statistic $\hat{\theta} = s(\mathbf{x})$ is:

$$\hat{\theta}_{(i)} = s(\mathbf{x}_{(i)}), \quad \forall i = 1, \dots, n$$

Jackknife replication of the mean

$$\begin{aligned} s(\mathbf{x}_{(i)}) &= \frac{1}{n-1} \sum_{j \neq i} x_j \\ &= \frac{(n\bar{x} - x_i)}{n-1} \\ &= \bar{x}_{(i)} \end{aligned}$$

Jackknife sampling

Jackknife estimation of the standard error

- 1 Compute the n jackknife subsamples $\mathbf{x}_{(1)}, \dots, \mathbf{x}_{(n)}$ from \mathbf{x} .
- 2 Evaluate the n jackknife replications $\hat{\theta}_{(i)} = s(\mathbf{x}_{(i)})$.
- 3 The **jackknife estimate of the standard error** is defined by:

$$\hat{se}_{jack} = \left[\frac{n-1}{n} \sum_{i=1}^n (\hat{\theta}_{(\cdot)} - \hat{\theta}_{(i)})^2 \right]^{1/2}$$

where $\hat{\theta}_{(\cdot)} = \frac{1}{n} \sum_{i=1}^n \hat{\theta}_{(i)}$.

Jackknife sampling

Jackknife estimation of the standard error of the mean

For $\hat{\theta} = \bar{x}$, it is easy to show that:

$$\begin{cases} \bar{x}_{(i)} = \frac{n\bar{x} - x_i}{n-1} \\ \bar{x}(\cdot) = \frac{1}{n} \sum_{i=1}^n \bar{x}_{(i)} = \bar{x} \end{cases}$$

Therefore:

$$\begin{aligned} \hat{se}_{jack} &= \left\{ \sum_{i=1}^n \frac{(x_i - \bar{x})^2}{(n-1)n} \right\}^{1/2} \\ &= \frac{\bar{\sigma}}{\sqrt{n}} \end{aligned}$$

where $\bar{\sigma}$ is the unbiased variance.

Jackknife sampling

Jackknife estimation of the standard error

- The factor $\frac{n-1}{n}$ is much larger than $\frac{1}{B-1}$ used in bootstrap.
- Intuitively this inflation factor is needed because jackknife deviation $(\hat{\theta}_{(i)} - \hat{\theta}_{(\cdot)})^2$ tend to be smaller than the bootstrap $(\hat{\theta}^*(b) - \hat{\theta}^*(\cdot))^2$ (the jackknife sample is more similar to the original data \mathbf{x} than the bootstrap).
- In fact, the factor $\frac{n-1}{n}$ is derived by considering the special case $\hat{\theta} = \bar{x}$ (somewhat arbitrary convention).

Jackknife sampling

Comparison of Jackknife and Bootstrap on an example

Example A: $\hat{\theta} = \bar{x}$

$F(x) = 0.2 \mathcal{N}(\mu=1, \sigma=2) + 0.8 \mathcal{N}(\mu=6, \sigma=1) \rightsquigarrow \mathbf{x} = (x_1, \dots, x_{100})$.

- Bootstrap standard error and bias w.r.t. the number B of bootstrap samples:

B	10	20	50	100	500	1000	10000
\widehat{se}_B	0.1386	0.2188	0.2245	0.2142	0.2248	0.2212	0.2187
\widehat{Bias}_B	0.0617	-0.0419	0.0274	-0.0087	-0.0025	0.0064	0.0025

- Jackknife: $\widehat{se}_{jack} = 0.2207$ and $\widehat{Bias}_{jack} = 0$
- Using textbook formulas: $se_{\hat{f}} = \frac{\hat{\sigma}}{\sqrt{n}} = 0.2196$ ($\frac{\bar{\sigma}}{\sqrt{n}} = 0.2207$).

Jackknife sampling

Jackknife estimation of the bias

- 1 Compute the n jackknife subsamples $\mathbf{x}_{(1)}, \dots, \mathbf{x}_{(n)}$ from \mathbf{x} .
- 2 Evaluate the n jackknife replications $\hat{\theta}_{(i)} = s(\mathbf{x}_{(i)})$.
- 3 The **jackknife estimation of the bias** is defined as:

$$\widehat{\text{Bias}}_{jack} = (n - 1)(\hat{\theta}_{(\cdot)} - \hat{\theta})$$

where $\hat{\theta}_{(\cdot)} = \frac{1}{n} \sum_{i=1}^n \hat{\theta}_{(i)}$.

Jackknife sampling

Jackknife estimation of the bias

- Note the inflation factor $(n - 1)$ (compared to the bootstrap bias estimate).
- $\hat{\theta} = \bar{x}$ is unbiased so the correspondence is done considering the plug-in estimate of the variance $\hat{\sigma}^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$.
- The jackknife estimate of the bias for the plug-in estimate of the variance is then:

$$\widehat{\text{Bias}}_{jack} = \frac{-\hat{\sigma}^2}{n}$$

Jackknife sampling

Histogram of the replications

Example A




Figure: Histograms of the bootstrap replications $\{\hat{\theta}^*(b)\}_{b \in \{1, \dots, B=1000\}}$ (left), and the jackknife replications $\{\hat{\theta}_{(i)}\}_{i \in \{1, \dots, n=100\}}$ (right).

Jackknife sampling

Histogram of the replications

Example A




Figure: Histograms of the bootstrap replications $\{\hat{\theta}^*(b)\}_{b \in \{1, \dots, B=1000\}}$ (left), and the inflated jackknife replications $\{\sqrt{n-1}(\hat{\theta}_{(i)} - \hat{\theta}_{(\cdot)}) + \hat{\theta}_{(\cdot)}\}_{i \in \{1, \dots, n=100\}}$ (right).

Jackknife sampling

Relationship between jackknife and bootstrap

- When n is small, it is easier (faster) to compute the n jackknife replications.
- However the jackknife uses less information (less samples) than the bootstrap.
- In fact, the jackknife is an approximation to the bootstrap!

Jackknife sampling

Relationship between jackknife and bootstrap

- Considering a linear statistic :

$$\begin{aligned}\hat{\theta} &= s(\mathbf{x}) = \mu + \frac{1}{n} \sum_{i=1}^n \alpha(x_i) \\ &= \mu + \frac{1}{n} \sum_{i=1}^n \alpha_i\end{aligned}$$

Mean $\hat{\theta} = \bar{x}$

The mean is linear $\mu = 0$ and $\alpha(x_i) = \alpha_i = x_i, \quad \forall i \in \{1, \dots, n\}$.

- There is no loss of information in using the jackknife to compute the standard error (compared to the bootstrap) for a linear statistic. Indeed the knowledge of the n jackknife replications $\{\hat{\theta}_{(i)}\}$, gives the value of $\hat{\theta}$ for any bootstrap data set.
- For non-linear statistics, the jackknife makes a linear approximation to the bootstrap for the standard error.

Jackknife sampling

Relationship between jackknife and bootstrap

- Considering a quadratic statistic

$$\hat{\theta} = s(\mathbf{x}) = \mu + \frac{1}{n} \sum_{i=1}^n \alpha(x_i) + \frac{1}{n^2} \beta(x_i, x_j)$$

Variance $\hat{\theta} = \hat{\sigma}^2$

$\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$ is a quadratic statistic.

- Again the knowledge of the n jackknife replications $\{s(\hat{\theta}_{(i)})\}$, gives the value of $\hat{\theta}$ for any bootstrap data set. The jackknife and bootstrap estimates of the bias agree for quadratic statistics.

Jackknife sampling

Relationship between jackknife and bootstrap

The Law school example: $\hat{\theta} = \widehat{\text{corr}}(\mathbf{x}, \mathbf{y})$.

The correlation is a non linear statistic.

- From $B=3200$ bootstrap replications, $\hat{se}_{B=3200} = 0.132$.
- From $n = 15$ jackknife replications, $\hat{se}_{jack} = 0.1425$.
- Textbook formula: $se_{\hat{f}} = (1 - \widehat{\text{corr}}^2) / \sqrt{n - 3} = 0.1147$

Jackknife sampling

Delete-d Jackknife samples

Definition

The **delete-d Jackknife** subsamples are computed by leaving out d observations from \mathbf{x} at a time.

- The dimension of the subsample is $n - d$.
- The number of possible subsamples now rises $\binom{n}{d} = \frac{n!}{d!(n-d)!}$.
- Choice: $\sqrt{n} < d < n - 1$

Jackknife sampling

Delete-d jackknife

- 1 Compute all $\binom{n}{d}$ d-jackknife subsamples $\mathbf{x}_{(1)}, \dots, \mathbf{x}_{(n)}$ from \mathbf{x} .
- 2 Evaluate the jackknife replications $\hat{\theta}_{(i)} = s(\mathbf{x}_{(i)})$.
- 3 Estimation of the standard error (when $n = r \cdot d$):

$$\hat{\text{se}}_{d\text{-jack}} = \left\{ \frac{r}{\binom{n}{d}} \sum_i (\hat{\theta}_{(i)} - \hat{\theta}(\cdot))^2 \right\}^{1/2}$$

$$\text{where } \hat{\theta}(\cdot) = \frac{\sum_i \hat{\theta}_{(i)}}{\binom{n}{d}}.$$

Jackknife sampling

Summary

- Bias and standard error estimates have been introduced using jackknife replications.
- The Jackknife standard error estimate is a linear approximation of the bootstrap standard error.
- The Jackknife bias estimate is a quadratic approximation of the bootstrap bias.
- Using smaller subsamples (delete-d jackknife) can improve for non-smooth statistics such as the median.

Jackknife sampling Matlab code

```
%% Jackknife Resampling
%

% Copyright 2015 The MathWorks, Inc.

%%
% Similar to the bootstrap is the jackknife, which uses resampling to
% estimate the bias of a sample statistic. Sometimes it is also used to
% estimate standard error of the sample statistic. The jackknife is
% implemented by the Statistics and Machine Learning Toolbox(TM) function
% |jackknife|.

%%
% The jackknife resamples systematically, rather than at random as the
% bootstrap does. For a sample with |n| points, the jackknife computes
% sample statistics on |n| separate samples of size |n|-1. Each sample is
% the original data with a single observation omitted.

%%
% In the bootstrap example, you measured the uncertainty in estimating the
% correlation coefficient. You can use the jackknife to estimate the bias,
% which is the tendency of the sample correlation to over-estimate or
% under-estimate the true, unknown correlation. First compute the sample
% correlation on the data.
load lawdata
rho_hat = corr(lsat,gpa)

%%
% Next compute the correlations for jackknife samples, and compute their
% mean.
rng default; % For reproducibility
jackrho = jackknife(@corr,lsat,gpa);
meanrho = mean(jackrho)

%%
% Now compute an estimate of the bias.
n = length(lsat);
biasrho = (n-1) * (meanrho-rho_hat)

%%
% The sample correlation probably underestimates the true correlation by
% about this amount.
```