


# Langmuir Turbulence I

→ Disparate Scale Interaction  
(see notes)

→ Deriving Zakharov Eqs.  
critically



Observe:  $\left\{ \begin{array}{l} \omega^2 = \omega_p^2 (1 + \kappa k^2 \lambda_{D0}^2) \\ \omega = k c_s \end{array} \right.$  Langmuir (Density)

→  $\left\{ \begin{array}{l} \omega = k c_s \\ \rightarrow 0 \end{array} \right.$

so basic interaction must be  $L + L \rightarrow \delta n$

i.e. interaction of Langmuir wave energy gas with low frequency perturbations.


⇒ As energy field  $\leftrightarrow \delta n$  evolution, must be envelope.



Now:  $\omega^2 = \omega_{pe}^2 (1 + \alpha k^2 \lambda_{D0}^2)$

$$(\omega_0 + i\gamma)^2 = \omega_{pe}^2 \left(1 + \frac{\delta n}{n_0}\right) \left(1 + \alpha (k + \frac{\delta k}{\lambda})^2 \lambda_{D0}^2\right)$$

$\downarrow$  slow evolution       $\downarrow$  modulation by density perturbation       $\downarrow$  envelope wave #



⇒

$$\omega_0^2 + 2i\gamma\omega_0 + (i\gamma)^2 = \omega_{pe}^2 + \frac{\delta n}{n_0} \omega_{pe}^2 + \alpha k^2 v_{th}^2 + \alpha v_{th}^2 (2k \cdot \underline{e}) + \alpha \epsilon^2 v_{th}^2$$

$$2i\gamma\omega_0 = \frac{\delta n}{n_0} \omega_{pe}^2 + \alpha (2k \cdot \underline{e}) v_{th}^2 + \alpha \epsilon^2 v_{th}^2$$

$$\omega_0 \approx \omega_{pe}$$

$$i\gamma\omega_0 = \frac{\omega_{pe}^2}{2} \frac{\delta n}{n_0} + \alpha (k \cdot \underline{e}) v_{th}^2 + \frac{\alpha}{2} \epsilon^2 v_{th}^2$$

$$i\gamma = \frac{\omega_{pe}}{2} \frac{\delta n}{n_0} + \frac{\alpha k v_{th}^2 \cdot \underline{e}}{\omega_0} + \alpha \frac{\epsilon^2 v_{th}^2}{\omega_0}$$



$$\omega = \omega_{p0} (1 + \alpha k^2 \lambda_D^2)^{1/2}$$

$$\frac{d\omega}{dk} = \omega_{p0} (\alpha k \lambda_D^2) / (1 + \alpha k^2 \lambda_D^2)^{1/2}$$

$$= \alpha k \frac{v_{Te}^2}{\omega_{p0}} = v_{gr}$$

$$\text{so } \alpha \frac{k v_{Te}^2}{\omega_0} \cdot \underline{z} = \underline{z} \cdot \underline{v}_{gr}$$

$$\text{Now } \psi = i \frac{\partial}{\partial t}$$

so if shift to co-moving frame

$$\underline{d.e} \quad \underline{x} \rightarrow \underline{x} - \underline{v}_{gr} t$$

then can eliminate  $\underline{z} \cdot \underline{v}_{gr}$  term.

∴

$$i \gamma = \frac{\omega_{p0}^2}{2} \frac{dN}{N_0} + \alpha \frac{z^2 v_{Te}^2}{\omega_0}$$

$$\Rightarrow \text{if } \underline{E} = \sum \underbrace{(\underline{A}_j(t))}_{\text{envelope}} e^{i(\underline{k} \cdot \underline{x} - \omega_j t)} \underbrace{e^{i(\underline{k} \cdot \underline{x} - \omega_j t)}}_{\text{carrier}}$$



$$i \frac{\partial \Sigma}{\partial t} = \frac{\omega_{pe}^2}{2} \frac{d\eta}{\eta} \Sigma - \frac{\omega_{pe}^2}{\omega_0} \nabla^2 \Sigma$$

$\downarrow$  refraction                       $\downarrow$  diffraction

~~Factor out~~

Can re-write?

$$i \omega_0 \frac{\partial \Sigma}{\partial t} = \frac{\omega_{pe}^2}{2} \frac{d\eta}{\eta_0} \Sigma - \omega_{pe}^2 \nabla^2 \Sigma$$

Now, for  $d\eta$

$$\frac{\partial \eta}{\partial t} + \nabla \cdot (n\mathbf{v}) = 0$$

$$n m_i \frac{d\mathbf{v}}{dt} = -\nabla P$$

$$P = P_{Th} + P_{rad} \approx c_s^2 m_i d\eta + \frac{|\mathbf{E}|^2}{8\pi}$$

$\downarrow$   
 ponderomotive  
 pressure



$$\partial_t \delta n + n_0 \underline{\nabla} \cdot \underline{\tilde{v}} = 0$$

$$n m_i \frac{\partial \underline{v}}{\partial t} = -\underline{\nabla} \left( c_s^2 m_i \delta n + \frac{|\underline{E}|^2}{8\pi} \right)$$

$$\frac{\partial \underline{v}}{\partial t} = -\underline{\nabla} \left( c_s^2 \frac{\delta n}{n_0} + \frac{|\underline{E}|^2}{8\pi n m_i} \right)$$

so

$$\partial_t (\underline{\nabla} \cdot \underline{\tilde{v}}) = -\nabla^2 \left( c_s^2 \frac{\delta n}{n_0} + \frac{|\underline{E}|^2}{8\pi n m_i} \right)$$

$$\partial_t^2 (\delta n / n_0) + \partial_t (\underline{\nabla} \cdot \underline{\tilde{v}}) = 0$$

so

$$\partial_t^2 \frac{\delta n}{n_0} - c_s^2 \nabla^2 \frac{\delta n}{n_0} = \nabla^2 \left( \frac{|\underline{E}|^2}{8\pi n m_i} \right)$$

so have Z-egns.



$$i\omega_0 \frac{\partial \epsilon}{\partial t} = \frac{\omega_{pe0}^2}{2} \frac{\delta n}{n_0} \epsilon - \alpha v_{th0}^2 \nabla^2 \epsilon$$

$$\frac{\partial^2}{\partial t^2} \frac{\delta n}{n_0} - c_s^2 \nabla^2 \frac{\delta n}{n_0} = \nabla^2 \left( \frac{|\epsilon|^2}{8\pi n m_0} \right)$$


For  $T \gg L/c_s$

$$-c_s^2 \nabla^2 \frac{\delta n}{n_0} = \nabla^2 \left( \frac{|\epsilon|^2}{8\pi n m_0} \right)$$

cavity  
↓

$$\therefore \frac{\delta n}{n_0} = - \frac{|\epsilon|^2}{8\pi n T}$$

State of (thermal + ponderomotive) Press  $\approx 0$



so plug into  $\epsilon$  eqn,



$$i\omega_0 \frac{\partial \epsilon}{\partial t} = -\frac{\omega_p^2}{2} \left( \frac{|\epsilon|^2}{8\pi\pi} \right) \epsilon - \alpha v_{th}^2 \nabla^2 \epsilon$$

i.e.

refraction

potential

$$i\omega_0 \frac{\partial \epsilon}{\partial t} = -\alpha^2 v_{th}^2 \nabla^2 \epsilon - \frac{\omega_p^2}{2} \left( \frac{|\epsilon|^2}{8\pi\pi} \right) \epsilon$$

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V \cdot \psi$$

$\Rightarrow$  NLS !  $V < 0$   
(attractive)

Also similar to self-focusing problem.

$\rightarrow$  Now, can simplify description to

$\rightarrow$  acoustic wave

$\rightarrow$  adiabatic (Action) eqn.



8.

## NOTES ON MODULATIONAL INTERACTION OF SECONDARY STRUCTURES, PART I

LECTURE BY P.H. DIAMOND AND NOTES BY RONGJIE HONG

### 1. SHORT INTRODUCTION ON DISPARATE SCALE INTERACTION

In this lecture, we will introduce one generic class of nonlinear processes called disparate scale interaction. Plasma turbulence itself has several explicit distinct characteristic length scales. For instance, ion gyroradius  $\rho_i$  and electron gyroradius  $\rho_e$  in magnetized plasmas. The Debye length for collective oscillation, and skin depth for magnetic perturbation.

One reason for the disparate scales is that  $m_e \ll m_i$ , thus plasmas have different fluctuation properties, e.g. plasma waves vs. ion-acoustic waves.

Nonlinear dynamics: unstable modes couple stable modes with common scale length. For example, drift waves are unstable when  $k_{\parallel} \ll k_{\theta}$ , then nonlinear interaction within like-scale will increase  $k_{\parallel}$ , allowing energy transfer to strongly damped modes.

Disparate scale interaction is in contrast to Kolmogorov cascade in neutral fluids. In cascade, the kinetic energy is transferred from large scale  $L$  to micro-scale  $l_d$  where the kinetic energy dissipated. And there is no preferred scale between  $L$  and  $l_d$ .

High  $\omega$  high  $k$  fluctuation (small)  $\rightarrow$  low  $\omega$  low  $k$  structure (large) by effective stress, and couple energy to large scale; Low  $\omega$  low  $k$  structure (large)  $\rightarrow$  high  $\omega$  high  $k$  fluctuation (small) by refraction or strain field, as shown in Fig. 1.1

Disparate scale interaction is a typical process for the generation of large scale structure [Diamond et al. PPCF 2005]. Formation of large scale by turbulence is similar to 'inverse cascade' in fluid dynamics. But in disparate scale interaction, the energy transfer directly to long-wavelength structure from small scales, while




FIGURE 1.1. Interaction between small-scale fluctuations and large-scale ones.

Key leverage: <sup>1</sup> Adiabatic Theory



in 'inverse cascade' case, the transfer occurs through a sequence of intermediate scales.

The disparate scale interactions has many examples. The simplest one is the interaction between Langmuir turbulence (plasma waves or plasmons) and ion-acoustic waves (phonons). In this case, the plasma waves form ponderomotive pressure field to ion-acoustic waves, and the density perturbation of ion-acoustic waves refracts plasma waves, so that the modulation of plasma waves grows.

The second example is the drift wave-zonal flow interaction in toroidal plasmas. In this case, small scale drift wave turbulence induces transport of momentum (Reynolds stress or vorticity flux), which amplifies the zonal flow shear. On the other hand, zonal flow shears stretch and tilt the drift wave packet. The coupling leads to an energy transfer from drift wave turbulence to zonal flows, which is an important nonlinear process for confinement of toroidal magnetized plasmas.

In the following contents, we will discuss these two examples.

## 2. WAVE KINETIC THEORY FOR LANGMUIR TURBULENCE

In general situations, plasma waves are excited as Langmuir turbulence, and the ion-acoustic waves may also be a broad-band spectrum. The evolution of envelope of Langmuir turbulence then may be comparable to the ion-acoustic speed. The coupling between Langmuir turbulence and ion-acoustic waves is studied using quasi-particle approach here.

The Langmuir turbulence field is characterized by action density  $N(\mathbf{k}, \mathbf{x}, t)$ , i.e population density of waves.

$$N = \frac{E_k}{\omega_k}, \quad E_k = \frac{\partial}{\partial \omega} (\omega \epsilon) |_{\omega_k} \frac{|\tilde{\mathbf{E}}_k|^2}{8\pi}$$

where  $E_k$  is energy density,  $\tilde{\mathbf{E}}_k$  is the electric field of plasma wave at wave number  $\mathbf{k}$ .

The ion-acoustic waves are described by the density and velocity perturbations  $\tilde{n}$  and  $\tilde{\mathbf{V}}$ , both of which vary slowly compared to  $\mathbf{k}$  and  $\omega_k$  of Langmuir turbulence.

Under the condition of scale separation, the  $N(\mathbf{k}, \mathbf{x}, t)$  is conserved along the trajectory. The wave kinetics equation for Langmuir turbulence under the influence of ion-acoustic waves is written as,

$$(2.1) \quad \frac{dN}{dt} = \frac{\partial N}{\partial t} + (\mathbf{v}_g + \tilde{\mathbf{V}}) \cdot \frac{\partial N}{\partial \mathbf{x}} - \frac{\partial}{\partial \mathbf{x}} (\omega_k + \mathbf{k} \cdot \tilde{\mathbf{V}}) \cdot \frac{\partial N}{\partial t} = 0$$

the trajectories determined by eikonal equations,

$$\frac{d\mathbf{x}}{dt} = \frac{\partial \omega_k}{\partial \mathbf{x}} + \tilde{\mathbf{V}}, \quad \frac{d\mathbf{k}}{dt} = -\frac{\partial}{\partial \mathbf{x}} (\omega_k + \mathbf{k} \cdot \tilde{\mathbf{V}})$$

In the presence of long-scale perturbations, wave frequency is modified,

$$\omega_k = \omega_{k0} + \tilde{\omega}_k$$

$\omega_{k0}$  is given in the absence of acoustic waves, and the unperturbed orbit of quasi-particles is then,

$$\frac{d\mathbf{x}_0}{dt} = \frac{\partial \omega_{k0}}{\partial \mathbf{x}} = \mathbf{v}_g, \quad \frac{d\mathbf{k}_0}{dt} = -\frac{\partial \omega_{k0}}{\partial \mathbf{x}}$$



**2.1. Evolution of the Langmuir wave action density.** Set the action density of plasmons  $N = \langle N \rangle + \tilde{N}$ . We analyze the case that the Langmuir turbulence is homogeneous in unperturbed state, and the Doppler shift is smaller than effects of modulation of refraction, then the dispersion relation for plasma waves is

$$\omega^2 = \omega_{p0}^2 \left(1 + \frac{\tilde{n}}{n_0}\right)$$

2.1.1. *Evolution of Langmuir wave energy density.* In interacting with acoustic waves, the action density  $N$  is conserved, then the change of energy density of plasma waves is

$$\frac{d}{dt} E_k = \frac{d}{dt} (\omega_k N) = N \frac{d}{dt} \omega_k$$

Noting the relations,

$$\frac{d}{dt} \omega_k = \frac{\partial \omega_k}{\partial \mathbf{k}} \cdot \frac{d\mathbf{k}}{dt} = \mathbf{v}_g \cdot \left(-\frac{\partial \omega_k}{\partial \mathbf{x}}\right) = -\mathbf{v}_g \cdot \frac{\partial}{\partial \mathbf{x}} \left(\frac{\omega_{p0}}{2} \frac{\tilde{n}}{n_0}\right)$$

where we use

$$\frac{d\mathbf{k}}{dt} = -\frac{\partial \omega_k}{\partial \mathbf{x}} = -\frac{\omega_{p0}}{2n_0} \frac{\partial \tilde{n}}{\partial \mathbf{x}}$$

then one has,

$$\frac{dE_k}{dt} = -\frac{N\omega_{p0}}{2n_0} \mathbf{v}_g \cdot \frac{\partial \tilde{n}}{\partial \mathbf{x}}$$

Putting  $N = \langle N \rangle + \tilde{N}$  into this relation, and 1st order terms vanishes in long time average, 2nd order terms survive,

$$\frac{d}{dt} \langle E_k \rangle = -\frac{\omega_{p0}}{2n_0} \mathbf{v}_g \cdot \langle \tilde{N} \frac{\partial \tilde{n}}{\partial \mathbf{x}} \rangle$$

This relation indicates that the change of plasma waves energy density, which is transferred to ion-acoustic waves, is given by the correlation  $\langle \tilde{N} \frac{\partial \tilde{n}}{\partial \mathbf{x}} \rangle$ .

2.1.2. *Wave kinetic equation of Langmuir action density.* Putting  $N = \langle N \rangle + \tilde{N}$  into equation 2.1, yields the response of  $\langle N \rangle$  and  $\tilde{N}$  to ion-acoustic waves,

$$(2.2) \quad \begin{aligned} \frac{\partial \tilde{N}}{\partial t} + \mathbf{v}_g \cdot \frac{\partial \tilde{N}}{\partial \mathbf{x}} - \frac{\partial \omega_{k0}}{\partial \mathbf{x}} \cdot \frac{\partial \tilde{N}}{\partial \mathbf{k}} &= \frac{\partial \tilde{\omega}_k}{\partial \mathbf{x}} \cdot \frac{\partial \langle N \rangle}{\partial \mathbf{k}} \\ \frac{\partial \langle N \rangle}{\partial t} &= \frac{\partial}{\partial \mathbf{k}} \cdot \left\langle \frac{\partial \tilde{\omega}_k}{\partial \mathbf{x}} \tilde{N} \right\rangle \end{aligned}$$

Here we neglect the  $\tilde{\mathbf{V}}$  and  $\mathbf{k} \cdot \tilde{\mathbf{V}}$ , since the Doppler shift by the ion fluid motion is smaller than the effect of the modulation of refraction  $\tilde{\omega}_k$ .

**2.2. Linear response of distribution of quasi-particles.** Set the fluctuations to be

$$\tilde{n} = \sum_{q, \Omega} n_{q, \Omega} \exp(i\mathbf{q} \cdot \mathbf{x} - i\Omega t), \quad \tilde{N} = \sum_{q, \Omega} N_{q, \Omega} \exp(i\mathbf{q} \cdot \mathbf{x} - i\Omega t)$$

$\mathbf{q}, \Omega$  stand for the slow spatiotemporal variation associated with ion-acoustic waves. Then we can get response from equation 2.2,

$$N_{\mathbf{q}, \Omega} = -\frac{\omega_{p0}}{\Omega - \mathbf{q} \cdot \mathbf{v}_g} \frac{n_{\mathbf{q}, \Omega}}{2n_0} \mathbf{q} \cdot \frac{\partial \langle N \rangle}{\partial \mathbf{k}}$$

end.



When the self-interaction of plasma waves is weaker than the decorrelation due to the wave dispersion, i.e.  $\tau_{ac} < \tau_{tr}, \tau_c$ , we can use quasi-linear theory to calculate the mean evolution of energy density,

$$(2.3) \quad \frac{d}{dt} \langle E_k \rangle = -D_N \frac{\partial \langle N \rangle}{\partial \mathbf{k}}$$

$$D_N = \left( \frac{\omega_{p0}}{2n_0} \right)^2 \Re \sum_{\mathbf{q}, \Omega} |n_{\mathbf{q}, \Omega}|^2 \frac{i}{\Omega - \mathbf{q} \cdot \mathbf{v}_g + i\gamma_N} \mathbf{v}_g \cdot \mathbf{q} \mathbf{q} \cdot \frac{\partial \langle N \rangle}{\partial \mathbf{k}}$$

$$\frac{i}{\Omega - \mathbf{q} \cdot \mathbf{v}_g + i\gamma_N} \rightarrow \pi \delta(\Omega - \mathbf{q} \cdot \mathbf{v}_g)$$

which is consistent with previous result.

*Relation to wave-wave interaction*

From Golden rule we know that

$$\text{Rate} \sim \delta(\omega_{k+q} - \omega_k - \omega_q) \approx \frac{i}{\omega_{k+q} - \omega_k - \omega_q}$$

In disparate scale interaction, we have  $\mathbf{q} \ll \mathbf{k}$ , then

$$\frac{i}{\omega_{k+q} - \omega_k - \omega_q} \approx \frac{i}{\omega_k + q \frac{d\omega}{dk} - \omega_k - \omega_q} = \frac{i}{qv_g - \omega_q}$$

The equation 2.3 describe the relation between action density (wave population density)  $\langle N \rangle$  and the energy transfer from plasma waves to ion-acoustic waves. Since the group velocity of plasma waves is  $\mathbf{v}_g = \frac{\partial \omega_k}{\partial \mathbf{k}} = \gamma_T v_{te}^2 \mathbf{k} / \omega_k > 0$ , the damping of plasma waves should satisfy the condition,

$$\frac{d}{dt} \langle E_k \rangle < 0 \Rightarrow \frac{\partial \langle N \rangle}{\partial \mathbf{k}} > 0$$

i.e. the energy transfer from plasma wave quasi-particles to ion-acoustic waves requires a population inversion, and the ion-acoustic waves grow in time at the expense of plasma waves.

**2.3. Growth of ion-acoustic waves.** The influence of Langmuir waves on ion-acoustic waves is due to electron pressure from fast oscillation by the plasma waves. But the ion kinetic energy associated with this rapid oscillation is  $m_e/m_i$  times smaller than that of electrons. In slow varying scales, which is relevant to ion-acoustic waves, the rapid electron oscillation induces an radiation pressure,

$$p_{rad} = \frac{\partial}{\partial \omega} (\omega \epsilon) |_{\omega_{p0}} \frac{|E|^2}{8\pi}$$

where  $E$  is the electric field,  $\epsilon$  is a dielectric function.  $p_{rad}$  is essentially plasma wave energy density. The gradient of  $p_{rad}$ , i.e. ponderomotive force, induces a slow-varying ion motion. In addition to thermal pressure  $p_{th} = c_s^2 \tilde{n} m_i$ , the linearized ion equation of motion is

$$m_i n_0 \frac{\partial \tilde{V}}{\partial t} = - \frac{\partial}{\partial x} (p_{th} + p_{rad})$$

plus the continuity equation,

$$\frac{\partial \tilde{n}}{\partial t} = -n_0 \frac{\partial \tilde{V}}{\partial x}$$



Wave kinetics	Envelope
phonon $N$	<b>E</b> envelope
Wave kinetic equation	Zakharov equation
phase space	real space
adiabatic $N$ conserved	envelope affected by IAWs
stochastic	coherent

TABLE 1. Wave kinetics approach versus envelope formalism for Langmuir turbulence.

eliminating  $\tilde{V}$ , we have dynamics equation for ion-acoustic waves

$$(2.4) \quad \frac{\partial^2 \tilde{n}}{\partial t^2 n_0} = \frac{\partial^2}{\partial x^2} \left( c_s^2 \frac{\tilde{n}}{n_0} + \frac{|E|^2}{8\pi n_0 m_i} \right)$$

write  $\frac{|E|^2}{8\pi n_0} = \int dk \omega_k \tilde{N}$  and use linear response of action density,

$$\tilde{N} = -\frac{q\omega_{p0}}{\Omega - qv_g} \frac{\tilde{n}}{2n_0} \frac{\partial \langle N \rangle}{\partial k}$$

then we have

$$\frac{\partial^2 \tilde{n}}{\partial t^2 n_0} = \frac{\partial^2}{\partial x^2} \left( c_s^2 \frac{\tilde{n}}{n_0} + \frac{1}{m_i} \int dk \omega_k \left( -\frac{q\omega_{p0}}{\Omega - qv_g} \frac{\tilde{n}}{2n_0} \frac{\partial \langle N \rangle}{\partial k} \right) \right)$$

then the dispersion relation is

$$\begin{aligned} \Omega^2 &= q^2 c_s^2 + q^2 \frac{\omega_{p0}^2}{2m_i} \int dk \left( -\frac{q}{\Omega - qv_g} \frac{\partial \langle N \rangle}{\partial k} \right) \\ &= q^2 c_s^2 + q^2 \frac{\omega_{p0}^2}{2m_i} \int dk \left( i\pi \delta(\Omega - qv_g) \frac{\partial \langle N \rangle}{\partial k} \right) \end{aligned}$$

set  $\Omega = qc_s + i\gamma_N$ ,  $\gamma_N \ll qc_s$ , then one has,

$$\Omega = qc_s + i\pi q^2 \frac{\omega_{p0}^2}{4m_i c_s} \int dk \delta(\Omega - qv_g) \frac{\partial \langle N \rangle}{\partial k}$$

so the ion-acoustic wave is unstable if

$$\frac{\partial \langle N \rangle}{\partial k} > 0, \quad \text{at } \Omega \simeq qv_g$$

We can compare the wave kinetics approach to envelope formalism using table 1

### 3. WAVE KINETICS FOR ZONAL FLOW GENERATION

It is worthwhile to note that the zonal flow growth is quite similar to the problem of Langmuir turbulence. In Langmuir turbulence, low frequency test phonons (i.e. ion-acoustic waves) grow by attracting energy from ambient plasmons (i.e. plasma waves). In this case, the zonal flow is the analogue of the ion-acoustic wave, while the drift waves are the analogue of plasma waves, and the test zonal flow interacts with a broad spectrum of drift wave fluctuations.

The essence of the theory for zonal flow growth is:

- Get mean field evolution equation of zonal flow, which relates  $\partial_t \phi_{ZF}$  to  $\langle \phi_{DW}^2 \rangle$ , in the presence of wave pressures and stresses.



- Then calculate the response of the drift wave spectrum to the test zonal flow shear.

This procedure is similar to modulational stability calculations. The time scale separation between low frequency zonal flow and high frequency drift waves enables the utilize of wave kinetics to calculate the response of the drift wave spectrum to the test zonal flow shear.

The zonal flow structure is essentially 2-dimensional. Thus in dimensionless form, the zonal flow potential evolves according to 2D vorticity equation,

$$\frac{\partial}{\partial t} \nabla_r^2 \phi_{ZF} = - \frac{\partial}{\partial r} \langle \tilde{v}_r \nabla^2 \tilde{\phi}_{DW} \rangle - \gamma_d \nabla_r^2 \phi_{ZF}$$

i.e. this equation relates the change of zonal flow vorticity to the drift wave vorticity flux. Zonal flow evolution is then a process driven by vorticity transport, as temperature and density evolution are driven by heat and particle fluxes.

Rewrite the drift wave vorticity flux  $\langle \tilde{v}_r \nabla^2 \tilde{\phi}_{DW} \rangle = B \partial_r \langle \tilde{v}_r \tilde{v}_\theta \rangle$  in this equation, noting  $\tilde{v}_r = -\partial_\theta \tilde{\phi}_{DW} / B$ , one have,

$$(3.1) \quad \frac{\partial}{\partial t} \nabla_r^2 \phi_{ZF} = \frac{1}{B} \nabla_r^2 \int d^2 k k_r k_\theta |\tilde{\phi}_k|^2 - \gamma_d \nabla_r^2 \phi_{ZF}$$

equation 3.1 directly relates the evolution of zonal flow potential to the slow-varying envelope of the drift wave intensity.

The drift wave energy density is

$$E_k = (1 + k_\perp^2 \rho_s^2) |\phi_k|^2$$

the potential enstrophy is

$$Z_k = (1 + k_\perp^2 \rho_s^2)^2 |\phi_k|^2$$

the drift wave dispersion relation is

$$\omega_k = \frac{\omega_{*e}}{1 + k_\perp^2 \rho_s^2}$$

thus the wave action density is

$$N = \frac{E_k}{\omega_k} = (1 + k_\perp^2 \rho_s^2)^2 \frac{|\phi_k|^2}{\omega_{*e}}$$

the  $\omega_{*e} = k_\theta V_*$  here is constant, since  $k_\theta$  is unchanged by zonal flow shearing, i.e.  $\frac{dk_y}{dt} = -\frac{\partial}{\partial y} (k_\theta V_{ZF}(x)) = 0$ . Thus we can relate wave action density to drift wave fluctuation intensity, noting  $\nabla_r^2 \phi_{ZF} = iq \tilde{V}_{ZF}$ , the equation 3.1 becomes

$$(3.2) \quad iq \frac{\partial}{\partial t} \tilde{V}_{ZF} = \frac{1}{B^2} \frac{\partial^2}{\partial r^2} \int d^2 k \frac{k_r k_\theta}{(1 + k_\perp^2 \rho_s^2)^2} \tilde{N} - \gamma_d (iq \tilde{V}_{ZF})$$

The modulational response  $\tilde{N}$  now can be calculated using linearized WKE for zonal flow shears,

$$\frac{\partial \tilde{N}}{\partial t} + v_g \frac{\partial \tilde{N}}{\partial r} + \gamma_k \tilde{N} = \frac{\partial}{\partial r} (k_\theta \tilde{V}_{ZF}) \frac{\partial \langle N \rangle}{\partial k_r}$$

then the modulation  $\tilde{N}$  induced by  $\tilde{V}_{ZF}$  is given by

$$\tilde{N} = - \frac{q k_\theta \tilde{V}_{ZF}}{\Omega - q v_g + i \gamma_k} \frac{\partial \langle N \rangle}{\partial k_r}$$



	Plasma waves and IAWs	DWs and ZFs
High freq fluctuation	plasma wave (plasmon)	drift wave
Low freq structure	ion-acoustic wave (phonon)	zonal flow
Drive mechanism	ponderomotive pressure	Reynolds stress
wave action distribution	plasmon number	potential enstrophy $N = (1 + k_{\perp}^2 \rho_s^2)^2  \phi_k ^2$
Modulational instability	population inversion needed	population inversion unnecessary
Regulator	ion Landau damping of phonon	collisional damping for ZFs

TABLE 2. Langmuir turbulence case versus zonal flow generation case.

so the theory conserves the energy which gives rise to a predator-prey model. Then drift wave turbulence energy is transferred into the energy of zonal flow via the modulational instability.

Now we have introduced both plasma wave-sound wave interaction (i.e. plasmon-phonon) and the drift wave-zonal flow interaction. The comparison is listed in table 2.

#### 4. NONLINEAR SCHRÖDINGER EQUATION FOR LANGMUIR WAVES

4.1. **Influence of ion-acoustic waves on plasma waves.** A heuristic description can be developed by applying the envelope formalism to plasmon-phonon interaction. We can write inhomogeneous plasma waves as

$$\tilde{\mathbf{E}} \sim E(x, t) \mathbf{e}_0 \exp(i\mathbf{k} \cdot \mathbf{x} - i\omega t)$$

where the  $E(x, t)$  indicates the slow-varying envelope, the  $\mathbf{e}_0$  denotes the polarization of the wave field, the  $\exp(i\mathbf{k} \cdot \mathbf{x} - i\omega t)$  is the fast oscillating plasma wave carrier. For plasma waves, the dispersion relation is

$$\omega^2 = \omega_{pe}^2 + \alpha k^2 v_{t,e}^2$$

set  $\omega = \omega_{p0} + i\gamma$  ( $\gamma \ll \omega_{p0}$ ), plug it back, then we have

$$\omega_{p0}^2 + 2i\omega_{p0}\gamma = \omega_{p0}^2 + \alpha v_{t,e}^2 k^2 + \omega_{p0}^2 \frac{\tilde{n}}{n_0}$$

$$2i\omega_{p0}\gamma = \alpha v_{t,e}^2 k^2 + \omega_{p0}^2 \frac{\tilde{n}}{n_0}$$

since  $\gamma \rightarrow \partial_t$  and  $k^2 \rightarrow -\nabla^2$ , we get

$$(4.1) \quad 2i\omega_{p0} \frac{\partial}{\partial t} E = -\alpha v_{t,e}^2 \nabla^2 E + \omega_{p0}^2 \frac{\tilde{n}}{n_0} E$$

together with equation 2.4 which we repost here,

$$(4.2) \quad \frac{\partial^2 \tilde{n}}{\partial t^2} \frac{1}{n_0} = c_s^2 \nabla^2 \left( \frac{\tilde{n}}{n_0} + \frac{|E|^2}{8\pi n_0 m_i c_s^2} \right)$$

the set of equations 4.1 and 4.2 is known as Zakharov equations. They are coupled envelope equation. In the absence of nonlinear coupling, i.e. if  $\tilde{n} \rightarrow 0$ , Eq. 4.1 becomes the Schrodinger equation for a free particle, and if  $|E|^2 \rightarrow 0$ , Eq. 4.2 reduces to the ion-acoustic wave equation.

See previous



put this back into zonal flow evolution equation, then the modulational instability eigenfrequency is,

$$\Omega = \frac{q^2}{B^2} \int d^2k \frac{k_\theta^2 k_r}{\Omega - qv_g + i\gamma_k} \frac{1}{(1 + k_\perp^2 \rho_s^2)^2} \frac{\partial \langle N \rangle}{\partial k_r} - i\gamma_d$$

the imaginary part gives the zonal flow growth rate

$$(3.3) \quad \Gamma = -\frac{q^2}{B^2} \int d^2k \frac{\gamma_k}{(\Omega - qv_g)^2 + \gamma_k^2} \frac{k_\theta^2 k_r}{(1 + k_\perp^2 \rho_s^2)^2} \frac{\partial \langle N \rangle}{\partial k_r} - \gamma_d$$

The growth of zonal flow requires  $\frac{\partial \langle N \rangle}{\partial k_r} < 0$ , which is satisfied for any realistic equilibrium spectrum of drift wave turbulence. In contrast to Langmuir turbulence, there is no population inversion here for zonal flow growth. This is because for drift waves are backward wave,

$$v_g = \frac{\partial \omega_k}{\partial k_r} = \frac{\partial}{\partial k_r} \frac{k_\theta V_*}{1 + k_\perp^2 \rho_s^2} = -\frac{2k_\theta k_r}{(1 + k_\perp^2 \rho_s^2)^2} V_* \Rightarrow \frac{v_g}{v_p} < 0$$

while the plasma waves are forward wave, i.e.  $v_g/v_p > 0$ .

On the other hand,

$$\frac{d}{dt} E_k = N \frac{d\omega_k}{dt}$$

while

$$\begin{aligned} \frac{dk_r}{dt} &= -\frac{\partial}{\partial r} (k_\theta V_{ZF}(x)) \\ &= -k_\theta V'_{ZF} \end{aligned}$$

$$\frac{\partial \langle N \rangle}{\partial t} = \frac{\partial}{\partial k_r} \langle k_\theta \tilde{V}'_{ZF} \tilde{N} \rangle$$

$$(3.4) \quad \begin{aligned} \frac{d \langle E_k \rangle}{dt} &= \omega_k \frac{\partial}{\partial k_r} \langle k_\theta \tilde{V}'_{ZF} \tilde{N} \rangle \\ &= -\sum_q \int d^2k \frac{\partial \omega_k}{\partial k_r} \frac{q^2 k_\theta^2}{\Omega - qv_g + i\gamma_k} |\tilde{V}_{ZF}|^2 \frac{\partial \langle N \rangle}{\partial k_r} \end{aligned}$$

$$(3.5) \quad = \frac{2}{B^2} \sum_q \int d^2k \frac{\gamma_k}{(\Omega - qv_g)^2 + \gamma_k^2} \frac{q^2 k_\theta^2 k_r}{(1 + k_\perp^2 \rho_s^2)^2} |\tilde{V}_{ZF}|^2 \frac{\partial \langle N \rangle}{\partial k_r}$$

since  $\frac{\partial \omega_k}{\partial k_r} < 0$ , growth of zonal flow by depleting energy from drift waves also requires  $\frac{\partial \langle N \rangle}{\partial k_r} < 0$ .

From Eq. 3.2 and 3.3, neglecting collisional damping, we have

$$\begin{aligned} \frac{d}{dt} |\tilde{V}_{ZF}|^2 &= \sum_q 2\Gamma_q |\tilde{V}_{ZF}|^2 \\ &= -\frac{2}{B^2} \sum_q \int d^2k \frac{\gamma_k}{(\Omega - qv_g)^2 + \gamma_k^2} \frac{q^2 k_\theta^2 k_r}{(1 + k_\perp^2 \rho_s^2)^2} |\tilde{V}_{ZF}|^2 \frac{\partial \langle N \rangle}{\partial k_r} \end{aligned}$$

It is thus apparently that

$$\frac{d}{dt} \left( |\tilde{V}_{ZF}|^2 + \langle E_k \rangle \right) = 0$$