

PHYS 201 Mathematical Physics, Fall 2017, Homework 2

Due date: Tuesday, October 17th, 2017

1. Compute the integral around the unit circle $|z| = 1$ for the following functions using Cauchy's Integral Formula. Explain the choice of branch cuts if necessary.

i. $\cot z$

ii. $\frac{ze^z}{z^2+1}$

2. In class, we derived Cauchy's Integral Formula for a point z on the contour C as $\text{PV} \oint_C \frac{f(\zeta)}{\zeta-z} d\zeta = \pi i f(z)$ by integrating on a semicircle around z with the semicircle lying inside the region enclosed by C (for an illustration, see Fig 2-2 of Carrier et al). Show that we obtain the same result when the semicircle is outside of the region enclosed by C .

3. In this exercise, we prove the converse of Cauchy's theorem, called *Morera's theorem*: If a function is continuous in a simple connected region and has the property that $\oint f(z)dz = 0$ on any closed contour lying in that region, then $f(z)$ must be analytic in that region. To prove this, first show that if $\oint f(z)dz = 0$ on any closed contour, then the integral $\int_{z_1}^{z_2} f(z)dz$ depends only on the end points z_1 and z_2 . Next, suppose $F(z_2) - F(z_1) = \int_{z_1}^{z_2} f(z)dz$. Show that the function F is analytic and explain why this implies that f is analytic.

4. In this exercise, we prove the complex analog of the theorem in electrostatics that if a charge-free domain has a constant potential on the boundary, then the potential is a constant inside the domain.

i. Show that if a function $f(z)$ is analytic and has a constant modulus $|f(z)|$ inside a domain D , then $f(z)$ is constant in that domain. (**Hint**: Write $f(z) = u(z) + iv(z)$. Differentiate $|f(z)|^2$ w.r.t the real and imaginary components (x, y) of z , and show that if $|f(z)|$ is a constant, then the Jacobian of the mapping $(x, y) \rightarrow (u, v)$ is zero).

ii. Using the above result, show that if $f(z)$ is analytic and non-vanishing inside D and is constant on the boundary of D , then f is constant everywhere in D . Assume that f is continuous on the closed domain $D \cup \partial D$ i.e., domain plus boundary. (**Hint**: Assume the contrary. By the max/min modulus theorem and continuity, the maximum and minimum of $|f|$ can only lie on the boundary.).

iii. Show that if a function $u(x, y)$ is harmonic and non-constant in a domain D , then u has neither a maximum nor a minimum in D . Using ideas from the previous results,

show that this implies that if u is constant on the boundary, then it is constant everywhere in D . Further, show that if two harmonic functions are equal on the boundary, then they are equal everywhere i.e., given the boundary conditions, the potential is unique. (**Hint:** For the first part, write $g(z) = e^{f(z)}$ where u is the real part of f and use max/min modulus theorem).