

## Problem Set 3: due 05 Nov 2018

- 1) Consider an ensemble of  $M$  stationary test particles in a plasma of  $N$  particles,  $N \gg M$  and  $n\lambda_D^3 > 1$ . Calculate the electrostatic energy of the field produced by the  $M$  test particles, assuming their positions are uncorrelated.
  
- 2) Derive the growth rate of the gentle bump-on-tail instability.
  - a) Develop the system as an off-shoot of the beam-plasma system. Explain when kinetic processes become relevant.
  - b) Derive the wave frequency and growth rate.
  
- 3) Derive the growth rate of the current driven ion acoustic instability.
  - a) Derive the ion acoustic wave kinetically. Calculate frequency and growth. Assume no electron current.
  - b) Now allow electron current, so  $\langle f_e \rangle$  is a shifted Maxwellian. When is instability possible? Take ions as unshifted Maxwellian and  $\delta f_e = |e| \frac{\tilde{\phi}}{T_e} \langle f_e \rangle + \hat{h}_e$ .
  - c) Derive the marginality condition for the CDIA. What parameters control stability?
  - d) Calculate the electron and ion heating.

- 4) a) Consider a chunk of collisionless, self-gravitating matter in one dimension. Here, take a "chunk" to be:

$$f = \begin{cases} f_0, & u_0 - \Delta v < v < u_0 + \Delta v \\ 0, & \text{elsewhere} \end{cases}.$$

Here,  $f_0$  is constant. Take  $u_0, \Delta v$  fixed. Using the Vlasov-Poisson system, calculate the marginal stability criterion for Jeans instability. Compare your result to the case for a self-gravitating gas.

- b) Now consider a plasma, with

$$f = \begin{cases} f_{max} + f_0, & u_0 - \Delta v < v < u_0 + \Delta v \\ f_{max}, & \text{elsewhere} \end{cases}$$

Consider  $f_0 > 0$  and  $f_0 < 0$ .  $f_{max}$  is the usual Maxwellian. Of course  $f_{max} + f_0 > 0$ , for all  $v$ . What is the marginality condition now? Relate your result to the bunching condition discussed in class for the beam-plasma interaction. Hint: Consider the sign of the dielectric function.

- c) For collisionless, self-gravitating matter with an initially Jeans unstable distribution, discuss how the instability might saturate. Hint: Consider simple quasi-linear analysis.
- 5) Consider an electron and ion plasma which is stable, but in which the electrons carry a current, i.e. assume a drift  $u_0$ . Take  $T_i$  finite.
- a) What are the collective resonances? When are they weakly damped, and approaching marginality?
- b) Estimate the thermal fluctuation spectrum  $(\langle E^2 \rangle_{k,\omega} / 8\pi)$  for the system described in Part (a).
- c) *Quantitatively* discuss the breakdown of the test particle model assumptions as the system approaches marginality as  $u_0$  increases.

- 6) For the system of Problem 5:
  - a) Derive the rate of electron-ion momentum transfer. What are the key dimensionless parameters determining this? Assume parameters such that the system is stable.
  - b) How does increasing drift affect the transfer? Assume the system remains stable, but approaches marginality from below.
- 7) Read and summarize the posted article by Rostoker and Rosenbluth on the Test Particle Model. Describe the key ideas of the Test Particle Model and how they are developed.