

Contents

- Contents i

- List of Tables iii

- List of Figures iii

- 7 Mean Field Theory of Phase Transitions 1**
 - 7.1 References 1
 - 7.2 The van der Waals system 2
 - 7.2.1 Equation of state 2
 - 7.2.2 Analytic form of the coexistence curve near the critical point 6
 - 7.2.3 History of the van der Waals equation 10
 - 7.3 Fluids, Magnets, and the Ising Model 11
 - 7.3.1 Lattice gas description of a fluid 11
 - 7.3.2 Phase diagrams and critical exponents 13
 - 7.3.3 Gibbs-Duhem relation for magnetic systems 15
 - 7.3.4 Order-disorder transitions 16
 - 7.4 Mean Field Theory 16
 - 7.4.1 The mean field *Ansatz* 16
 - 7.4.2 Zero external field 18
 - 7.4.3 Finite external field 20
 - 7.4.4 Magnetization dynamics 23

7.4.5	Beyond nearest neighbors	26
7.4.6	Ising model with long-ranged forces	26
7.5	Variational Density Matrix Method	28
7.5.1	The variational principle	28
7.5.2	Variational density matrix for the Ising model	29
7.5.3	q -state Potts model	32
7.5.4	XY Model	33
7.6	Landau Theory of Phase Transitions	37
7.6.1	Quartic free energy with Ising symmetry	37
7.6.2	Cubic terms in Landau theory : first order transitions	39
7.6.3	Magnetization dynamics	40
7.6.4	Sixth order Landau theory : tricritical point	42
7.6.5	Hysteresis for the sextic potential	44
7.7	Mean Field Theory of Fluctuations	45
7.7.1	Correlation and response in mean field theory	45
7.7.2	Calculation of the response functions	47
7.7.3	Beyond the Ising model	50
7.8	Global Symmetries	53
7.8.1	Symmetries and symmetry groups	53
7.8.2	Lower critical dimension	54
7.8.3	Continuous symmetries	56
7.8.4	Random systems : Imry-Ma argument	57
7.9	Ginzburg-Landau Theory	59
7.9.1	Ginzburg-Landau free energy	59
7.9.2	Domain wall profile	60
7.9.3	Derivation of Ginzburg-Landau free energy	61
7.9.4	Ginzburg criterion	64
7.10	Appendix I : Equivalence of the Mean Field Descriptions	66

7.10.1	Variational density matrix	67
7.10.2	Mean field approximation	68
7.11	Appendix II : Additional Examples of Mean Field Theory	69
7.11.1	Blume-Capel model	69
7.11.2	Ising antiferromagnet in an external field	70
7.11.3	Canted quantum antiferromagnet	74
7.11.4	Coupled order parameters	76

List of Tables

7.1	van der Waals parameters for some common gases	4
7.2	Critical exponents	22

List of Figures

7.1	Pressure <i>versus</i> volume for the van der Waals gas	3
7.2	Molar free energy $f(T, v)$ of the van der Waals system	5
7.3	Pressure-volume isotherms for the van der Waals system, with Maxwell construction	6
7.4	Maxwell construction in the (v, p) plane	7
7.5	Universality of the liquid-gas transition	9
7.6	The lattice gas model	12
7.7	Comparison of liquid-gas and Ising ferromagnet phase diagrams	14

7.8	Order-disorder transition on the square lattice	17
7.9	Ising mean field theory at $h = 0$	19
7.10	Ising mean field theory at $h = 0.1$	21
7.11	Dissipative magnetization dynamics $\dot{m} = -f'(m)$	24
7.12	Magnetization hysteresis	25
7.13	Variational field free energy <i>versus</i> magnetization	30
7.14	Phase diagram for the quartic Landau free energy	38
7.15	Behavior of the quartic free energy $f(m) = \frac{1}{2}am^2 - \frac{1}{3}ym^3 + \frac{1}{4}bm^4$	40
7.16	Fixed points for $\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{3}u^3 + \frac{1}{4}u^4$ and flow $\dot{u} = -\varphi'(u)$	41
7.17	Behavior of the sextic free energy $f(m) = \frac{1}{2}am^2 + \frac{1}{4}bm^4 + \frac{1}{6}cm^6$	43
7.18	Sextic free energy $\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{4}u^4 + \frac{1}{6}u^6$ for different values of r	44
7.19	Fixed points $\varphi'(u^*) = 0$ and flow $\dot{u} = -\varphi'(u)$ for the sextic potential	46
7.20	A domain wall in a one-dimensional Ising model	54
7.21	Domain walls in the two and three dimensional Ising model	55
7.22	A domain wall in an XY ferromagnet	57
7.23	Imry-Ma domains and free energy <i>versus</i> domain size	58
7.24	Mean field phase diagram for the Blume-Capel model	71
7.25	Mean field solution for an Ising antiferromagnet in an external field	72
7.26	Mean field phase diagram for an Ising antiferromagnet in an external field	73
7.27	Mean field phase diagram for the model of eqn. 7.340	76
7.28	Phase diagram for the model of eqn. 7.352	80

Chapter 7

Mean Field Theory of Phase Transitions

7.1 References

- M. Kardar, *Statistical Physics of Particles* (Cambridge, 2007)
A superb modern text, with many insightful presentations of key concepts.
- M. Plischke and B. Bergersen, *Equilibrium Statistical Physics* (3rd edition, World Scientific, 2006)
An excellent graduate level text. Less insightful than Kardar but still a good modern treatment of the subject. Good discussion of mean field theory.
- G. Parisi, *Statistical Field Theory* (Addison-Wesley, 1988)
An advanced text focusing on field theoretic approaches, covering mean field and Landau-Ginzburg theories before moving on to renormalization group and beyond.
- J. P. Sethna, *Entropy, Order Parameters, and Complexity* (Oxford, 2006)
An excellent introductory text with a very modern set of topics and exercises. Available online at <http://www.physics.cornell.edu/sethna/StatMech>

7.2 The van der Waals system

7.2.1 Equation of state

Recall the van der Waals equation of state,

$$\left(p + \frac{a}{v^2}\right)(v - b) = RT \quad , \quad (7.1)$$

where $v = N_A V/N$ is the molar volume. Solving for $p(v, T)$, we have

$$p = \frac{RT}{v - b} - \frac{a}{v^2} \quad . \quad (7.2)$$

Let us fix the temperature T and examine the function $p(v)$. Clearly $p(v)$ is a decreasing function of volume for v just above the minimum allowed value $v = b$, as well as for $v \rightarrow \infty$. But is $p(v)$ a monotonic function for all $v \in [b, \infty]$?

We can answer this by computing the derivative,

$$\left(\frac{\partial p}{\partial v}\right)_T = \frac{2a}{v^3} - \frac{RT}{(v - b)^2} \quad . \quad (7.3)$$

Setting this expression to zero for finite v , we obtain the equation¹

$$\frac{2a}{bRT} = \frac{u^3}{(u - 1)^2} \quad , \quad (7.4)$$

where $u \equiv v/b$ is dimensionless. It is easy to see that the function $f(u) = u^3/(u - 1)^2$ has a unique minimum for $u > 1$. Setting $f'(u^*) = 0$ yields $u^* = 3$, and so $f_{\min} = f(3) = \frac{27}{4}$. Thus, for $T > T_c = 8a/27bR$, the LHS of eqn. 7.4 lies below the minimum value of the RHS, and there is no solution. This means that $p(v, T > T_c)$ is a monotonically decreasing function of v .

At $T = T_c$ there is a saddle-node bifurcation. Setting $v_c = bu^* = 3b$ and evaluating $p_c = p(v_c, T_c)$, we have that the location of the *critical point* for the van der Waals system is²

$$p_c = \frac{a}{27b^2} \quad , \quad v_c = 3b \quad , \quad T_c = \frac{8a}{27bR} \quad . \quad (7.5)$$

For $T < T_c$, there are two solutions to eqn. 7.4, corresponding to a local minimum and a local maximum of the function $p(v)$. The locus of points in the (v, p) plane for which $(\partial p/\partial v)_T = 0$ is obtained by setting eqn. 7.3 to zero and solving for T , then substituting this into eqn. 7.2. The result is

$$p^*(v) = \frac{a}{v^2} - \frac{2ab}{v^3} \quad . \quad (7.6)$$

¹There is always a solution to $(\partial p/\partial v)_T = 0$ at $v = \infty$.

²Equivalently, one can obtain the location of the saddle-node bifurcation in the local extrema of $p(T, v)$ by setting $(\partial p/\partial v)_T = 0$ and $(\partial^2 p/\partial v^2)_T = 0$. This yields two equations for the two unknowns (T_c, v_c) . Invoking the van der Waals equation of state then yields $p_c = p(T_c, v_c)$.




Figure 7.1: Pressure *versus* molar volume for the van der Waals gas at temperatures in equal intervals from $T = 1.10 T_c$ (red) to $T = 0.85 T_c$ (blue). The purple curve is $\bar{p}^*(\bar{v})$.

Expressed in terms of dimensionless quantities $\bar{p} = p/p_c$ and $\bar{v} = v/v_c$, this equation becomes

$$\bar{p}^*(\bar{v}) = \frac{3}{\bar{v}^2} - \frac{2}{\bar{v}^3} . \quad (7.7)$$

Along the curve $p = p^*(v)$, the isothermal compressibility, $\kappa_T = -v^{-1}(\partial v/\partial p)_T$ diverges, heralding a *thermodynamic instability*. To understand better, let us compute the free energy of the van der Waals system, $F = E - TS$. Regarding the energy E , we showed back in chapter 2 that

$$\left(\frac{\partial \varepsilon}{\partial v}\right)_T = T \left(\frac{\partial p}{\partial T}\right)_v - p = \frac{a}{v^2} , \quad (7.8)$$

which entails

$$\varepsilon(T, v) = \frac{1}{2}fRT - \frac{a}{v} , \quad (7.9)$$

where $\varepsilon = E/\nu$ is the molar internal energy. The first term is the molar energy of an ideal gas, where f is the number of molecular freedoms, which is the appropriate low density limit. The molar specific heat is then $c_V = (\partial \varepsilon/\partial T)_v = \frac{1}{2}fR$, which means that the molar entropy is

$$s(T, v) = \int^T dT' \frac{c_V}{T'} = \frac{1}{2}fR \ln(T/T_c) + s_1(v) . \quad (7.10)$$

We then write $f = \varepsilon - Ts$, and we fix the function $s_1(v)$ by demanding that $p = -(\partial f/\partial v)_T$. This yields $s_1(v) = R \ln(v - b) + s_0$, where s_0 is a constant. Thus³,

$$f(T, v) = \frac{1}{2}fRT \left(1 - \ln(T/T_c)\right) - \frac{a}{v} - RT \ln(v - b) - Ts_0 . \quad (7.11)$$

gas	$a \left(\frac{\text{L}^2 \cdot \text{bar}}{\text{mol}^2} \right)$	$b \left(\frac{\text{L}}{\text{mol}} \right)$	p_c (bar)	T_c (K)	v_c (L/mol)
Acetone	14.09	0.0994	52.82	505.1	0.2982
Argon	1.363	0.03219	48.72	150.9	0.0966
Carbon dioxide	3.640	0.04267	7404	304.0	0.1280
Ethanol	12.18	0.08407	63.83	516.3	0.2522
Freon	10.78	0.0998	40.09	384.9	0.2994
Helium	0.03457	0.0237	2.279	5.198	0.0711
Hydrogen	0.2476	0.02661	12.95	33.16	0.0798
Mercury	8.200	0.01696	1055	1723	0.0509
Methane	2.283	0.04278	46.20	190.2	0.1283
Nitrogen	1.408	0.03913	34.06	128.2	0.1174
Oxygen	1.378	0.03183	50.37	154.3	0.0955
Water	5.536	0.03049	220.6	647.0	0.0915

Table 7.1: van der Waals parameters for some common gases. (Source: Wikipedia)

We know that under equilibrium conditions, f is driven to a minimum by spontaneous processes. Now suppose that $(\partial^2 f / \partial v^2)_T < 0$ over some range of v at a given temperature T . This would mean that one mole of the system at volume v and temperature T could lower its energy by rearranging into two half-moles, with respective molar volumes $v \pm \delta v$, each at temperature T . The total volume and temperature thus remain fixed, but the free energy changes by an amount $\Delta f = \frac{1}{2}(\partial^2 f / \partial v^2)_T (\delta v)^2 < 0$. This means that the system is unstable – it can lower its energy by dividing up into two subsystems each with different densities (*i.e.* molar volumes). Note that the onset of stability occurs when

$$\left. \frac{\partial^2 f}{\partial v^2} \right|_T = - \left. \frac{\partial p}{\partial v} \right|_T = \frac{1}{v \kappa_T} = 0 \quad , \quad (7.12)$$

which is to say when $\kappa_T = \infty$. As we saw, this occurs at $p = p^*(v)$, given in eqn. 7.6.

However, this condition, $(\partial^2 f / \partial v^2)_T < 0$, is in fact too strong. That is, the system can be unstable even at molar volumes where $(\partial^2 f / \partial v^2)_T > 0$. The reason is shown graphically in fig. 7.2. At the fixed temperature T , for any molar volume v between $v_{\text{liquid}} \equiv v_1$ and $v_{\text{gas}} \equiv v_2$, the system can lower its free energy by *phase separating* into regions of different molar volumes. In general we can write

$$v = (1 - x)v_1 + xv_2 \quad , \quad (7.13)$$

so $v = v_1$ when $x = 0$ and $v = v_2$ when $x = 1$. The free energy upon phase separation is simply

$$f = (1 - x)f_1 + xf_2 \quad , \quad (7.14)$$

where $f_j = f(v_j, T)$. This function is given by the straight black line connecting the points at volumes v_1 and v_2 in fig. 7.2.

³Don't confuse the molar free energy (f) with the number of molecular degrees of freedom (f)!




Figure 7.2: The molar free energy $f(T, v)$ of the van der Waals system at $T = 0.85 T_c$. The dot-dashed black line shows the Maxwell construction connecting molar volumes $v_{1,2}$ on opposite sides of the coexistence curve.

The two equations which give us v_1 and v_2 are

$$\left. \frac{\partial f}{\partial v} \right|_{v_1, T} = \left. \frac{\partial f}{\partial v} \right|_{v_2, T} = \frac{f(T, v_2) - f(T, v_1)}{(v_2 - v_1)} \quad . \quad (7.15)$$

Equivalently, in terms of the pressure, $p = -(\partial f / \partial v)_T$, these equations are equivalent to

$$p(T, v_1) = p(T, v_2) = \frac{1}{v_2 - v_1} \int_{v_1}^{v_2} dv p(T, v) \quad . \quad (7.16)$$

This procedure is known as the *Maxwell construction*, and is depicted graphically in fig. 7.4. When the Maxwell construction is enforced, the isotherms resemble the curves in fig. 7.3. In this figure, all points within the purple shaded region have $\partial^2 f / \partial v^2 < 0$, hence this region is unstable to infinitesimal fluctuations. The boundary of this region is called the *spinodal*, and the spontaneous phase separation into two phases is a process known as *spinodal decomposition*. The dot-dashed orange curve, called the *coexistence curve*, marks the instability boundary for *nucleation*. In a nucleation process, an energy barrier must be overcome in order to achieve the lower free energy state. There is no energy barrier for spinodal decomposition – it is a spontaneous process.

Suppose we follow along an isotherm starting from the high molar volume (gas) phase. If $T > T_c$, the volume v decreases continuously as the pressure p increases. If $T < T_c$, then at the instant the isotherm first intersects the orange boundary curve in fig. 7.3, there is a discontinuous change in the molar volume from high (gas) to low (liquid). This discontinuous change is the hallmark of a *first order phase transition*. Note that the volume discontinuity, $\Delta v = v_- - (T_c - T)^{1/2}$. This is an example of a *critical behavior* in which the *order parameter* ϕ , which in this case may be taken to be the difference $\phi = v_G - v_L$, behaves as a power law in $|T - T_c|$, where T_c is the *critical temperature*. In this case, we have $\phi(T) \propto (T_c - T)_+^\beta$, where $\beta = \frac{1}{2}$ is the exponent, and where $(T_c - T)_+$ is defined to be $T_c - T$ if $T < T_c$ and 0 otherwise. Recall




Figure 7.3: Pressure-volume isotherms for the van der Waals system, as in fig. 7.1, but corrected to account for the Maxwell construction. The boundary of the purple shaded region is the spinodal line $\bar{p}^*(\bar{v})$. The boundary of the orange shaded region is the stability boundary with respect to phase separation, and is called the *coexistence curve*.

the isothermal compressibility is $\kappa_T = -v^{-1}(\partial v/\partial p)_T$. This is finite along the coexistence curve – it diverges only along the spinodal. It therefore diverges at the critical point, which lies at the intersection of the spinodal and the coexistence curve.

It is convenient to express the equation of state and the coexistence curve in terms of dimensionless variables. Writing

$$\bar{p} = \frac{p}{p_c} \quad , \quad \bar{v} = \frac{v}{v_c} \quad , \quad \bar{T} = \frac{T}{T_c} \quad , \quad (7.17)$$

the dimensionless van der Waals equation of state takes the form

$$\bar{p} = \frac{8\bar{T}}{3\bar{v} - 1} - \frac{3}{\bar{v}^2} \quad . \quad (7.18)$$

7.2.2 Analytic form of the coexistence curve near the critical point

We write $v_L = v_c + w_L$ and $v_G = v_c + w_G$. One of our equations is $p(v_c + w_L, T) = p(v_c + w_G, T)$. Taylor expanding in powers of w_L and w_G , we have

$$0 = p_v(v_c, T)(w_G - w_L) + \frac{1}{2} p_{vv}(v_c, T)(w_G^2 - w_L^2) + \frac{1}{6} p_{vvv}(v_c, T)(w_G^3 - w_L^3) + \dots \quad , \quad (7.19)$$

where

$$p_v \equiv \frac{\partial p}{\partial v} \quad , \quad p_{vv} \equiv \frac{\partial^2 p}{\partial v^2} \quad , \quad p_{vvv} \equiv \frac{\partial^3 p}{\partial v^3} \quad , \quad p_{vT} \equiv \frac{\partial^2 p}{\partial v \partial T} \quad , \quad \text{etc.} \quad (7.20)$$




Figure 7.4: Maxwell construction in the (v, p) plane. The system is absolutely unstable between volumes v_d and v_e . For $v \in [v_a, v_d]$ or $v \in [v_e, v_c]$, the solution is unstable with respect to phase separation. Source: *Wikipedia*.

The second equation we write as

$$\int_{w_L}^{w_G} dw p(v_c + w, T) = \frac{1}{2}(w_G - w_L) \left(p(v_c + w_L, T) + p(v_c + w_G, T) \right) . \quad (7.21)$$

Expanding in powers of w_L and w_G , this becomes

$$\begin{aligned} & p(v_c, T) (w_G - w_L) + \frac{1}{2} p_{vv}(v_c, T) (w_G^2 - w_L^2) + \frac{1}{6} p_{vvv}(v_c, T) (w_G^3 - w_L^3) \\ & + \frac{1}{24} p_{vvvv}(v_c, T) (w_G^4 - w_L^4) + \frac{1}{120} p_{vvvvv}(v_c, T) (w_G^5 - w_L^5) + \dots \\ & = \frac{1}{2}(w_G - w_L) \left\{ 2p(v_c, T) + p_v(v_c, T) (w_G + w_L) + \frac{1}{2} p_{vv}(v_c, T) (w_G^2 + w_L^2) \right. \\ & \quad \left. + \frac{1}{6} p_{vvv}(v_c, T) (w_G^3 + w_L^3) + \frac{1}{24} p_{vvvv}(v_c, T) (w_G^4 + w_L^4) + \dots \right\} \end{aligned} \quad (7.22)$$

Subtracting the LHS from the RHS, we find that we can then divide by $\frac{1}{6}(w_G^2 - w_L^2)$, resulting in

$$0 = p_{vv}(v_c, T) + \frac{1}{2} p_{vvv}(v_c, T) (w_G + w_L) + \frac{1}{20} p_{vvvv}(v_c, T) (3w_G^2 + 4w_G w_L + 3w_L^2) + \mathcal{O}(w_{G,L}^3) . \quad (7.23)$$

We now define $w_{\pm} \equiv w_G \pm w_L$. In terms of these variables, eqns. 7.19 and 7.23 become

$$\begin{aligned} 0 &= p_v(v_c, T) + \frac{1}{2} p_{vv}(v_c, T) w_+ + \frac{1}{8} p_{vvv}(v_c, T) (w_+^2 + \frac{1}{3} w_-^2) + \mathcal{O}(w_{\pm}^3) \\ 0 &= p_{vv}(v_c, T) + \frac{1}{2} p_{vvv}(v_c, T) w_+ + \frac{1}{8} p_{vvvv}(v_c, T) (w_+^2 + \frac{1}{5} w_-^2) + \mathcal{O}(w_{\pm}^3) . \end{aligned} \quad (7.24)$$

We now write $T = T_c + \Theta$ and evaluate w_{\pm} as expansions in Θ . Note that $p_v(v_c, T_c) = p_{vv}(v_c, T_c) = 0$, since the critical point is an inflection point in the (v, p) plane. Thus, we have $p_v(v_c, T) = p_{vT}^0 \Theta + \mathcal{O}(\Theta^2)$, where $p_{vT}^0 = p_{vT}(v_c, T_c)$. We can then see that $w_- \propto \sqrt{-\Theta}$, while $w_+ \propto \Theta$ as $\Theta \rightarrow 0^-$, and we have

$$\begin{aligned} 0 &= p_{vT}^0 \Theta + \frac{1}{24} p_{vvv}^0 w_-^2 + \mathcal{O}(\Theta^2) \\ 0 &= p_{vvT}^0 \Theta + \frac{1}{2} p_{vvv}^0 w_+ + \frac{1}{40} p_{vvvv}^0 w_-^2 + \mathcal{O}(\Theta^2) \quad . \end{aligned} \quad (7.25)$$

Thus,

$$\begin{aligned} w_- &= \left(\frac{24 p_{vT}^0}{p_{vvv}^0} \right)^{1/2} \sqrt{-\Theta} + \dots \\ w_+ &= \left(\frac{6 p_{vT}^0 p_{vvvv}^0 - 10 p_{vvv}^0 p_{vvT}^0}{5 (p_{vvv}^0)^2} \right) \Theta + \dots \quad . \end{aligned} \quad (7.26)$$

We then have

$$\begin{aligned} w_L &= - \left(\frac{6 p_{vT}^0}{p_{vvv}^0} \right)^{1/2} \sqrt{-\Theta} + \left(\frac{3 p_{vT}^0 p_{vvvv}^0 - 5 p_{vvv}^0 p_{vvT}^0}{5 (p_{vvv}^0)^2} \right) \Theta + \mathcal{O}(\Theta^{3/2}) \\ w_G &= \left(\frac{6 p_{vT}^0}{p_{vvv}^0} \right)^{1/2} \sqrt{-\Theta} + \left(\frac{3 p_{vT}^0 p_{vvvv}^0 - 5 p_{vvv}^0 p_{vvT}^0}{5 (p_{vvv}^0)^2} \right) \Theta + \mathcal{O}(\Theta^{3/2}) \quad . \end{aligned} \quad (7.27)$$

Close to the critical point, the dimensionless equation of state may be written as $\pi = \pi(\epsilon, t)$, where

$$\bar{p} = 1 + \pi \quad , \quad \bar{v} = 1 + \epsilon \quad , \quad \bar{T} = 1 + t \quad , \quad (7.28)$$

where $\pi(0, 0) = 0$. Equivalently,

$$\pi = \frac{p - p_c}{p_c} \quad , \quad \epsilon = \frac{v - v_c}{v_c} \quad , \quad t = \frac{T - T_c}{T_c} = \frac{\Theta}{T_c} \quad . \quad (7.29)$$

Here π , ϵ , and t are, respectively, the dimensionless deviations of pressure, molar volume, and temperature from their critical point values. For the van der Waals equation of state in eqn. 7.18, we have

$$\begin{aligned} \pi(\epsilon, t) &= \frac{8(1+t)}{2+3\epsilon} - \frac{3}{(1+\epsilon)^2} - 1 \\ &= 4t - 6t\epsilon + 9\epsilon^2 t - \frac{3}{2}\epsilon^3 - \frac{27}{2}\epsilon^3 t + \frac{21}{4}\epsilon^4 + \frac{81}{4}\epsilon^4 t - \frac{99}{8}\epsilon^5 + \dots \quad . \end{aligned} \quad (7.30)$$

Expressed in these dimensionless quantities, eqns. 7.27 take the form

$$\epsilon_{L,G} = \mp \left(\frac{6 \pi_{\epsilon t}^0}{\pi_{\epsilon \epsilon \epsilon}^0} \right)^{1/2} (-t)^{1/2} + \left(\frac{3 \pi_{\epsilon t}^0 \pi_{\epsilon \epsilon \epsilon \epsilon}^0 - 5 \pi_{\epsilon \epsilon \epsilon}^0 \pi_{\epsilon \epsilon t}^0}{5 (\pi_{\epsilon \epsilon \epsilon}^0)^2} \right) t + \mathcal{O}((-t)^{3/2}) \quad . \quad (7.31)$$

For the van der Waals system, we have

$$\pi_{\epsilon t}^0 = -6 \quad , \quad \pi_{\epsilon \epsilon t}^0 = 18 \quad , \quad \pi_{\epsilon \epsilon \epsilon}^0 = -9 \quad , \quad \pi_{\epsilon \epsilon \epsilon \epsilon}^0 = 126 \quad , \quad (7.32)$$




Figure 7.5: Universality of the liquid-gas transition for eight different atomic and molecular fluids, from E. A. Guggenheim, *J. Chem. Phys.* **13**, 253 (1945). Dimensionless temperature $\bar{T} = T/T_c$ versus dimensionless number density $\bar{n} = n/n_c = v_c/v$ is shown. The van der Waals / mean field theory gives $\Delta n = n_{\text{liquid}} - n_{\text{gas}} \propto (-t)^{1/2}$, while experiments show a result closer to $\Delta n \propto (-t)^{1/3}$. Here $t \equiv \bar{T} - 1 = (T - T_c)/T_c$ is the dimensionless temperature deviation with respect to the critical point. (Image adapted from Matthew Schwartz's Harvard lecture notes, adapted from Guggenheim 1945.)

where the derivatives are evaluated at critical point, where $\epsilon = t = \pi = 0$. Thus, for van der Waals,

$$\epsilon_{L,G} = \mp 2\sqrt{-t} - \frac{5}{18}t + \dots \quad (7.33)$$

We identify the difference $\Delta v \equiv v_L - v_G$ as the *order parameter* for the transition which occurs at $T = T_c$. We see that the order parameter behaves as a power law for T just below the critical point, with $\Delta v \propto (-t)^\beta$ and $\beta = \frac{1}{2}$, which is the *order parameter critical exponent*.

The spinodal boundary for the vdW system is then given by the solution to

$$\frac{\partial \pi}{\partial \epsilon} = -6t + 18\epsilon t - \frac{9}{2}\epsilon^2 - \frac{81}{2}\epsilon^2 t + 21\epsilon^3 + \dots \quad (7.34)$$

For the spinodal, it is easy to see that the lowest order nontrivial solution is $\epsilon = \mp \frac{2}{\sqrt{3}}\sqrt{-t}$. For the coexistence curve, we found $\epsilon_{L,G} = \mp 2\sqrt{-t}$.

Fig. 7.5 shows the universality of the liquid-gas transition for eight different fluids: Ne, Ar, Kr, Xe, N₂, O₂, CO, and CH₄. The experimental coexistence curve expressed in dimensionless variables $\bar{n} = 1/\bar{v}$

and $\bar{T} = T/T_c$ is fairly well-fit to the curve⁴

$$\bar{n}_{L,G}(t) = 1 \pm \frac{7}{4}(1 - \bar{T})^{1/3} + \frac{3}{4}(1 - \bar{T}) \quad (7.35)$$

which shows that the critical exponent β is much closer to $\beta = \frac{1}{3}$ than to the vdW value $\beta^{\text{vdW}} = \frac{1}{2}$. The van der Waals equation is in essence a *mean field theory* of the liquid-gas transition.

7.2.3 History of the van der Waals equation

The van der Waals equation of state first appears in van der Waals' 1873 PhD thesis⁵, "*Over de Continuïteit van den Gas - en Vloeistoestand*" ("On the continuity of the gas and liquid state"). In his Nobel lecture⁶, van der Waals writes of how he was inspired by Rudolf Clausius' 1857 treatise on the nature of heat, where it is posited that a gas in fact consists of microscopic particles whizzing around at high velocities. van der Waals reasoned that liquids, which result when gases are compressed, also consist of 'small moving particles': "*Thus I conceived the idea that there is no essential difference between the gaseous and the liquid state of matter...*"

Clausius' treatise showed how his kinetic theory of heat was consistent with Boyle's law for gases ($pV = \text{constant}$ at fixed temperature). van der Waals pondered why this might fail for the non-dilute liquid phase, and he reasoned that there were two principal differences: inter-particle attraction and excluded volume. These considerations prompted him to posit his famous equation,

$$p = \frac{RT}{v - b} - \frac{a}{v^2} \quad (7.36)$$

The first term on the RHS accounts for excluded volume effects, and the second for mutual attractions.

In the limiting case of $p \rightarrow \infty$, the molar volume approaches $v = b$. On physical grounds, one might expect $b = v_0/\zeta$, where $v_0 = N_A \omega_0$ is N_A times the volume ω_0 of a single molecule, and the *packing fraction* is $\zeta = N\omega_0/V = v_0/v$, which is the ratio of the total molecular volume to the total system volume. In three dimensions, the maximum possible packing fraction is for fcc and hcp lattices, each of which have coordination number 12, with $\zeta_{\text{max}} = \frac{\pi}{3\sqrt{2}} = 0.74078$. Dense random packing results in $\zeta_{\text{drp}} = 0.634$. Expanding the vdW equation of state in inverse powers of v yields

$$p = \frac{RT}{v} + \left(b - \frac{a}{RT}\right) \cdot \frac{RT}{v^2} + \mathcal{O}(v^{-3}) \quad (7.37)$$

and we read of the second virial coefficient $B_2 = (b - \frac{a}{RT})/N_A$. For hard spheres, $a = 0$, and the result $B_2 = 4\omega_0$ from the Mayer cluster expansion corresponds to $b_{\text{Mayer}} = 4v_0$, which is larger than the result from even the loosest regular sphere packing, *i.e.* that for a cubic lattice, with $\zeta_{\text{cub}} = \frac{\pi}{6}$.

⁴See M. Schwartz, <https://scholar.harvard.edu/files/schwartz/files/9-phases.pdf>.

⁵Johannes Diderik van der Waals, the eldest of ten children, was the son of a carpenter. As a child he received only a primary school education. He worked for a living until age 25, and was able to enroll in a three-year industrial evening school for working class youth. Afterward he continued his studies independently, in his spare time, working as a teacher. By the time he obtained his PhD, he was 36 years old. He received the Nobel Prize for Physics in 1910.

⁶See http://www.nobelprize.org/nobel_prizes/physics/laureates/1910/waals-lecture.pdf

The law of corresponding states

Another of van der Waals' great achievements was his articulation of the *law of corresponding states*. Recall that the van der Waals equation of state, when written in terms of dimensionless quantities $\bar{p} = p/p_c$, $\bar{v} = v/v_c$, and $\bar{T} = T/T_c$, takes the form of eqn. 7.18. Thus, while the a and b parameters are specific to each fluid – see Tab. 7.1 – when written in terms of these scaled dimensionless variables, the equation of state and all its consequent properties (*i.e.* the liquid-gas phase transition) are *universal*.

The van der Waals equation is best viewed as semi-phenomenological. Interaction and excluded volume effects surely are present, but the van der Waals equation itself only captures them in a very approximate way. It is applicable to gases, where it successfully predicts features that are not present in ideal systems (*e.g.* throttling). It is of only qualitative and pedagogical use in the study of fluids, the essential physics of which lies in the behavior of quantities like the pair distribution function $g(r)$. As we saw in chapter 6, any adequate first principles derivation of $g(r)$ – a function which can be measured in scattering experiments – involves rather complicated approximation schemes to close the BBGKY hierarchy. Else one must resort to numerical simulations such as the Monte Carlo method. Nevertheless, the lessons learned from the van der Waals system are invaluable and they provide us with a first glimpse of what is going on in the vicinity of a phase transition, and how nonanalytic behavior, such as $v_G - v_L \propto (T_c - T)^\beta$ with noninteger exponent β may result due to singularities in the free energy at the critical point.

7.3 Fluids, Magnets, and the Ising Model

7.3.1 Lattice gas description of a fluid

The usual description of a fluid follows from a continuum Hamiltonian of the form

$$\hat{H}(\mathbf{p}, \mathbf{x}) = \sum_{i=1}^N \frac{\mathbf{p}_i^2}{2m} + \sum_{i<j} u(\mathbf{x}_i - \mathbf{x}_j) \quad . \quad (7.38)$$

The potential $u(r)$ is typically central, depending only on the magnitude $|r|$, and short-ranged. Now consider a discretized version of the fluid, in which we divide up space into cells (cubes, say), each of which can accommodate at most one fluid particle (due to excluded volume effects). That is, each cube has a volume on the order of a^3 , where a is the diameter of the fluid particles. In a given cube i we set the occupancy $n_i = 1$ if a fluid particle is present and $n_i = 0$ if there is no fluid particle present. We then have that the potential energy is

$$U = \sum_{i<j} u(\mathbf{x}_i - \mathbf{x}_j) = \frac{1}{2} \sum_{\mathbf{R} \neq \mathbf{R}'} V_{\mathbf{R}\mathbf{R}'} n_{\mathbf{R}} n_{\mathbf{R}'} \quad , \quad (7.39)$$

where $V_{\mathbf{R}\mathbf{R}'} \approx v(\mathbf{R} - \mathbf{R}')$, where R_k is the position at the center of cube k . The grand partition function is then approximated as

$$\Xi(T, V, \mu) \approx \sum_{\{n_{\mathbf{R}}\}} \left(\prod_{\mathbf{R}} \xi^{n_{\mathbf{R}}} \right) \exp \left(-\frac{1}{2} \beta \sum_{\mathbf{R} \neq \mathbf{R}'} V_{\mathbf{R}\mathbf{R}'} n_{\mathbf{R}} n_{\mathbf{R}'} \right) \quad , \quad (7.40)$$




Figure 7.6: The lattice gas model. An occupied cell corresponds to $n = 1$ ($\sigma = +1$), and a vacant cell to $n = 0$ ($\sigma = -1$).

where $\xi = e^{\beta\mu} \lambda_T^{-d} a^d$, and where a is the side length of each cube (chosen to be on the order of the hard sphere diameter). The λ_T^{-d} factor arises from the integration over the momenta. Note $\sum_{\mathbf{R}} n_{\mathbf{R}} = N$ is the total number of fluid particles, so

$$\prod_{\mathbf{R}} \xi^{n_{\mathbf{R}}} = \xi^N = e^{\beta\mu N} \lambda_T^{-Nd} a^{Nd} . \quad (7.41)$$

Thus, we can write a lattice Hamiltonian,

$$\begin{aligned} \hat{H} &= \frac{1}{2} \sum_{\mathbf{R} \neq \mathbf{R}'} V_{\mathbf{R}\mathbf{R}'} n_{\mathbf{R}} n_{\mathbf{R}'} - k_{\text{B}} T \ln \xi \sum_{\mathbf{R}} n_{\mathbf{R}} \\ &= -\frac{1}{2} \sum_{\mathbf{R} \neq \mathbf{R}'} J_{\mathbf{R}\mathbf{R}'} \sigma_{\mathbf{R}} \sigma_{\mathbf{R}'} - H \sum_{\mathbf{R}} \sigma_{\mathbf{R}} + E_0 , \end{aligned} \quad (7.42)$$

where $\sigma_{\mathbf{R}} \equiv 2n_{\mathbf{R}} - 1$ is a spin variable taking the possible values $\{-1, +1\}$, and

$$J_{\mathbf{R}\mathbf{R}'} = -\frac{1}{4} V_{\mathbf{R}\mathbf{R}'} , \quad H = \frac{1}{2} k_{\text{B}} T \ln \xi - \frac{1}{4} \sum'_{\mathbf{R}'} V_{\mathbf{R}\mathbf{R}'} , \quad E_0 = \frac{1}{2} \sum_{\mathbf{R} \neq \mathbf{R}'} V_{\mathbf{R}\mathbf{R}'} , \quad (7.43)$$

where the prime on the sum indicates that $\mathbf{R}' = \mathbf{R}$ is to be excluded. For the Lennard-Jones system, $V_{\mathbf{R}\mathbf{R}'} = v(\mathbf{R} - \mathbf{R}') < 0$ is due to the attractive tail of the potential, hence $J_{\mathbf{R}\mathbf{R}'}$ is positive, which prefers alignment of the spins $\sigma_{\mathbf{R}}$ and $\sigma_{\mathbf{R}'}$. This interaction is therefore *ferromagnetic*. The spin Hamiltonian in eqn. 7.42 is known as the Ising model.

7.3.2 Phase diagrams and critical exponents

The physics of the liquid-gas transition in fact has a great deal in common with that of the transition between a magnetized and unmagnetized state of a magnetic system. The correspondences are⁷

$$p \longleftrightarrow H \quad , \quad v \longleftrightarrow m \quad ,$$

where m is the magnetization density, defined here to be the total magnetization M divided by the number of lattice sites N_s .⁸

$$m = \frac{M}{N_s} = \frac{1}{N_s} \sum_{\mathbf{R}} \langle \sigma_{\mathbf{R}} \rangle \quad . \quad (7.44)$$

Sketches of the phase diagrams are reproduced in fig. 7.7. Of particular interest is the *critical point*, which occurs at (T_c, p_c) in the fluid system and (T_c, H_c) in the magnetic system, with $H_c = 0$ by symmetry.

In the fluid, the coexistence curve in the (p, T) plane separates high density (liquid) and low density (vapor) phases. The specific volume v (or the density $n = v^{-1}$) jumps discontinuously across the coexistence curve. In the magnet, the coexistence curve in the (H, T) plane separates positive magnetization and negative magnetization phases. The magnetization density m jumps discontinuously across the coexistence curve. For $T > T_c$, the latter system is a *paramagnet*, in which the magnetization varies smoothly as a function of H . This behavior is most apparent in the bottom panel of the figure, where $v(p)$ and $m(H)$ curves are shown.

For $T < T_c$, the fluid exists in a *two phase region*, which is spatially inhomogeneous, supporting local regions of high and low density. There is no stable homogeneous thermodynamic phase for (T, v) within the two phase region shown in the middle left panel. Similarly, for the magnet, there is no stable homogeneous thermodynamic phase at fixed temperature T and magnetization m if (T, m) lies within the coexistence region. Rather, the system consists of blobs where the spin is predominantly up, and blobs where the spin is predominantly down.

Note also the analogy between the isothermal compressibility κ_T and the isothermal susceptibility χ_T :

$$\kappa_T = -\frac{1}{v} \left(\frac{\partial v}{\partial p} \right)_T \quad , \quad \chi_T = \left(\frac{\partial m}{\partial H} \right)_T \quad , \quad (7.45)$$

with $\kappa_T(T_c, p_c) = \infty$ and $\chi_T(T_c, H_c) = \infty$.

The *order parameter* for a second order phase transition is a quantity which vanishes in the disordered phase and is finite in the ordered phase. For the fluid, the order parameter can be chosen to be $\Psi \propto (v_{\text{vap}} - v_{\text{liq}})$, the difference in the specific volumes of the vapor and liquid phases. In the vicinity of the critical point, the system exhibits power law behavior in many physical quantities, *viz.*

$$m(T, H_c) \sim (T_c - T)_+^\beta \quad , \quad \chi(T, H_c) \sim |T - T_c|^{-\gamma} \quad , \quad C_M(T, H_c) \sim |T - T_c|^{-\alpha} \quad , \quad m(T_c, H) \sim \pm |H|^{1/\delta} \quad . \quad (7.46)$$

⁷ H is more properly analogous to μ . However, since $\mu = \mu(p, T)$, H can also be regarded as analogous to p . Note also that $\beta p = z \lambda_T^{-d}$ for the ideal gas, in which case $\xi = z(a/\lambda_T)^d$ is proportional to $p/k_B T$.

⁸Note the distinction between the number of lattice sites N_s and the number of occupied cells N . According to our definitions, $N = \frac{1}{2}(M + N_s)$.




Figure 7.7: Comparison of the liquid-gas phase diagram with that of the Ising ferromagnet.

The quantities α , β , γ , and δ are the *critical exponents* associated with the transition. These exponents satisfy certain equalities, such as the Rushbrooke and Griffiths relations:

$$\alpha + 2\beta + \gamma = 2 \quad (\text{Rushbrooke}) \quad , \quad \beta + \gamma = \beta\delta \quad (\text{Griffiths}) \quad . \quad (7.47)$$

Originally such relations were derived as inequalities, and only after the advent of scaling and renormalization group theories it was realized that they held as equalities.

In addition to the exponents α , β , γ , and δ , one defines the *correlation length exponent* ν from the behavior of the two-point correlation function $C(\mathbf{r}, T) = \langle \psi(0) \psi(\mathbf{r}) \rangle$, where $\psi(\mathbf{r})$ is a local operator, such as the

local density in a fluid or the local spin polarization in a magnet⁹. In the limit $T \rightarrow T_c$, one has

$$C(\mathbf{r}, T, h) = r^{-(d-2+\eta)} \phi(r/\xi(T), H/\xi_H(T)) \quad , \quad (7.48)$$

where η is the *anomalous exponent*, $\phi(r/\xi, H/\xi_H)$ is a *scaling function*, $\xi(T) \propto |T - T_c|^{-\nu}$ is the *correlation length*, and $\xi_H(T) \propto |T - T_c|^\Delta$, with $\Delta = \beta\delta$, is a field scale. As we have seen, for the lattice gas system the effective magnetic field H is a proxy for the pressure or the chemical potential. Along with the new exponents η and ν come additional exponent relations,

$$(2 - \eta)\nu = \gamma \quad , \quad d\nu = 2 - \alpha \quad (\text{hyperscaling}) \quad . \quad (7.49)$$

Thus there are three relations among the six critical exponents $\alpha, \beta, \gamma, \delta, \eta$, and ν , which entails that there are three independent values among the six.

7.3.3 Gibbs-Duhem relation for magnetic systems

Homogeneity of $E(S, M, N_s)$ means $E = TS + HM + \mu N_s$, by Euler's theorem. After invoking the First Law $dE = T dS + H dM + \mu dN_s$, we have

$$S dT + M dH + N_s d\mu = 0 \quad . \quad (7.50)$$

Now consider two magnetic phases in coexistence. We must have $d\mu_1 = d\mu_2$, hence

$$d\mu_1 = -s_1 dT - m_1 dH = -s_2 dT - m_2 dH = d\mu_2 \quad , \quad (7.51)$$

where $m = M/N_s$ is the magnetization per site and $s = S/N_s$ is the specific entropy. Thus, we obtain the Clapeyron equation for magnetic systems,

$$\left(\frac{dH}{dT} \right)_{\text{coex}} = -\frac{s_1 - s_2}{m_1 - m_2} \quad . \quad (7.52)$$

Thus, if $m_1 \neq m_2$ and $(dH/dT)_{\text{coex}} = 0$, then we must have $s_1 = s_2$, which says that there is *no latent heat associated with the transition*. This absence of latent heat is a consequence of the *symmetry* which guarantees that $G(T, H, N_s) = G(T, -H, N_s)$.

Recall our discussion in §2.12.2 of the Clausius-Clapeyron relation for liquid-gas systems. From $G = E - TS + pV = G(T, p, N)$, the differential of the Gibbs free energy per particle, $\mu = G/N$, is given by $d\mu = -s dT + v dp$, where $v = V/N$ is the volume per particle¹⁰. This leads to the Clapeyron relation,

$$\left(\frac{dp}{dT} \right)_{\text{coex}} = \frac{s_2 - s_1}{v_2 - v_1} = \frac{\ell}{T \Delta v} \quad , \quad (7.53)$$

which determines the slope of the coexistence curve in the (T, p) plane.

⁹The local 'order parameter field' $\psi(\mathbf{r})$ may carry vector or tensor indices. In general, it transforms as the fundamental representation of the global symmetry group G .

¹⁰In §2.12.2 we considered, equivalently, the differential of the molar free energy $g = N_A \mu$, with v the molar volume.

7.3.4 Order-disorder transitions

Another application of the Ising model lies in the theory of order-disorder transitions in alloys. Examples include Cu_3Au , CuZn , and other compounds. In CuZn , the Cu and Zn atoms occupy sites of a body centered cubic (BCC) lattice, forming an alloy known as β -brass. Below $T_c \simeq 740\text{ K}$, the atoms are ordered, with the Cu preferentially occupying one simple cubic sublattice and the Zn preferentially occupying the other.

The energy is a sum of pairwise interactions, with a given link contributing ε_{AA} , ε_{BB} , or ε_{AB} , depending on whether it is an A-A, B-B, or A-B/B-A link. Here A and B represent Cu and Zn, respectively. Thus, we can write the energy of the link $\langle ij \rangle$ as

$$E_{ij} = \varepsilon_{AA} P_i^A P_j^A + \varepsilon_{BB} P_i^B P_j^B + \varepsilon_{AB} (P_i^A P_j^B + P_i^B P_j^A) \quad , \quad (7.54)$$

where

$$P_i^A = \frac{1}{2}(1 + \sigma_i) = \begin{cases} 1 & \text{if site } i \text{ contains Cu} \\ 0 & \text{if site } i \text{ contains Zn} \end{cases} \quad , \quad P_i^B = \frac{1}{2}(1 - \sigma_i) = \begin{cases} 1 & \text{if site } i \text{ contains Zn} \\ 0 & \text{if site } i \text{ contains Cu} \end{cases} \quad .$$

The Hamiltonian is then

$$\begin{aligned} \hat{H} &= \sum_{\langle ij \rangle} E_{ij} = \sum_{\langle ij \rangle} \left\{ \frac{1}{4}(\varepsilon_{AA} + \varepsilon_{BB} - 2\varepsilon_{AB}) \sigma_i \sigma_j + \frac{1}{4}(\varepsilon_{AA} - \varepsilon_{BB}) (\sigma_i + \sigma_j) + \frac{1}{4}(\varepsilon_{AA} + \varepsilon_{BB} + 2\varepsilon_{AB}) \right\} \\ &= -J \sum_{\langle ij \rangle} \sigma_i \sigma_j - H \sum_i \sigma_i + E_0 \quad , \end{aligned} \quad (7.55)$$

where the exchange constant J and the magnetic field H are given by

$$J = \frac{1}{4}(2\varepsilon_{AB} - \varepsilon_{AA} - \varepsilon_{BB}) \quad , \quad H = \frac{1}{4}(\varepsilon_{BB} - \varepsilon_{AA}) \quad , \quad (7.56)$$

and $E_0 = \frac{1}{8}Nz(\varepsilon_{AA} + \varepsilon_{BB} + 2\varepsilon_{AB})$, where N is the total number of lattice sites and $z = 8$ is the *lattice coordination number*, which is the number of nearest neighbors of any given site.

Note that

$$\begin{aligned} 2\varepsilon_{AB} > \varepsilon_{AA} + \varepsilon_{BB} &\implies J > 0 \quad (\text{ferromagnetic}) \\ 2\varepsilon_{AB} < \varepsilon_{AA} + \varepsilon_{BB} &\implies J < 0 \quad (\text{antiferromagnetic}) \quad . \end{aligned} \quad (7.57)$$

The antiferromagnetic case is depicted in fig. 7.8.

7.4 Mean Field Theory

7.4.1 The mean field Ansatz

Consider the Ising model Hamiltonian,

$$\hat{H} = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j - H \sum_i \sigma_i \quad , \quad (7.58)$$




Figure 7.8: Order-disorder transition on the square lattice. Below $T = T_c$, order develops spontaneously on the two $\sqrt{2} \times \sqrt{2}$ sublattices. There is perfect sublattice order at $T = 0$ (left panel).

where the first sum on the RHS is over all links of the lattice. Each spin can be either ‘up’ ($\sigma = +1$) or ‘down’ ($\sigma = -1$). We further assume that the spins are located on a Bravais lattice¹¹ and that the coupling $J_{ij} = J(|\mathbf{R}_i - \mathbf{R}_j|)$, where \mathbf{R}_i is the position of the i^{th} spin.

On each site i we decompose σ_i into a contribution from its thermodynamic average and a fluctuation:

$$\sigma_i = \langle \sigma_i \rangle + \delta\sigma_i \quad . \quad (7.59)$$

We will write $\langle \sigma_i \rangle \equiv m$, the local magnetization (dimensionless), and assume that m is independent of position i . Then

$$\begin{aligned} \sigma_i \sigma_j &= (m + \delta\sigma_i)(m + \delta\sigma_j) \\ &= m^2 + m(\delta\sigma_i + \delta\sigma_j) + \delta\sigma_i \delta\sigma_j \\ &= -m^2 + m(\sigma_i + \sigma_j) + \delta\sigma_i \delta\sigma_j \quad . \end{aligned} \quad (7.60)$$

The last term on the RHS of the second equation above is quadratic in the fluctuations, and we assume this to be negligibly small. This neglect of the fluctuations is the *mean field Ansatz*, and results in the *mean field Hamiltonian*

$$\hat{H}_{\text{MF}} = \frac{1}{2}NzJm^2 - (H + zJm) \sum_i \sigma_i \quad , \quad (7.61)$$

where N is the total number of lattice sites. The first term is a constant, although the value of m is yet to be determined. The Boltzmann weights are then completely determined by the second term, which is just what we would write down for a Hamiltonian of *noninteracting* spins in an effective ‘mean field’

$$H_{\text{eff}} = H + zJm \quad . \quad (7.62)$$

In other words, $H_{\text{eff}} = H_{\text{ext}} + H_{\text{int}}$, where the external field is applied field $H_{\text{ext}} = H$, and the ‘internal field’ is $H_{\text{int}} = zJm$. The internal field accounts for the interaction with the *average* values of all other

¹¹A Bravais lattice is one in which any site is equivalent to any other site through an appropriate discrete translation. Examples of Bravais lattices include the linear chain, square, triangular, simple cubic, face-centered cubic, *etc.* lattices. The honeycomb lattice is not a Bravais lattice, because there are two sets of inequivalent sites – those in the center of a Y and those in the center of an upside down Y.

spins coupled to a spin at a given site, hence it is often called the ‘mean field’. Since the spins are noninteracting, we have

$$m = \frac{e^{\beta H_{\text{eff}}} - e^{-\beta H_{\text{eff}}}}{e^{\beta H_{\text{eff}}} + e^{-\beta H_{\text{eff}}}} = \tanh\left(\frac{H + zJm}{k_{\text{B}}T}\right) . \quad (7.63)$$

It is a simple matter to solve for the free energy, given the noninteracting Hamiltonian \hat{H}_{MF} . The partition function is

$$Z = \text{Tr} e^{-\beta \hat{H}_{\text{MF}}} = e^{-\frac{1}{2}\beta N z J m^2} \left(\sum_{\sigma} e^{\beta(H+zJm)\sigma} \right)^N = e^{-\beta F} . \quad (7.64)$$

We now define dimensionless variables:

$$f \equiv \frac{F}{N z J} , \quad \theta \equiv \frac{k_{\text{B}}T}{zJ} , \quad h \equiv \frac{H}{zJ} , \quad (7.65)$$

and obtain the dimensionless free energy

$$f(m, h, \theta) = \frac{1}{2}m^2 - \theta \ln \cosh\left(\frac{m+h}{\theta}\right) - \theta \ln 2 . \quad (7.66)$$

Differentiating with respect to m gives the mean field equation,

$$m = \tanh\left(\frac{m+h}{\theta}\right) , \quad (7.67)$$

which is equivalent to the self-consistency requirement, $m = \langle \sigma_i \rangle$. In terms of the dimensionless variables θ and h , the physics is universal, and independent of details such as the magnitude of J , the value of z , and Boltzmann’s constant.

7.4.2 Zero external field

When $h = 0$ the mean field equation becomes $m = \tanh(m/\theta)$. This nonlinear equation can be solved graphically, as in the top panel of fig. 7.9. The RHS is a tanh function which gets steeper with decreasing dimensionless temperature θ . If, at $m = 0$, the slope of $\tanh(m/\theta)$ is smaller than unity, then the curve $y = \tanh(m/h)$ will intersect $y = m$ only at $m = 0$. However, if the slope is larger than unity, there will be three such intersections. Since the slope is $1/\theta$, we identify $\theta_c = 1$ as the *mean field transition temperature*.

The mean field free energies are plotted in the bottom panel of fig. 7.9. It is possible to make analytical progress by assuming m is small and Taylor expanding the free energy $f(m, \theta)$ in powers of m when we are very close to the critical point, *i.e.* when $|\theta - \theta_c| \ll 1$. Then we have

$$f(m, \theta) = -\theta \ln 2 + \frac{1}{2}m^2 - \theta \ln \cosh\left(\frac{m}{\theta}\right) = f_0 + \frac{1}{2}(\theta - \theta_c)m^2 + \frac{m^4}{12} + \dots , \quad (7.68)$$

with $f_0 = -\theta \ln 2$. Note that the sign of the quadratic term is positive for $\theta > \theta_c$ and negative for $\theta < \theta_c$. Thus, the shape of the free energy $f(m, \theta)$ as a function of m qualitatively changes at $\theta = \theta_c = 1$, the mean field transition temperature, also known as the (dimensionless) *critical temperature*. Within our mean field theory, the predicted critical temperature is $T_c = zJ\theta_c = zJ$.




Figure 7.9: Results for $h = 0$. Upper panels: graphical solution to self-consistency equation $m = \tanh(m/\theta)$ at temperatures $\theta = 0.65$ (blue) and $\theta = 1.5$ (dark red). Lower panel: mean field free energy, with energy shifted by $\theta \ln 2$ so that $f(m = 0, \theta) = 0$.

In the high temperature phase, $\theta > \theta_c$, and there is a unique minimum to $f(\theta, m)$ lying at $m = 0$. This is the disordered phase, where the *order parameter* m vanishes. By contrast, in the low temperature phase $\theta < \theta_c$, there are three solutions to the mean field equations. One solution is always at $m = 0$. The other two solutions must be related by the \mathbb{Z}_2 symmetry of the free energy ($m \rightarrow -m$ with $h = 0$). For $\theta > \theta_c$, the free energy $f(m, \theta)$ has a single minimum at $m = 0$. Below θ_c , the curvature at $m = 0$ reverses, and $m = 0$ becomes a local maximum. There are then two equivalent minima symmetrically displaced on either side of $m = 0$. Differentiating with respect to m , we find these additional local minima to lie at $m^2 = 3(\theta_c - \theta) + \mathcal{O}((\Delta\theta)^2)$. Thus, we find for $|\theta - \theta_c| \ll 1$,

$$m(\theta, h = 0) = \pm\sqrt{3} (\theta_c - \theta)_+^{1/2} \quad , \quad (7.69)$$

where the $+$ subscript indicates that this solution is only for $\theta_c - \theta > 0$. As the blue curve in fig. 7.9 shows, these nonzero solutions for m in the low temperature phase lie at a lower value of the free energy f than $m = 0$.

Again, for $\theta > \theta_c$ the only solution is $m = 0$. The high temperature phase is thus one where the \mathbb{Z}_2 (*i.e.* $m \rightarrow -m$) symmetry is *unbroken*. In the low temperature phase, the magnetization m is nonzero, and takes on one of two possible values which are degenerate in free energy. The degeneracy is guaranteed by the \mathbb{Z}_2 symmetry present when $h = 0$. But the system must somehow choose! This is the phenomenon of *spontaneous symmetry breaking* (SSB). The exponent with which $m(\theta)$ vanishes as $\theta \rightarrow \theta_c^-$ is denoted as

β . That is, $m(\theta, h = 0) \propto (\theta_c - \theta)_+^\beta$ with $\beta = \frac{1}{2}$ for our mean field theory.

Specific heat

We can now expand the free energy $f(\theta, h = 0)$. We find

$$f(\theta, h = 0) = \begin{cases} -\theta \ln 2 & \text{if } \theta > \theta_c \\ -\theta \ln 2 - \frac{3}{4}(\theta_c - \theta)^2 + \mathcal{O}((\theta_c - \theta)^4) & \text{if } \theta < \theta_c \end{cases} . \quad (7.70)$$

Thus, if we compute the heat capacity, we find in the vicinity of $\theta = \theta_c$

$$c_V = -\theta \frac{\partial^2 f}{\partial \theta^2} = \begin{cases} 0 & \text{if } \theta > \theta_c \\ \frac{3}{2} & \text{if } \theta < \theta_c \end{cases} . \quad (7.71)$$

Thus, the specific heat is *discontinuous* at $\theta = \theta_c$. We emphasize that our results here are only valid for $|\theta - \theta_c| \ll 1$. The general result valid for all θ (within our mean field theory) is¹²

$$c_V(\theta) = \frac{1}{\theta} \cdot \frac{m^2(\theta) - m^4(\theta)}{\theta - 1 + m^2(\theta)} , \quad (7.72)$$

With this expression one can check both limits $\theta \rightarrow 0$ and $\theta \rightarrow \theta_c$. As $\theta \rightarrow 0$ the magnetization saturates and one has $m^2(\theta) \simeq 1 - 4e^{-2/\theta}$. The numerator then vanishes as $e^{-2/\theta}$, which overwhelms the denominator that itself vanishes as θ^2 . As a result, $c_V(\theta \rightarrow 0) = 0$, as expected. As $\theta \rightarrow m^2 = 3(\theta_c - \theta) + \dots$, invoking $m^2 \simeq 3(m^2 = 3(\theta_c - \theta) + \dots - \theta)$ we recover $c_V(\theta_c^-) = \frac{3}{2}$. In the theory of critical phenomena, $c_V(\theta) \propto |\theta - \theta_c|^{-\alpha}$ as $\theta \rightarrow \theta_c$. We see that mean field theory yields $\alpha = 0$.

7.4.3 Finite external field

Let us first assume $h \ll |\theta - 1| \ll 1$, *i.e.* that we are very close to the critical point $(\theta_c, h_c) = (1, 0)$. The mean field solution for $m(\theta, h)$ then be small, and we may expand the free energy from eqn. 7.66 in m and h , *viz.*

$$\begin{aligned} f(m, h, \theta) &= -\theta \ln 2 + \frac{1}{2}(1 - \theta^{-1})m^2 + \frac{m^4}{12\theta^3} - \frac{hm}{\theta} \\ &= f_0 + \frac{1}{2}(\theta - \theta_c)m^2 + \frac{1}{12}m^4 - hm + \dots \end{aligned} \quad (7.73)$$

Note that we have only gone to linear order in h . Setting $\partial f / \partial m = 0$, we obtain

$$\frac{1}{3}m^3 + (\theta - \theta_c) \cdot m - h = 0 \quad . \quad (7.74)$$

¹²To obtain this result, one writes $f = f(\theta, m(\theta))$ and then differentiates twice with respect to θ , using the chain rule. Along the way, any naked (*i.e.* undifferentiated) term proportional to $\partial f / \partial m$ may be dropped, since this vanishes at any θ by the mean field equation.




Figure 7.10: $f(m, \theta, h)$ for $h = 0.1$. Upper panels: graphical solution to the self-consistency equation $m = \tanh((m+h)/\theta)$ at temperatures $\theta = 0.65$ (blue), $\theta = 0.9$ (dark green), and $\theta = 1.5$ (dark red). Lower panel: mean field free energy, with energy shifted by $\theta \ln 2$ so that $f(\theta, m = 0) = 0$.

If $\theta > \theta_c$ then we have a solution $m = h/(\theta - \theta_c)$. The m^3 term can be ignored because it is higher order in h , and we have assumed $h \ll |\theta - \theta_c| \ll 1$. This is known as the *Curie-Weiss law*¹³. The magnetic susceptibility behaves as

$$\chi(\theta) = \frac{\partial m}{\partial h} = \frac{1}{\theta - \theta_c} \propto |\theta - \theta_c|^{-\gamma} \quad , \quad (7.75)$$

where the magnetization critical exponent γ is $\gamma = 1$. If $\theta < \theta_c$ then while there is still a solution at $m = h/(\theta - \theta_c)$, it lies at a local maximum of the free energy, as shown in fig. 7.10. The minimum of the free energy occurs close to the $h = 0$ solution $m = m_0(\theta) \equiv \sqrt{3}(\theta_c - \theta)$, and writing $m = m_0 + \delta m$ we

¹³Pierre Curie was a pioneer in the fields of crystallography, magnetism, and radiation physics. In 1880, Pierre and his older brother Jacques discovered piezoelectricity. He was 21 years old at the time. It was in 1895 that Pierre made the first systematic studies of the effects of temperature on magnetic materials, and he formulated what is known as *Curie's Law*, $\chi = C/T$, where C is a constant. Curie married Marie Sklodowska in the same year. Their research turned toward radiation, recently discovered by Becquerel and Röntgen. In 1898, Pierre and Marie Curie discovered radium. They shared the 1903 Nobel Prize in Physics with Becquerel. Marie went on to win the 1911 Nobel Prize in Chemistry and was the first person ever awarded two Nobel Prizes. Their daughter Irène Joliot Curie shared the 1935 Prize in Chemistry (with her husband), also for work on radioactivity. Pierre Curie met an untimely and unfortunate end in the Spring of 1906. Walking across the Place Dauphine, he slipped and fell under a heavy horse-drawn wagon carrying military uniforms. His skull was crushed by one of the wagon wheels, killing him instantly. Later that year, Pierre-Ernest Weiss proposed a modification of Curie's Law to account for ferromagnetism. This became known as the Curie-Weiss law, $\chi = C/(T - T_c)$.

Exponent	MFT	2D Ising (exact)	3D Ising (numerical)	binary liquid (3D)
α	0	0	0.11008(1)	0.113(5)
β	1/2	1/8	0.326419(3)	0.316(8)
γ	1	7/4	1.237075(10)	1.240(7)
δ	3	15	4.78984(1)	–
η	0	1/3	0.036298(2)	0.016(7)
ν	1/2	1	0.629971(4)	0.625(5)

Table 7.2: Critical exponents from mean field theory as compared with exact results for the two-dimensional Ising model, numerical results for the three-dimensional Ising model, and experiments on the liquid-gas transition in the binary fluids triethylamine and water (α : D. Beysens and A. Bourgou, *Phys. Rev. A* **19**, 2407 (1979)), isobutyric acid and water (β : S. C. Greer, *Phys. Rev. A* **14**, 1770 (1976)), and 3-methylpentane-nitroethane (γ, η, ν : R. F. Chang *et al.*, *Phys. Rev. Lett.* **37**, 1481 (1976)).

find δm to linear order in h as $\delta m(\theta, h) = h/2(\theta_c - \theta)$. Thus,

$$m(\theta, h) = \sqrt{3}(\theta_c - \theta)^{1/2} + \frac{h}{2(\theta_c - \theta)} \quad . \quad (7.76)$$

Once again, we find that $\chi(\theta)$ diverges as $|\theta - \theta_c|^{-\gamma}$ with $\gamma = 1$. The exponent γ on either side of the transition is the same.

Finally, we can set $\theta = \theta_c$ and examine $m(h)$. We find, from eqn. 7.74,

$$m(\theta = \theta_c, h) = (3h)^{1/3} \propto h^{1/\delta} \quad , \quad (7.77)$$

where δ is a new critical exponent. Mean field theory gives $\delta = 3$. Note that at $\theta = \theta_c = 1$ we have $m = \tanh(m + h)$, and inverting we find

$$h(m, \theta = \theta_c) = \frac{1}{2} \ln \left(\frac{1+m}{1-m} \right) - m = \frac{m^3}{3} + \frac{m^5}{5} + \dots \quad , \quad (7.78)$$

which is consistent with what we just found for $m(h, \theta = \theta_c)$.

How well does mean field theory do in describing the phase transition of the Ising model? In table 7.2 we compare our mean field results for the exponents α, β, γ , and δ with exact values for the two-dimensional Ising model, numerical work on the three-dimensional Ising model, and experiments on the liquid-gas transition in CO_2 . The first thing to note is that the exponents are dependent on the dimension of space, and this is something that mean field theory completely misses. In fact, it turns out that the mean field exponents are exact provided $d > d_u$, where d_u is the *upper critical dimension* of the theory. For the Ising model, $d_u = 4$, and above four dimensions (which is of course unphysical) the mean field exponents are in fact exact. We see that all in all the MFT results compare better with the three dimensional exponent values than with the two-dimensional ones – this makes sense since MFT does better in higher dimensions. The reason for this is that higher dimensions means more nearest neighbors, which effectively reduces the relative importance of the fluctuations we neglected to include.

Metastable states in the ordered phase

Consider the free energy $f(m, h, \theta)$ in eqn. 7.66, now for general θ not restricted to the immediate vicinity of θ_c . When $\theta < \theta_c$ and h is sufficiently small – just how small we are about to find out – the free energy as a function of m has one local maximum and two local minima, one of which is the thermodynamically stable state (*i.e.* the one for which $mh > 0$), and the other (with $mh < 0$) is metastable. Consider the case $h > 0$. As the temperature is raised, the metastable local minimum at $m < 0$ eventually vanishes, annihilating with the local maximum in a saddle-node bifurcation. To find where this happens, one sets $\partial f / \partial m = 0$ and $\partial^2 f / \partial m^2 = 0$ simultaneously. From eqn. 7.66, we have

$$\frac{\partial f}{\partial m} = m - \tanh\left(\frac{m+h}{\theta}\right) \quad , \quad \frac{\partial^2 f}{\partial m^2} = 1 - \theta^{-1} \operatorname{sech}^2\left(\frac{m+h}{\theta}\right) \quad . \quad (7.79)$$

Thus $h = \theta \tanh^{-1}(m) - m$, and using $\operatorname{sech}^2 x = 1 - \tanh^2 x$, we have $m^2 = 1 - \theta$ and

$$h^*(\theta) = \sqrt{1 - \theta} - \frac{\theta}{2} \ln\left(\frac{1 + \sqrt{1 - \theta}}{1 - \sqrt{1 - \theta}}\right). \quad (7.80)$$

The solutions lie at $h = \pm h^*(\theta)$. For $\theta < \theta_c = 1$ and $h \in [-h^*(\theta), +h^*(\theta)]$, there are three solutions to the mean field equation. Equivalently we could in principle invert the above expression to obtain $\theta^*(h)$. For $\theta > \theta^*(h)$, there is only a single global minimum in the free energy $f(m)$ and there is no local minimum. Note $\theta^*(h = 0) = 1$. Note that we could in principle invert the above relation to obtain $\theta^*(h)$, but alas this is not analytically possible.

7.4.4 Magnetization dynamics

Dissipative processes drive physical systems to minimum energy states. We can crudely model the dissipative dynamics of a magnet by writing the phenomenological equation

$$\frac{dm}{ds} = \frac{1}{\Gamma} \frac{dm}{dt} = -\frac{\partial f}{\partial m} \quad , \quad (7.81)$$

where Γ has the dimensions of frequency and $s = \Gamma t$ is the (dimensionless) rescaled time. Under these dynamics, the free energy is never increasing:

$$\frac{df}{ds} = \frac{\partial f}{\partial m} \frac{dm}{ds} = -\left(\frac{\partial f}{\partial m}\right)^2 \leq 0 \quad . \quad (7.82)$$

Clearly the *fixed point* of these dynamics, where $\dot{m} = 0$, is a solution to the mean field equation $\frac{\partial f}{\partial m} = 0$. The time dependent $m(s)$ thus evolves until it reaches the first fixed point encountered, which is to say the first local extremum of the function $f(m)$. This extremum could be a global minimum, but it could also be a local minimum or even an inflection point¹⁴.

The phase flow for the equation $\dot{m} = -f'(m)$ is shown in fig. 7.11. As we have seen, for any value of h there is a temperature θ^* below which the free energy $f(m)$ has two local minima and one local

¹⁴Since $f(s)$ is never increasing, the extremum cannot be a local maximum.




Figure 7.11: Dissipative magnetization dynamics $\dot{m} = -f'(m)$. Bottom panel shows $h^*(\theta)$ from eqn. 7.80. For (θ, h) within the blue shaded region, the free energy $f(m)$ has a global minimum plus a local minimum and a local maximum. Otherwise $f(m)$ has only a single global minimum. Top panels show an imperfect bifurcation in the magnetization dynamics at $h = 0.0215$, for which $\theta^* = 0.90$. Temperatures shown: $\theta = 0.65$ (blue), $\theta = \theta^*(h) = 0.90$ (green), and $\theta = 1.2$. The rightmost stable fixed point corresponds to the global minimum of the free energy. The bottom of the middle two upper panels shows $h = 0$, where both of the attractive fixed points and the repulsive fixed point coalesce into a single attractive fixed point (supercritical pitchfork bifurcation).

maximum. When $h = 0$ the minima are degenerate, but at finite h one of the minima is a global minimum. Thus, for $\theta < \theta^*(h)$ there are three solutions to the mean field equations. In the language of dynamical systems, under the dynamics of eqn. 7.81, minima of $f(m)$ correspond to attractive fixed points and maxima to repulsive fixed points. If $h > 0$, the rightmost of these fixed points corresponds to the global minimum of the free energy. As θ is increased, this fixed point evolves smoothly. At $\theta = \theta^*$, the (metastable) local minimum and the local maximum coalesce and annihilate in a saddle-node bifurcation. However at $h = 0$ all three fixed points coalesce at $\theta = \theta_c$ and the bifurcation is a supercritical pitchfork. As a function of t at finite h , the dynamics are said to exhibit an *imperfect bifurcation*, which is a deformed supercritical pitchfork.

The solution set for the mean field equation is simply expressed by inverting the tanh function to obtain $h(\theta, m)$. One readily finds

$$h(\theta, m) = \frac{\theta}{2} \ln \left(\frac{1+m}{1-m} \right) - m \quad . \quad (7.83)$$




Figure 7.12: Top panel : hysteresis as a function of ramping the dimensionless magnetic field h at $\theta = 0.40$. Dark red arrows below the curve follow evolution of the magnetization on slow increase of h . Dark grey arrows above the curve follow evolution of the magnetization on slow decrease of h . Bottom panel : solution set for $m(\theta, h)$ as a function of h at temperatures $\theta = 0.40$ (blue), $\theta = \theta_c = 1.0$ (dark green), and $t = 1.25$ (red).

As we see in the bottom panel of fig. 7.12, $m(h)$ becomes multivalued for $h \in [-h^*(\theta), +h^*(\theta)]$, where $h^*(\theta)$ is given in eqn. 7.80. Now imagine that $\theta < \theta_c$ and we slowly ramp the field h from a large negative value to a large positive value, and then slowly back down to its original value. On the time scale of the magnetization dynamics, we can regard $h(s)$ as a constant. (Remember the time variable is s here.) Thus, $m(s)$ will flow to the nearest stable fixed point. Initially the system starts with $m = -1$ and h large and negative, and there is only one fixed point, at $m^* \approx -1$. As h slowly increases, the fixed point value m^* also slowly increases. As h exceeds $-h^*(\theta)$, a saddle-node bifurcation occurs, and two new fixed points are created at positive m , one stable and one unstable. The global minimum of the free energy still lies at the fixed point with $m^* < 0$. However, when h crosses $h = 0$, the global minimum of the free energy lies at the most positive fixed point m^* . The dynamics, however, keep the system stuck in what is a metastable phase. This persists until $h = +h^*(\theta)$, at which point another saddle-node bifurcation occurs, and the attractive fixed point at $m^* < 0$ annihilates with the repulsive fixed point. The dynamics then act quickly to drive m to the only remaining fixed point. This process is depicted in the top panel of fig. 7.12. As one can see from the figure, the the system follows a stable fixed point until the fixed point disappears, even though that fixed point may not always correspond to a global minimum of the free energy. The resulting $m(h)$ curve is then not reversible as a function of time, and it possesses a characteristic shape known as a *hysteresis loop*. Etymologically, the word *hysteresis* derives from the Greek $\upsilon\sigma\tau\epsilon\rho\eta\sigma\iota\varsigma$, which means ‘lagging behind’ (and not from $\iota\sigma\tau\omicron\rho\iota\alpha$, which means

‘inquiry’). Systems which are hysteretic exhibit a *history-dependence* to their status, which is not uniquely determined by external conditions. Hysteresis may be exhibited with respect to changes in applied magnetic field, changes in temperature, or changes in other externally determined parameters.

7.4.5 Beyond nearest neighbors

Up to this point we have assumed nearest-neighbor interactions on a lattice of coordination number z . Suppose, instead, that we had started with the more general model,

$$\hat{H} = -\frac{1}{2} \sum_{i \neq j} J_{ij} \sigma_i \sigma_j - H \sum_i \sigma_i \quad , \quad (7.84)$$

where $J_{ij} = J(|\mathbf{R}_i - \mathbf{R}_j|)$ is the coupling between spins on sites i and j ¹⁵. In the top equation above, each pair (ij) is counted once in the interaction term; this may be replaced by a sum over all i and j if we include a factor of $\frac{1}{2}$.¹⁶ The resulting mean field Hamiltonian is then

$$\hat{H}_{\text{MF}} = \frac{1}{2} N \hat{J}(0) m^2 - (H + \hat{J}(0) m) \sum_i \sigma_i \quad . \quad (7.85)$$

Here, $\hat{J}(\mathbf{q})$ is the lattice Fourier transform of the interaction function $J(\mathbf{R})$:¹⁷

$$\hat{J}(\mathbf{q}) = \sum_{\mathbf{R}} J(\mathbf{R}) e^{-i\mathbf{q} \cdot \mathbf{R}} \quad . \quad (7.86)$$

For nearest neighbor interactions only, one has $\hat{J}(0) = zJ$, where z is the *lattice coordination number*, i.e. the number of nearest neighbors of any given site. The scaled free energy is as in eqn. 7.66, with $f = F/N\hat{J}(0)$, $\theta = k_{\text{B}}T/\hat{J}(0)$, and $h = H/\hat{J}(0)$. The analysis proceeds precisely as before, and we conclude $\theta_c = 1$, i.e. $k_{\text{B}}T_c^{\text{MF}} = \hat{J}(0)$. For example, on the simple cubic lattice there are six nearest neighbors and twelve next-nearest neighbors. Thus, if J is the nearest neighbor coupling and J' the next-nearest neighbor coupling, we have $\hat{J}(0) = 6J + 12J'$.

7.4.6 Ising model with long-ranged forces

Consider an Ising model where $J_{ij} = J/N$ for all i and j , so that there is a very weak interaction between every pair of spins. The Hamiltonian is then

$$\hat{H} = -\frac{J}{2N} \left(\sum_i \sigma_i \right)^2 - H \sum_k \sigma_k \quad . \quad (7.87)$$

¹⁵Note that we may write $\frac{1}{2} \sum_{i \neq j} A_{ij}$ as $\sum_{i < j} A_{ij}$ provided $A = A^t$ is symmetric. Clearly $A_{ij} = \sigma_i \sigma_j$ is symmetric under interchange of indices i and j . Each version of the sum counts all unique *pairs* $(i, j) = (j, i)$ exactly once.

¹⁶The self-interaction terms with $i = j$ contribute a constant to \hat{H} and may be either included or excluded. However, this property only pertains to the $\sigma_i = \pm 1$ model. For higher spin versions of the Ising model, say where $S_i \in \{-1, 0, +1\}$, then S_i^2 is not constant and we should explicitly exclude the self-interaction terms.

¹⁷The sum in the discrete Fourier transform is over all ‘direct Bravais lattice vectors’ and the wavevector \mathbf{q} may be restricted to the ‘first Brillouin zone’. These terms are familiar from elementary solid state physics.

The partition function is

$$Z = \text{Tr}_{\{\sigma_i\}} \exp \left[\frac{\beta J}{2N} \left(\sum_i \sigma_i \right)^2 + \beta H \sum_i \sigma_i \right] . \quad (7.88)$$

We now invoke the Gaussian integral,

$$\int_{-\infty}^{\infty} dx e^{-\alpha x^2 - \beta x} = \sqrt{\frac{\pi}{\alpha}} e^{\beta^2/4\alpha} . \quad (7.89)$$

Thus,

$$\exp \left[\frac{\beta J}{2N} \left(\sum_i \sigma_i \right)^2 \right] = \left(\frac{N\beta J}{2\pi} \right)^{1/2} \int_{-\infty}^{\infty} dm e^{-\frac{1}{2}N\beta J m^2 + \beta J m \sum_i \sigma_i} , \quad (7.90)$$

and we can write the partition function as

$$Z = \left(\frac{N\beta J}{2\pi} \right)^{1/2} \int_{-\infty}^{\infty} dm e^{-\frac{1}{2}N\beta J m^2} \left(\sum_{\sigma} e^{\beta(H+Jm)\sigma} \right)^N = \left(\frac{N}{2\pi\theta} \right)^{1/2} \int_{-\infty}^{\infty} dm e^{-NA(m)/\theta} , \quad (7.91)$$

where $\theta = k_B T/J$, $h = H/J$, and

$$A(m) = \frac{1}{2}m^2 - \theta \ln \left[2 \cosh \left(\frac{h+m}{\theta} \right) \right] . \quad (7.92)$$

Since $N \rightarrow \infty$, we can perform the integral using the method of steepest descents. Thus, we must set

$$\left. \frac{dA}{dm} \right|_{m^*} = 0 \quad \implies \quad m^* = \tanh \left(\frac{m^* + h}{\theta} \right) . \quad (7.93)$$

Expanding about $m = m^*$, we write

$$A(m) = A(m^*) + \frac{1}{2}A''(m^*) (m - m^*)^2 + \frac{1}{6}A'''(m^*) (m - m^*)^3 + \dots . \quad (7.94)$$

Performing the integrations, we obtain

$$\begin{aligned} Z &= \left(\frac{N}{2\pi\theta} \right)^{1/2} e^{-NA(m^*)/\theta} \int_{-\infty}^{\infty} d\nu \exp \left[-\frac{NA''(m^*)}{2\theta} \nu^2 - \frac{NA'''(m^*)}{6\theta} \nu^3 + \dots \right] \\ &= \frac{1}{\sqrt{A''(m^*)}} e^{-NA(m^*)/\theta} \cdot \left\{ 1 + \mathcal{O}(N^{-1}) \right\} . \end{aligned} \quad (7.95)$$

The corresponding free energy per site

$$f = \frac{F}{NJ} = A(m^*) + \frac{\theta}{2N} \ln A''(m^*) + \mathcal{O}(N^{-2}) , \quad (7.96)$$

where m^* is the solution to the mean field equation which minimizes $A(m)$. Mean field theory is *exact* for this model!

7.5 Variational Density Matrix Method

7.5.1 The variational principle

Suppose we are given a Hamiltonian \hat{H} . From this we construct the free energy, F :

$$F = E - TS = \text{Tr}(\varrho \hat{H}) + k_B T \text{Tr}(\varrho \ln \varrho) \quad . \quad (7.97)$$

Here, ϱ is the *density matrix*¹⁸. A physical density matrix must be (i) normalized (*i.e.* $\text{Tr} \varrho = 1$), (ii) Hermitian, and (iii) non-negative definite (*i.e.* all the eigenvalues of ϱ must be non-negative).

Our goal is to extremize the free energy subject to the various constraints on ϱ . Let us assume that ϱ is diagonal in the basis of eigenstates of \hat{H} , *i.e.*

$$\varrho = \sum_{\gamma} P_{\gamma} |\gamma\rangle\langle\gamma| \quad , \quad (7.98)$$

where P_{γ} is the probability that the system is in state $|\gamma\rangle$. Then

$$F = \sum_{\gamma} E_{\gamma} P_{\gamma} + k_B T \sum_{\gamma} P_{\gamma} \ln P_{\gamma} \quad . \quad (7.99)$$

Thus, the free energy is a function of the set $\{P_{\gamma}\}$. We now extremize F subject to the normalization constraint. This means we form the extended function

$$F^*(\{P_{\gamma}\}, \lambda) = F(\{P_{\gamma}\}) + \lambda \left(\sum_{\gamma} P_{\gamma} - 1 \right) \quad , \quad (7.100)$$

and then freely extremize over both the probabilities $\{P_{\gamma}\}$ as well as the Lagrange multiplier λ . This yields the Boltzmann distribution,

$$P_{\gamma}^{\text{eq}} = \frac{1}{Z} \exp(-E_{\gamma}/k_B T) \quad , \quad (7.101)$$

where $Z = \sum_{\gamma} e^{-E_{\gamma}/k_B T} = \text{Tr} e^{-\hat{H}/k_B T}$ is the canonical partition function, which is related to λ through

$$\lambda = k_B T (\ln Z - 1) \quad . \quad (7.102)$$

Note that the Boltzmann weights are, appropriately, all positive.

If the spectrum of \hat{H} is bounded from below, our extremum should in fact yield a minimum for the free energy F . Furthermore, since we have freely minimized over all the probabilities, subject to the single normalization constraint, *any distribution* $\{P_{\gamma}\}$ *other than the equilibrium one must yield a greater value of* F .

Alas, the Boltzmann distribution, while exact, is often intractable to evaluate. For one-dimensional systems, there are general methods such as the transfer matrix approach which do permit an exact evaluation of the free energy. However, beyond one dimension the situation is in general hopeless. A family of solvable (“integrable”) models exists in two dimensions, but their solutions require specialized techniques and are extremely difficult. The idea behind the variational density matrix approximation is to construct a tractable *trial* density matrix ϱ which depends on a set of variational parameters $\{x_{\alpha}\}$, and to minimize with respect to this (finite) set.

¹⁸How do we take the logarithm of a matrix? The rule is this: $A = \ln B$ if $B = \exp(A)$. The exponential of a matrix may be evaluated via its Taylor expansion.

7.5.2 Variational density matrix for the Ising model

Consider once again the Ising model Hamiltonian,

$$\hat{H} = - \sum_{i<j} J_{ij} \sigma_i \sigma_j - H \sum_i \sigma_i \quad . \quad (7.103)$$

The states of the system $|\gamma\rangle$ may be labeled by the values of the spin variables: $|\gamma\rangle \longleftrightarrow |\sigma_1, \sigma_2, \dots\rangle$. We assume the density matrix is diagonal in this basis, *i.e.*

$$\varrho_N(\gamma | \gamma') \equiv \varrho(\gamma) \delta_{\gamma, \gamma'} \quad , \quad (7.104)$$

where $\delta_{\gamma, \gamma'} = \prod_i \delta_{\sigma_i, \sigma'_i}$. Indeed, this is the case for the exact density matrix, which is to say the Boltzmann weight,

$$\varrho_N(\sigma_1, \sigma_2, \dots) = \frac{1}{Z} e^{-\beta \hat{H}(\sigma_1, \dots, \sigma_N)} \quad . \quad (7.105)$$

We now write a *trial density matrix* which is a product over contributions from independent single sites:

$$\varrho_N(\sigma_1, \sigma_2, \dots) = \prod_i \varrho(\sigma_i) \quad , \quad (7.106)$$

where

$$\varrho(\sigma) = \left(\frac{1+m}{2} \right) \delta_{\sigma, 1} + \left(\frac{1-m}{2} \right) \delta_{\sigma, -1} \quad . \quad (7.107)$$

Note that we've changed our notation slightly. We are denoting by $\varrho(\sigma)$ the corresponding diagonal element of the matrix

$$\varrho = \begin{pmatrix} \frac{1+m}{2} & 0 \\ 0 & \frac{1-m}{2} \end{pmatrix} \quad , \quad (7.108)$$

and the full density matrix is a tensor product over the single site matrices: $\varrho_N = \varrho \otimes \varrho \otimes \dots \otimes \varrho$.

Note that ϱ and hence ϱ_N are appropriately normalized. The variational parameter here is m , which, if ρ is to be non-negative definite, must satisfy $-1 \leq m \leq 1$. The quantity m has the physical interpretation of the average spin on any given site, since

$$\langle \sigma_i \rangle = \sum_{\sigma} \varrho(\sigma) \sigma = m. \quad (7.109)$$

We may now evaluate the average energy:

$$\begin{aligned} E &= \text{Tr}(\varrho_N \hat{H}) = - \sum_{i<j} J_{ij} m^2 - H \sum_i m \\ &= -\frac{1}{2} N \hat{J}(0) m^2 - N H m \quad , \end{aligned} \quad (7.110)$$

where once again $\hat{J}(0)$ is the discrete Fourier transform of $J(\mathbf{R})$ at wavevector $\mathbf{q} = 0$. The entropy is given by

$$\begin{aligned} S &= -k_B \text{Tr}(\varrho_N \ln \varrho_N) = -N k_B \text{Tr}(\varrho \ln \varrho) \\ &= -N k_B \left\{ \left(\frac{1+m}{2} \right) \ln \left(\frac{1+m}{2} \right) + \left(\frac{1-m}{2} \right) \ln \left(\frac{1-m}{2} \right) \right\} \quad . \end{aligned} \quad (7.111)$$




Figure 7.13: Variational field free energy $\Delta f = f(m, h, \theta) + \theta \ln 2$ versus magnetization m at six equally spaced temperatures interpolating between ‘high’ ($\theta = 1.25$, red) and ‘low’ ($\theta = 0.75$, blue) values. Top panel: $h = 0$. Bottom panel: $h = 0.06$.

We now define the dimensionless free energy per site: $f \equiv F/N\hat{J}(0)$. We have

$$f(m, h, \theta) = -\frac{1}{2}m^2 - hm + \theta \left\{ \left(\frac{1+m}{2} \right) \ln \left(\frac{1+m}{2} \right) + \left(\frac{1-m}{2} \right) \ln \left(\frac{1-m}{2} \right) \right\} , \quad (7.112)$$

where $\theta \equiv k_B T / \hat{J}(0)$ is the dimensionless temperature, and $h \equiv H / \hat{J}(0)$ the dimensionless magnetic field, as before. We extremize $f(m)$ by setting

$$\frac{\partial f}{\partial m} = 0 = -m - h + \frac{\theta}{2} \ln \left(\frac{1+m}{1-m} \right) . \quad (7.113)$$

Solving for m , we obtain

$$m = \tanh \left(\frac{m+h}{\theta} \right) , \quad (7.114)$$

which is precisely what we found in eqn. 7.67.

Note that the optimal value of m indeed satisfies the requirement $|m| \leq 1$ of non-negative probability. This nonlinear equation may be solved graphically. For $h = 0$, the unmagnetized solution $m = 0$ always applies. However, for $\theta < 1$ there are two additional solutions at $m = \pm m_A(\theta)$, with $m_A(\theta) = \sqrt{3(1-\theta)} + \mathcal{O}((1-\theta)^{3/2})$ for t close to (but less than) one. These solutions, which are related by the \mathbb{Z}_2 symmetry of the $h = 0$ model, are in fact the low energy solutions. This is shown clearly in figure

7.13, where the variational free energy $f(m, t)$ is plotted as a function of m for a range of temperatures interpolating between ‘high’ and ‘low’ values. At the *critical temperature* $\theta_c = 1$, the lowest energy state changes from being unmagnetized (high temperature) to magnetized (low temperature).

For $h > 0$, there is no longer a \mathbb{Z}_2 symmetry (*i.e.* $\sigma_i \rightarrow -\sigma_i \forall i$). The high temperature solution now has $m > 0$ (or $m < 0$ if $h < 0$), and this smoothly varies as t is lowered, approaching the completely polarized limit $m = 1$ as $\theta \rightarrow 0$. At very high temperatures, the argument of the tanh function is small, and we may approximate $\tanh(x) \simeq x$, in which case

$$m(h, \theta) = \frac{h}{\theta - \theta_c} . \quad (7.115)$$

This is called the *Curie-Weiss law*. One can infer θ_c from the high temperature susceptibility $\chi(\theta) = (\partial m / \partial h)_{h=0}$ by plotting χ^{-1} versus θ and extrapolating to obtain the θ -intercept. In our case, $\chi(\theta) = (\theta - \theta_c)^{-1}$. For low θ and weak h , there are two inequivalent minima in the free energy.

When m is small, it is appropriate to expand $f(m, h, \theta)$, obtaining

$$f(m, h, \theta) = -\theta \ln 2 - hm + \frac{1}{2} (\theta - 1) m^2 + \frac{\theta}{12} m^4 + \frac{\theta}{30} m^6 + \frac{\theta}{56} m^8 + \dots . \quad (7.116)$$

This is known as the *Landau expansion* of the free energy in terms of the *order parameter* m . An order parameter is a thermodynamic variable ϕ which distinguishes ordered and disordered phases. Typically $\phi = 0$ in the disordered (high temperature) phase, and $\phi \neq 0$ in the ordered (low temperature) phase. When the order sets in continuously, *i.e.* when ϕ is continuous across θ_c , the phase transition is said to be *second order*. When ϕ changes abruptly, the transition is *first order*. It is also quite commonplace to observe phase transitions between two ordered states. For example, a crystal, which is an ordered state, may change its lattice structure, say from a high temperature tetragonal phase to a low temperature orthorhombic phase. When the high T phase possesses the same symmetries as the low T phase, as in the tetragonal-to-orthorhombic example, the transition may be second order. When the two symmetries are completely unrelated, for example in a hexagonal-to-tetragonal transition, or in a transition between a ferromagnet and an antiferromagnet, the transition is in general first order.

Throughout this discussion, we have assumed that the interactions J_{ij} are predominantly *ferromagnetic*, *i.e.* $J_{ij} > 0$, so that all the spins prefer to align. When $J_{ij} < 0$, the interaction is said to be *antiferromagnetic* and prefers anti-alignment of the spins (*i.e.* $\sigma_i \sigma_j = -1$). Clearly not every pair of spins can be anti-aligned – there are two possible spin states and a thermodynamically extensive number of spins. But on the square lattice, for example, if the only interactions J_{ij} are between nearest neighbors and the interactions are antiferromagnetic, then the lowest energy configuration ($T = 0$ ground state) will be one in which spins on opposite sublattices are anti-aligned. The square lattice is *bipartite* – it breaks up into two interpenetrating sublattices A and B (which are themselves square lattices, rotated by 45° with respect to the original, and with a larger lattice constant by a factor of $\sqrt{2}$), such that any site in A has nearest neighbors in B, and *vice versa*. The honeycomb lattice is another example of a bipartite lattice. So is the simple cubic lattice. The triangular lattice, however, is not bipartite (it is *tripartite*). Consequently, with nearest neighbor antiferromagnetic interactions, the triangular lattice Ising model is highly *frustrated*. The moral of the story is this: antiferromagnetic interactions can give rise to complicated magnetic ordering, and, when frustrated by the lattice geometry, may have finite specific entropy even at $T = 0$.

7.5.3 q -state Potts model

The Hamiltonian for the Potts model is

$$\hat{H} = - \sum_{i < j} J_{ij} \delta_{\sigma_i, \sigma_j} - H \sum_i \delta_{\sigma_i, 1} \quad . \quad (7.117)$$

Here, $\sigma_i \in \{1, \dots, q\}$, with integer q . This is the so-called ' q -state Potts model'. The quantity H is analogous to an external magnetic field, and preferentially aligns (for $H > 0$) the local spins in the $\sigma = 1$ direction. We will assume $H \geq 0$.

The q -component set is conveniently taken to be the integers from 1 to q , but it could be anything, such as

$$\sigma_i \in \{\text{tomato, penny, ostrich, Grateful Dead ticket from 1987, ...}\} \quad . \quad (7.118)$$

The interaction energy is $-J_{ij}$ if sites i and j contain the same object (q possibilities), and 0 if i and j contain different objects ($q^2 - q$ possibilities).

The two-state Potts model is equivalent to the Ising model. Let the allowed values of σ be ± 1 . Then the quantity

$$\delta_{\sigma, \sigma'} = \frac{1}{2} + \frac{1}{2} \sigma \sigma' \quad (7.119)$$

equals 1 if $\sigma = \sigma'$, and is zero otherwise. The three-state Potts model cannot be written as a simple three-state Ising model, *i.e.* one with a bilinear interaction $\sigma \sigma'$ where $\sigma \in \{-1, 0, +1\}$. However, it is straightforward to verify the identity

$$\delta_{\sigma, \sigma'} = 1 + \frac{1}{2} \sigma \sigma' + \frac{3}{2} \sigma^2 \sigma'^2 - (\sigma^2 + \sigma'^2) \quad . \quad (7.120)$$

Thus, the $q = 3$ -state Potts model is equivalent to a $S = 1$ (three-state) Ising model which includes both bilinear ($\sigma \sigma'$) and biquadratic ($\sigma^2 \sigma'^2$) interactions, as well as a local field term which couples to the square of the spin, σ^2 . In general one can find such correspondences for higher q Potts models, but, as should be expected, the interactions become increasingly complex, with bi-cubic, bi-quartic, bi-quintic, *etc.* terms. Such a formulation, however, obscures the beautiful S_q symmetry inherent in the model, where S_q is the permutation group on q symbols, which has $q!$ elements.

Getting back to the mean field theory, we write the single site variational density matrix ϱ as a diagonal matrix with entries

$$\varrho(\sigma) = x \delta_{\sigma, 1} + \left(\frac{1-x}{q-1} \right) (1 - \delta_{\sigma, 1}) \quad , \quad (7.121)$$

with $\varrho_N(\sigma_1, \dots, \sigma_N) = \varrho(\sigma_1) \cdots \varrho(\sigma_N)$. Note that $\text{Tr}(\varrho) = 1$. The variational parameter is x . When $x = q^{-1}$, all states are equally probable. But for $x > q^{-1}$, the state $\sigma = 1$ is preferred, and the other $(q-1)$ states have identical but smaller probabilities. It is a simple matter to compute the energy and entropy:

$$\begin{aligned} E &= \text{Tr}(\varrho_N \hat{H}) = -\frac{1}{2} N \hat{J}(0) \left\{ x^2 + \frac{(1-x)^2}{q-1} \right\} - NHx \\ S &= -k_B \text{Tr}(\varrho_N \ln \varrho_N) = -Nk_B \left\{ x \ln x + (1-x) \ln \left(\frac{1-x}{q-1} \right) \right\} \quad . \end{aligned} \quad (7.122)$$

The dimensionless free energy per site is then

$$f(x, \theta, h) = -\frac{1}{2} \left\{ x^2 + \frac{(1-x)^2}{q-1} \right\} + \theta \left\{ x \ln x + (1-x) \ln \left(\frac{1-x}{q-1} \right) \right\} - hx \quad , \quad (7.123)$$

where $h = H/\hat{J}(0)$. We now extremize with respect to x to obtain the mean field equation,

$$\frac{\partial f}{\partial x} = 0 = -x + \frac{1-x}{q-1} + \theta \ln x - \theta \ln \left(\frac{1-x}{q-1} \right) - h \quad . \quad (7.124)$$

Note that for $h = 0$, $x = q^{-1}$ is a solution, corresponding to a disordered state in which all states are equally probable. At high temperatures, for small h , we expect $x - q^{-1} \propto h$. Indeed, using `Mathematica` one can set $x \equiv q^{-1} + s$ and expand the mean field equation in powers of s . One obtains

$$h = \frac{q(q\theta - 1)}{q-1} s + \frac{q^3(q-2)\theta}{2(q-1)^2} s^2 + \mathcal{O}(s^3) \quad . \quad (7.125)$$

For weak fields, $|h| \ll 1$, and we have

$$s(\theta) = \frac{(q-1)h}{q(q\theta - 1)} + \mathcal{O}(h^2) \quad , \quad (7.126)$$

which again is of the Curie-Weiss form. The difference $s = x - q^{-1}$ is the order parameter for the transition.

Finally, one can expand the free energy in powers of s , obtaining the Landau expansion,

$$\begin{aligned} f(s, \theta, h) = & -\frac{2h+1}{2q} - \theta \ln q - hs + \frac{q(q\theta - 1)}{2(q-1)} s^2 - \frac{(q-2)q^3\theta}{6(q-1)^2} s^3 \\ & + \frac{q^3\theta}{12} [1 + (q-1)^{-3}] s^4 - \frac{q^4\theta}{20} [1 - (q-1)^{-4}] s^5 + \frac{q^5\theta}{30} [1 + (q-1)^{-5}] s^6 + \dots \quad . \end{aligned} \quad (7.127)$$

Note that, for $q = 2$, the coefficients of s^3 , s^5 , and higher order odd powers of s vanish in the Landau expansion. This is consistent with what we found for the Ising model, and is related to the \mathbb{Z}_2 symmetry of that model. For $q > 3$, there is a cubic term in the mean field free energy, and thus we generically expect a first order transition, as we shall see below when we discuss Landau theory.

7.5.4 XY Model

Variational density matrix

Consider the so-called XY model, in which each site contains a continuous planar spin, represented by an angular variable $\phi_i \in [-\pi, \pi]$:

$$\hat{H} = -\frac{1}{2} \sum_{i \neq j} J_{ij} \cos(\phi_i - \phi_j) - H \sum_i \cos \phi_i \quad . \quad (7.128)$$

We write the (diagonal elements of the) full density matrix once again as a product:

$$\varrho_N(\phi_1, \phi_2, \dots) = \prod_i \varrho(\phi_i) \quad . \quad (7.129)$$

Our goal will be to extremize the free energy with respect to the *function* $\varrho(\phi)$. To this end, we compute

$$E = \text{Tr}(\varrho_N \hat{H}) = -\frac{1}{2}N\hat{J}(0) \left| \text{Tr}(\varrho e^{i\phi}) \right|^2 - NH \text{Tr}(\varrho \cos \phi) \quad . \quad (7.130)$$

The entropy is $S = -Nk_B \text{Tr}(\varrho \ln \varrho)$. Note that for any function $A(\phi)$, we have¹⁹

$$\text{Tr}(\varrho A) \equiv \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \varrho(\phi) A(\phi) \quad . \quad (7.131)$$

We now extremize the *functional* $F[\varrho(\phi)] = E - TS$ with respect to $\varrho(\phi)$, under the condition that $\text{Tr} \varrho = 1$. We therefore use Lagrange's method of undetermined multipliers, writing

$$F^* = F - Nk_B T \lambda (\text{Tr} \varrho - 1) \quad . \quad (7.132)$$

Note that F^* is a *function* of the Lagrange multiplier λ and a *functional* of the density matrix $\varrho(\phi)$. The prefactor $Nk_B T$ which multiplies λ is of no mathematical consequence – we could always redefine the multiplier to be $\lambda' \equiv Nk_B T \lambda$. It is present only to maintain homogeneity and proper dimensionality of F^* with λ itself dimensionless and of order N^0 . We now have

$$\begin{aligned} \frac{\delta F^*}{\delta \varrho(\phi)} = \frac{\delta}{\delta \varrho(\phi)} \left\{ -\frac{1}{2}N\hat{J}(0) \left| \text{Tr}(\varrho e^{i\phi}) \right|^2 - NH \text{Tr}(\varrho \cos \phi) \right. \\ \left. + Nk_B T \text{Tr}(\varrho \ln \varrho) - Nk_B T \lambda (\text{Tr} \varrho - 1) \right\} \quad . \end{aligned} \quad (7.133)$$

To this end, we note that

$$\frac{\delta}{\delta \varrho(\phi)} \text{Tr}(\varrho A) = \frac{\delta}{\delta \varrho(\phi)} \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \varrho(\phi) A(\phi) = \frac{1}{2\pi} A(\phi) \quad . \quad (7.134)$$

Thus, we have

$$\begin{aligned} \frac{\delta \tilde{F}}{\delta \varrho(\phi)} = -\frac{1}{2}N\hat{J}(0) \cdot \frac{1}{2\pi} \left[\text{Tr}_{\phi'}(\varrho e^{i\phi'}) e^{-i\phi} + \text{Tr}_{\phi'}(\varrho e^{-i\phi'}) e^{i\phi} \right] - NH \cdot \frac{\cos \phi}{2\pi} \\ + Nk_B T \cdot \frac{1}{2\pi} \left[\ln \varrho(\phi) + 1 \right] - Nk_B T \cdot \frac{\lambda}{2\pi} \quad . \end{aligned} \quad (7.135)$$

¹⁹The denominator of 2π in the measure is not necessary, and in fact it is even slightly cumbersome. It divides out whenever we take a ratio to compute a thermodynamic average. I introduce this factor to preserve the relation $\text{Tr} 1 = 1$. I personally find unnormalized traces to be profoundly unsettling on purely aesthetic grounds.

Now let us define

$$\mathrm{Tr}_\phi(\varrho e^{i\phi}) = \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \varrho(\phi) e^{i\phi} \equiv m e^{i\phi_0} \quad . \quad (7.136)$$

We then have

$$\ln \varrho(\phi) = \frac{\hat{J}(0)}{k_B T} m \cos(\phi - \phi_0) + \frac{H}{k_B T} \cos \phi + \lambda - 1. \quad (7.137)$$

Clearly the free energy will be reduced if $\phi_0 = 0$ so that the mean field is maximal and aligns with the external field, which prefers $\phi = 0$. Thus, we conclude

$$\varrho(\phi) = \mathcal{C} \exp\left(\frac{H_{\mathrm{eff}}}{k_B T} \cos \phi\right) \quad , \quad (7.138)$$

where $H_{\mathrm{eff}} = \hat{J}(0) m + H$ and $\mathcal{C} = e^{\lambda-1}$. The value of λ is then determined by invoking the constraint,

$$\mathrm{Tr} \varrho = 1 = \mathcal{C} \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \exp\left(\frac{H_{\mathrm{eff}}}{k_B T} \cos \phi\right) = \mathcal{C} I_0(H_{\mathrm{eff}}/k_B T) \quad , \quad (7.139)$$

where $I_0(z)$ is the Bessel function. We are free to define $\varepsilon \equiv H_{\mathrm{eff}}/k_B T$, and treat ε as our single variational parameter. We then have the normalized single site density matrix

$$\varrho(\phi) = \frac{\exp(\varepsilon \cos \phi)}{\int_{-\pi}^{\pi} \frac{d\phi'}{2\pi} \exp(\varepsilon \cos \phi')} = \frac{\exp(\varepsilon \cos \phi)}{I_0(\varepsilon)} \quad . \quad (7.140)$$

We next compute the following averages:

$$\langle e^{\pm i\phi} \rangle = \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \varrho(\phi) e^{\pm i\phi} = \frac{I_1(\varepsilon)}{I_0(\varepsilon)} \quad (7.141)$$

$$\langle \cos(\phi - \phi') \rangle = \mathrm{Re} \langle e^{i\phi} e^{-i\phi'} \rangle = \left(\frac{I_1(\varepsilon)}{I_0(\varepsilon)}\right)^2 \quad , \quad (7.142)$$

as well as

$$\mathrm{Tr}(\varrho \ln \varrho) = \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \frac{e^{\varepsilon \cos \phi}}{I_0(\varepsilon)} \left\{ \varepsilon \cos \phi - \ln I_0(\varepsilon) \right\} = \varepsilon \frac{I_1(\varepsilon)}{I_0(\varepsilon)} - \ln I_0(\varepsilon) \quad . \quad (7.143)$$

The dimensionless free energy per site is therefore

$$f(\varepsilon, h, \theta) = -\frac{1}{2} \left(\frac{I_1(\varepsilon)}{I_0(\varepsilon)}\right)^2 + (\theta\varepsilon - h) \frac{I_1(\varepsilon)}{I_0(\varepsilon)} - \theta \ln I_0(\varepsilon) \quad , \quad (7.144)$$

with $\theta = k_B T/\hat{J}(0)$ and $h = H/\hat{J}(0)$ and $f = F/N\hat{J}(0)$ as before. Note that the mean field equation is $m = \theta\varepsilon - h = \langle e^{i\phi} \rangle$, i.e.

$$\theta\varepsilon - h = \frac{I_1(\varepsilon)}{I_0(\varepsilon)} \quad . \quad (7.145)$$

For small ε , we may expand the Bessel functions, using

$$I_\nu(z) = \left(\frac{1}{2}z\right)^\nu \sum_{k=0}^{\infty} \frac{\left(\frac{1}{4}z^2\right)^k}{k! \Gamma(k + \nu + 1)} \quad , \quad (7.146)$$

to obtain

$$f(\varepsilon, h, \theta) = \frac{1}{4} \left(\theta - \frac{1}{2}\right) \varepsilon^2 + \frac{1}{64} (2 - 3\theta) \varepsilon^4 - \frac{1}{2} h \varepsilon + \frac{1}{16} h \varepsilon^3 + \dots \quad . \quad (7.147)$$

This predicts a second order phase transition at $\theta_c = \frac{1}{2}$.²⁰ Note also the Curie-Weiss form of the susceptibility at high θ :

$$\frac{\partial f}{\partial \varepsilon} = 0 \quad \implies \quad \varepsilon = \frac{h}{\theta - \theta_c} + \dots \quad . \quad (7.148)$$

Neglect of fluctuations method

Consider again the Hamiltonian of eqn. 7.128. Define $z_i \equiv \exp(i\phi_i)$ and write $z_i = w + \delta z_i$, where $w \equiv \langle z_i \rangle$ and $\delta z_i \equiv z_i - w$. Of course we also have the complex conjugate relations $z_i^* = w^* + \delta z_i^*$ and $w^* = \langle z_i^* \rangle$. Writing $\cos(\phi_i - \phi_j) = \text{Re}(z_i^* z_j)$, by neglecting the terms proportional to $\delta z_i^* \delta z_j$ in \hat{H} we arrive at the mean field Hamiltonian,

$$\hat{H}^{\text{MF}} = \frac{1}{2} N \hat{J}(0) |w|^2 - \frac{1}{2} \hat{J}(0) |w| \sum_i (w^* z_i + w z_i^*) - \frac{1}{2} H \sum_i (z_i^* + z_i) \quad (7.149)$$

It is clear that the free energy will be minimized if the mean field w breaks the O(2) symmetry in the same direction as the external field H , which means $w \in \mathbb{R}$ and

$$\hat{H}^{\text{MF}} = \frac{1}{2} N \hat{J}(0) |w|^2 - (H + \hat{J}(0) |w|) \sum_i \cos \phi_i \quad . \quad (7.150)$$

The dimensionless free energy per site is then

$$f = \frac{1}{2} |w|^2 - \theta \ln I_0 \left(\frac{h + |w|}{\theta} \right) \quad . \quad (7.151)$$

Differentiating with respect to $|w|$, one obtains

$$|w| \equiv m = I_1 \left(\frac{h + m}{\theta} \right) / I_0 \left(\frac{h + m}{\theta} \right) \quad , \quad (7.152)$$

which is the same equation as eqn. 7.145. The two mean field theories yield the same results in every detail (see §7.10).

²⁰Note that the coefficient of the quartic term in ε is negative for $\theta > \frac{2}{3}$. At $\theta = \theta_c = \frac{1}{2}$, the coefficient is positive, but for larger θ one must include higher order terms in the Landau expansion.

7.6 Landau Theory of Phase Transitions

Landau's theory of phase transitions is based on an expansion of the free energy of a thermodynamic system in terms of an *order parameter*, which is nonzero in an ordered phase and zero in a disordered phase. For example, the magnetization M of a ferromagnet in zero external field but at finite temperature typically vanishes for temperatures $T > T_c$, where T_c is the *critical temperature*, also called the *Curie temperature* in a ferromagnet. A low order expansion in powers of the order parameter is appropriate sufficiently close to the phase transition, *i.e.* at temperatures such that the order parameter, if nonzero, is still small.

7.6.1 Quartic free energy with Ising symmetry

The simplest example is the quartic free energy,

$$f(m, h = 0, \theta) = f_0 + \frac{1}{2}am^2 + \frac{1}{4}bm^4 \quad , \quad (7.153)$$

where $f_0 = f_0(\theta)$, $a = a(\theta)$, and $b = b(\theta)$. Here, θ is a dimensionless measure of the temperature²¹. We assume $b > 0$, which is necessary if the free energy is to be bounded from below²².

The equation of state, which relates the intensive quantities m and θ at $h = 0$, is then

$$\frac{\partial f}{\partial m} = 0 = am + bm^3 \quad , \quad (7.154)$$

has three solutions in the complex m plane: (i) $m = 0$, (ii) $m = \sqrt{-a/b}$, and (iii) $m = -\sqrt{-a/b}$. The latter two solutions lie along the (physical) real axis provided $a < 0$. We assume that there exists a unique temperature θ_c where $a(\theta_c) = 0$. Minimizing f , we find

$$\begin{aligned} a < 0 \quad (\theta < \theta_c) & : \quad f(\theta) = f_0 - \frac{a^2}{4b} \\ a > 0 \quad (\theta > \theta_c) & : \quad f(\theta) = f_0 \quad . \end{aligned} \quad (7.155)$$

Thus $a(\theta)$ changes sign at $\theta = \theta_c$, where the free energy is continuous, since $a(\theta_c) = 0$. The specific heat, however, is discontinuous across the transition, with

$$c(\theta_c^+) - c(\theta_c^-) = -\theta_c \left. \frac{\partial^2}{\partial \theta^2} \right|_{\theta=\theta_c} \left(\frac{a^2}{4b} \right) = -\frac{\theta_c [a'(\theta_c)]^2}{2b(\theta_c)} \quad . \quad (7.156)$$

The presence of a magnetic field h breaks the \mathbb{Z}_2 symmetry of $m \rightarrow -m$. The free energy becomes

$$f(m, h, \theta) = f_0 + \frac{1}{2}am^2 + \frac{1}{4}bm^4 - hm \quad , \quad (7.157)$$

²¹For example in an Ising ferromagnet we might define $\theta = k_B T / \hat{J}(0)$, as before.

²²It is always the case that f is bounded from below, on physical grounds. Were b negative, we'd have to consider higher order terms in the Landau expansion.




Figure 7.14: Phase diagram for the quartic Landau free energy $f = f_0 + \frac{1}{2}am^2 + \frac{1}{4}bm^4 - hm$, with $b > 0$. There is a first order line at $h = 0$ extending from $a = -\infty$ and terminating in a critical point at $a = 0$. For $|h| < h^*(a)$ (dashed red line) there are three solutions to the mean field equation, corresponding to one global minimum, one local minimum, and one local maximum. Insets show behavior of the free energy $f(m)$.

and the mean field equation is

$$bm^3 + am - h = 0 \quad . \quad (7.158)$$

This is a cubic equation for m with real coefficients, and as such it can either have three real solutions or one real solution and two complex solutions related by complex conjugation. Clearly we must have $a < 0$ in order to have three real roots, since $bm^3 + am$ is monotonically increasing otherwise. The boundary between these two classes of solution sets occurs when two roots coincide, which means $f''(m) = 0$ as well as $f'(m) = 0$. Simultaneously solving these two equations, we find

$$h^*(a) = \pm \frac{2}{3^{3/2}} \frac{(-a)^{3/2}}{b^{1/2}} \quad , \quad (7.159)$$

or, equivalently,

$$a^*(h) = -\frac{3}{2^{2/3}} b^{1/3} |h|^{2/3}. \quad (7.160)$$

If, for fixed h , we have $a < a^*(h)$, then there will be three real solutions to the mean field equation $f'(m) = 0$, one of which is a global minimum (the one for which $m \cdot h > 0$). For $a > a^*(h)$ there is only a single global minimum, at which m also has the same sign as h . If we solve the mean field equation

perturbatively in h/a , we find

$$\begin{aligned} m(a, h) &= \frac{h}{a} - \frac{b}{a^4} h^3 + \mathcal{O}(h^5) & (a > 0) \\ &= \pm \frac{|a|^{1/2}}{b^{1/2}} + \frac{h}{2|a|} \pm \frac{3b^{1/2}}{8|a|^{5/2}} h^2 + \mathcal{O}(h^3) & (a < 0) . \end{aligned} \quad (7.161)$$

7.6.2 Cubic terms in Landau theory : first order transitions

Next, consider a free energy with a cubic term,

$$f = f_0 + \frac{1}{2}am^2 - \frac{1}{3}ym^3 + \frac{1}{4}bm^4 , \quad (7.162)$$

with $b > 0$ for stability. Without loss of generality, we may assume $y > 0$ (else send $m \rightarrow -m$). Note that we no longer have $m \rightarrow -m$ (*i.e.* \mathbb{Z}_2) symmetry. The cubic term favors positive m . What is the phase diagram in the (a, y) plane?

Extremizing the free energy with respect to m , we obtain

$$\frac{\partial f}{\partial m} = 0 = am - ym^2 + bm^3 . \quad (7.163)$$

This cubic equation factorizes into a linear and quadratic piece, and hence may be solved simply. The three solutions are $m = 0$ and

$$m = m_{\pm} \equiv \frac{y}{2b} \pm \sqrt{\left(\frac{y}{2b}\right)^2 - \frac{a}{b}} . \quad (7.164)$$

We now see that for $y^2 < 4ab$ there is only one real solution, at $m = 0$, while for $y^2 > 4ab$ there are three real solutions. Which solution has lowest free energy? To find out, we compare the energy $f(0)$ with $f(m_+)$ ²³. Thus, we set

$$f(m) = f(0) \implies \frac{1}{2}am^2 - \frac{1}{3}ym^3 + \frac{1}{4}bm^4 = 0 , \quad (7.165)$$

and we now have two quadratic equations to solve simultaneously:

$$0 = a - ym + bm^2 , \quad 0 = \frac{1}{2}a - \frac{1}{3}ym + \frac{1}{4}bm^2 = 0 . \quad (7.166)$$

Eliminating the quadratic term gives $m = 3a/y$. Finally, substituting $m = m_+$ gives us a relation between a , b , and y :

$$y^2 = \frac{9}{2} ab . \quad (7.167)$$

Thus, we have the following:

$$\begin{aligned} a > \frac{y^2}{4b} & : \quad 1 \text{ real root } m = 0 \\ \frac{y^2}{4b} > a > \frac{2y^2}{9b} & : \quad 3 \text{ real roots; minimum at } m = 0 \\ \frac{2y^2}{9b} > a & : \quad 3 \text{ real roots; minimum at } m = \frac{y}{2b} + \sqrt{\left(\frac{y}{2b}\right)^2 - \frac{a}{b}} \end{aligned} \quad (7.168)$$

²³We needn't waste our time considering the $m = m_-$ solution, since the cubic term prefers positive m .




Figure 7.15: Behavior of the quartic free energy $f(m) = \frac{1}{2}am^2 - \frac{1}{3}ym^3 + \frac{1}{4}bm^4$. A: $y^2 < 4ab$; B: $4ab < y^2 < \frac{9}{2}ab$; C and D: $y^2 > \frac{9}{2}ab$. The thick black line denotes a line of first order transitions, where the order parameter is discontinuous across the transition.

The solution $m = 0$ lies at a local minimum of the free energy for $a > 0$ and at a local maximum for $a < 0$. Over the range $y^2/4b > a > 2y^2/9b$, then, there is a global minimum at $m = 0$, a local minimum at $m = m_+$, and a local maximum at $m = m_-$, with $m_+ > m_- > 0$. For $y^2/9b > a > 0$, there is a local minimum at $a = 0$, a global minimum at $m = m_+$, and a local maximum at $m = m_-$, again with $m_+ > m_- > 0$. For $a < 0$, there is a local maximum at $m = 0$, a local minimum at $m = m_-$, and a global minimum at $m = m_+$, with $m_+ > 0 > m_-$. See fig. 7.15.

With $y = 0$, we have a second order transition at $a = 0$. With $y \neq 0$, there is a discontinuous (first order) transition at $a_c = 2y^2/9b > 0$ and $m_c = 2y/3b$. This occurs before a reaches the value $a = 0$ where the curvature at $m = 0$ turns negative. If we write $a = \alpha(T - T_0)$, then the expected second order transition at $T = T_0$ is preempted by a first order transition at $T_c = T_0 + 2y^2/9\alpha b$.

7.6.3 Magnetization dynamics

Suppose we now impose some dynamics on the system, of the simple relaxational type

$$\frac{dm}{dt} = -\Gamma \frac{\partial f}{\partial m} \quad , \quad (7.169)$$

where Γ is a phenomenological kinetic coefficient. Assuming $y > 0$ and $b > 0$, it is convenient to adimensionalize by writing

$$m \equiv \frac{y}{b} \cdot u \quad , \quad a \equiv \frac{y^2}{b} \cdot r \quad , \quad t \equiv \frac{b}{\Gamma y^2} \cdot s \quad . \quad (7.170)$$




Figure 7.16: Fixed points for $\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{3}u^3 + \frac{1}{4}u^4$ and flow under the dynamics $\dot{u} = -\varphi'(u)$. Solid curves represent stable fixed points and dashed curves unstable fixed points. Magenta arrows show behavior under slowly increasing control parameter r and dark blue arrows show behavior under slowly decreasing r . For $u > 0$ there is a hysteresis loop. The thick black curve shows the equilibrium thermodynamic value of $u(r)$, *i.e.* that value which minimizes the free energy $\varphi(u)$. There is a first order phase transition at $r = \frac{2}{9}$, where the thermodynamic value of u jumps from $u = 0$ to $u = \frac{2}{3}$.

Then we obtain

$$\frac{du}{ds} = -\frac{\partial\varphi}{\partial u} \quad , \quad (7.171)$$

where the dimensionless free energy function is

$$\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{3}u^3 + \frac{1}{4}u^4 \quad . \quad (7.172)$$

We see that there is a single control parameter, r . The fixed points of the dynamics are then the stationary points of $\varphi(u)$, where $\varphi'(u) = 0$, with $\varphi'(u) = u(r - u + u^2)$. The solutions to $\varphi'(u) = 0$ are then given by

$$u^* = 0 \quad , \quad u^* = \frac{1}{2} \pm \left(\frac{1}{4} - r\right)^{1/2} \quad . \quad (7.173)$$

For $r > \frac{1}{4}$ there is one fixed point at $u = 0$, which is attractive under the dynamics $\dot{u} = -\varphi'(u)$ since $\varphi''(0) = r$. At $r = \frac{1}{4}$ there occurs a saddle-node bifurcation and a pair of fixed points is generated, one stable and one unstable. As we see from fig. 7.14, the interior fixed point is always unstable and the two exterior fixed points are always stable. At $r = 0$ there is a transcritical bifurcation where two fixed points of opposite stability collide and bounce off one another (metaphorically speaking).

At the saddle-node bifurcation, $r = \frac{1}{4}$ and $u = \frac{1}{2}$, and we find $\varphi(u = \frac{1}{2}; r = \frac{1}{4}) = \frac{1}{192}$, which is positive. Thus, the thermodynamic state of the system remains at $u = 0$ until the value of $\varphi(u_+)$ crosses zero. This occurs when $\varphi(u) = 0$ and $\varphi'(u) = 0$, the simultaneous solution of which yields $r = \frac{2}{9}$ and $u = \frac{2}{3}$.

Suppose we slowly ramp the control parameter r up and down as a function of the dimensionless time s . Under the dynamics of eqn. 7.171, $u(s)$ flows to the first stable fixed point encountered – this is always

the case for a dynamical system with a one-dimensional phase space. Then as r is further varied, u follows the position of whatever locally stable fixed point it initially encountered. Thus, $u(r(s))$ evolves smoothly until a bifurcation is encountered. The situation is depicted by the arrows in fig. 7.16. The equilibrium thermodynamic value for $u(r)$ is discontinuous; there is a first order phase transition at $r = \frac{2}{9}$, as we've already seen. As r is increased, $u(r)$ follows a trajectory indicated by the magenta arrows. For an negative initial value of u , the evolution as a function of r will be *reversible*. However, if $u(0)$ is initially positive, then the system exhibits *hysteresis*, as shown. Starting with a large positive value of r , $u(s)$ quickly evolves to $u = 0^+$, which means a positive infinitesimal value. Then as r is decreased, the system remains at $u = 0^+$ even through the first order transition, because $u = 0$ is an attractive fixed point. However, once r begins to go negative, the $u = 0$ fixed point becomes repulsive, and $u(s)$ quickly flows to the stable fixed point $u_+ = \frac{1}{2} + (\frac{1}{4} - r)^{1/2}$. Further decreasing r , the system remains on this branch. If r is later increased, then $u(s)$ remains on the upper branch past $r = 0$, until the u_+ fixed point annihilates with the unstable fixed point at $u_- = \frac{1}{2} - (\frac{1}{4} - r)^{1/2}$, at which time $u(s)$ quickly flows down to $u = 0^+$ again.

7.6.4 Sixth order Landau theory : tricritical point

Finally, consider a model with \mathbb{Z}_2 symmetry, with the Landau free energy

$$f = f_0 + \frac{1}{2}am^2 + \frac{1}{4}bm^4 + \frac{1}{6}cm^6 \quad , \quad (7.174)$$

with $c > 0$ for stability. We seek the phase diagram in the (a, b) plane. Extremizing f with respect to m ,

$$\frac{\partial f}{\partial m} = 0 = m(a + bm^2 + cm^4) \quad , \quad (7.175)$$

which is a quintic with five solutions over the complex m plane. One solution is obviously $m = 0$. The other four are

$$m = \pm \sqrt{-\frac{b}{2c} \pm \sqrt{\left(\frac{b}{2c}\right)^2 - \frac{a}{c}}} \quad . \quad (7.176)$$

For each \pm symbol in the above equation, there are two options, hence four roots in all.

If $a > 0$ and $b > 0$, then four of the roots are imaginary and there is a unique minimum at $m = 0$.

For $a < 0$, there are only three solutions to $f'(m) = 0$ for real m , since the $-$ choice for the \pm sign under the radical leads to imaginary roots. One of the solutions is $m = 0$. The other two are

$$m = \pm \sqrt{-\frac{b}{2c} + \sqrt{\left(\frac{b}{2c}\right)^2 - \frac{a}{c}}} \quad . \quad (7.177)$$

The most interesting situation is $a > 0$ and $b < 0$. If $a > 0$ and $b < -2\sqrt{ac}$, all five roots are real. There must be three minima, separated by two local maxima. Clearly if m^* is a solution, then so is $-m^*$. Thus, the only question is whether the outer minima are of lower energy than the minimum at $m = 0$. We




Figure 7.17: Behavior of the sextic free energy $f(m) = \frac{1}{2}am^2 + \frac{1}{4}bm^4 + \frac{1}{6}cm^6$. A: $a > 0$ and $b > 0$; B: $a < 0$ and $b > 0$; C: $a < 0$ and $b < 0$; D: $a > 0$ and $b < -\frac{4}{\sqrt{3}}\sqrt{ac}$; E: $a > 0$ and $-\frac{4}{\sqrt{3}}\sqrt{ac} < b < -2\sqrt{ac}$; F: $a > 0$ and $-2\sqrt{ac} < b < 0$. The thick dashed line is a line of second order transitions, which meets the thick solid line of first order transitions at the tricritical point, $(a, b) = (0, 0)$.

assess this by demanding $f(m^*) = f(0)$, where m^* is the position of the largest root (*i.e.* the rightmost minimum). This gives a second quadratic equation,

$$0 = \frac{1}{2}a + \frac{1}{4}bm^2 + \frac{1}{6}cm^4, \quad (7.178)$$

which together with equation 7.175 gives $b = -\frac{4}{\sqrt{3}}\sqrt{ac}$. Thus, we have the following, for fixed $a > 0$:

$$\begin{aligned} b > -2\sqrt{ac} & : 1 \text{ real root } m = 0 \\ -2\sqrt{ac} > b > -\frac{4}{\sqrt{3}}\sqrt{ac} & : 5 \text{ real roots; minimum at } m = 0 \\ -\frac{4}{\sqrt{3}}\sqrt{ac} > b & : 5 \text{ real roots; minima at } m = \pm \sqrt{-\frac{b}{2c} + \sqrt{\left(\frac{b}{2c}\right)^2 - \frac{a}{c}}} \end{aligned} \quad (7.179)$$

The point $(a, b) = (0, 0)$, which lies at the confluence of a first order line and a second order line, is known as a *tricritical point*.




Figure 7.18: Free energy $\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{4}u^4 + \frac{1}{6}u^6$ for different values of the control parameter r .

7.6.5 Hysteresis for the sextic potential

Once again, we consider the dissipative dynamics $\dot{m} = -\Gamma f'(m)$. We adimensionalize by writing

$$m \equiv \sqrt{\frac{|b|}{c}} \cdot u \quad , \quad a \equiv \frac{b^2}{c} \cdot r \quad , \quad t \equiv \frac{c}{\Gamma b^2} \cdot s \quad . \quad (7.180)$$

Then we obtain once again the dimensionless equation $du/ds = -\partial\varphi/\partial u$, where

$$\varphi(u) = \frac{1}{2}ru^2 \pm \frac{1}{4}u^4 + \frac{1}{6}u^6 \quad . \quad (7.181)$$

In the above equation, the coefficient of the quartic term is positive if $b > 0$ and negative if $b < 0$. That is, the coefficient is $\text{sgn}(b)$. When $b > 0$ we can ignore the sextic term for sufficiently small u , and we recover the quartic free energy studied earlier. There is then a second order transition at $r = 0$. The free energy curves for various values of r are plotted in fig. 7.18.

New and interesting behavior occurs for $b > 0$. The fixed points of the dynamics are obtained by setting $\varphi'(u) = 0$. We have

$$\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{4}u^4 + \frac{1}{6}u^6 \quad , \quad \varphi'(u) = u(r - u^2 + u^4) \quad . \quad (7.182)$$

Thus, the equation $\varphi'(u) = 0$ factorizes into a linear factor u and a quartic factor $u^4 - u^2 + r$ which is quadratic in u^2 . Thus, we can easily obtain the roots:

$$\begin{aligned} r < 0 & : \quad u^* = 0 \quad , \quad u^* = \pm \sqrt{\frac{1}{2} + \sqrt{\frac{1}{4} - r}} \\ 0 < r < \frac{1}{4} & : \quad u^* = 0 \quad , \quad u^* = \pm \sqrt{\frac{1}{2} + \sqrt{\frac{1}{4} - r}} \quad , \quad u^* = \pm \sqrt{\frac{1}{2} - \sqrt{\frac{1}{4} - r}} \\ r > \frac{1}{4} & : \quad u^* = 0 \quad . \end{aligned} \quad (7.183)$$

In fig. 7.19, we plot the fixed points and the hysteresis loops for this system. At $r = \frac{1}{4}$, there are two symmetrically located saddle-node bifurcations at $u = \pm \frac{1}{\sqrt{2}}$. We find $\varphi(u = \pm \frac{1}{\sqrt{2}}, r = \frac{1}{4}) = \frac{1}{48}$, which is positive, indicating that the stable fixed point $u^* = 0$ remains the thermodynamic minimum for the free energy $\varphi(u)$ as r is decreased through $r = \frac{1}{4}$. Setting $\varphi(u) = 0$ and $\varphi'(u) = 0$ simultaneously, we obtain $r = \frac{3}{16}$ and $u = \pm \frac{\sqrt{3}}{2}$. The thermodynamic value for u therefore jumps discontinuously from $u = 0$ to $u = \pm \frac{\sqrt{3}}{2}$ (either branch) at $r = \frac{3}{16}$; this is a first order transition.

Under the dissipative dynamics considered here, the system exhibits hysteresis, as indicated in the figure, where the arrows show the evolution of $u(s)$ for very slowly varying $r(s)$. When the control parameter r is large and positive, the flow is toward the sole fixed point at $u^* = 0$. At $r = \frac{1}{4}$, two simultaneous saddle-node bifurcations take place at $u^* = \pm \frac{1}{\sqrt{2}}$; the outer branch is stable and the inner branch unstable in both cases. At $r = 0$ there is a subcritical pitchfork bifurcation, and the fixed point at $u^* = 0$ becomes unstable.

Suppose one starts off with $r \gg \frac{1}{4}$ with some value $u > 0$. The flow $\dot{u} = -\varphi'(u)$ then rapidly results in $u \rightarrow 0^+$. This is the 'high temperature phase' in which there is no magnetization. Now let r increase slowly, using s as the dimensionless time variable. The scaled magnetization $u(s) = u^*(r(s))$ will remain pinned at the fixed point $u^* = 0^+$. As r passes through $r = \frac{1}{4}$, two new stable values of u^* appear, but our system remains at $u = 0^+$, since $u^* = 0$ is a stable fixed point. But after the subcritical pitchfork, $u^* = 0$ becomes unstable. The magnetization $u(s)$ then flows rapidly to the stable fixed point at $u^* = \frac{1}{\sqrt{2}}$, and follows the curve $u^*(r) = (\frac{1}{2} + (\frac{1}{4} - r)^{1/2})^{1/2}$ for all $r < 0$.

Now suppose we start increasing r (i.e. increasing temperature). The magnetization follows the stable fixed point $u^*(r) = (\frac{1}{2} + (\frac{1}{4} - r)^{1/2})^{1/2}$ past $r = 0$, beyond the first order phase transition point at $r = \frac{3}{16}$, and all the way up to $r = \frac{1}{4}$, at which point this fixed point is annihilated at a saddle-node bifurcation. The flow then rapidly takes $u \rightarrow u^* = 0^+$, where it remains as r continues to be increased further. Within the region $r \in [0, \frac{1}{4}]$ of control parameter space, the dynamics are said to be *irreversible* and the behavior of $u(s)$ is said to be *hysteretic*.

7.7 Mean Field Theory of Fluctuations

7.7.1 Correlation and response in mean field theory

We now consider the matter of correlation and response functions within mean field theory. A correlation function is a thermodynamic average of fields at various sites, such as $\langle \sigma_i \sigma_j \rangle$ (two-point correlation function), $\langle \sigma_i \sigma_j \sigma_k \sigma_l \rangle$ (four-point correlation function), etc. A response function describes the influence of external fields on thermodynamic averages, such as $\partial m_i / \partial H_j$, the variation of the local magnetization $m_i = \langle \sigma_i \rangle$ at site i due to the presence of a local field H_j at site j . We shall see that there is a direct relation between correlation and response functions.

Consider the Ising model,

$$\hat{H} = -\frac{1}{2} \sum_{i,j} J_{ij} \sigma_i \sigma_j - \sum_k H_k \sigma_k \quad , \quad (7.184)$$




Figure 7.19: Fixed points $\varphi'(u^*) = 0$ for the sextic potential $\varphi(u) = \frac{1}{2}ru^2 - \frac{1}{4}u^4 + \frac{1}{6}u^6$, and dynamical flow (arrows) under $\dot{u} = -\varphi'(u)$. Solid curves show stable fixed points and dashed curves show unstable fixed points. The thick solid black and solid grey curves indicate the equilibrium thermodynamic values for u ; note the overall $u \rightarrow -u$ symmetry. Within the region $r \in [0, \frac{1}{4}]$ the dynamics are irreversible and the system exhibits the phenomenon of hysteresis. There is a first order phase transition at $r = \frac{3}{16}$.

where the local magnetic field on site k is now H_k . We assume without loss of generality that the diagonal terms vanish: $J_{ii} = 0$. Now consider the partition function $Z = \text{Tr} e^{-\beta\hat{H}}$ as a function of the temperature T and the local field values $\{H_i\}$. We have

$$\begin{aligned} \frac{\partial Z}{\partial H_i} &= \beta \text{Tr} [\sigma_i e^{-\beta\hat{H}}] = \beta Z \cdot \langle \sigma_i \rangle \\ \frac{\partial^2 Z}{\partial H_i \partial H_j} &= \beta^2 \text{Tr} [\sigma_i \sigma_j e^{-\beta\hat{H}}] = \beta^2 Z \cdot \langle \sigma_i \sigma_j \rangle \quad . \end{aligned} \quad (7.185)$$

Thus,

$$\begin{aligned} m_i &= -\frac{\partial F}{\partial H_i} = \langle \sigma_i \rangle \\ \chi_{ij} &= \frac{\partial m_i}{\partial H_j} = -\frac{\partial^2 F}{\partial H_i \partial H_j} = \frac{1}{k_B T} \cdot \left\{ \langle \sigma_i \sigma_j \rangle - \langle \sigma_i \rangle \langle \sigma_j \rangle \right\} \quad . \end{aligned} \quad (7.186)$$

Expressions such as $\langle \sigma_i \rangle$, $\langle \sigma_i \sigma_j \rangle$, etc. are in general called *correlation functions*. For example, we define the *spin-spin correlation function* C_{ij} as

$$C_{ij} \equiv \langle \sigma_i \sigma_j \rangle - \langle \sigma_i \rangle \langle \sigma_j \rangle \quad . \quad (7.187)$$

Expressions such as $\partial F / \partial H_i$ and $\partial^2 F / \partial H_i \partial H_j$ are called *response functions*. The above relation between correlation functions and response functions, $C_{ij} = k_B T \chi_{ij}$, is valid *only for the equilibrium distribution*.

In particular, this relationship is *invalid* if one uses an approximate distribution, such as the variational density matrix formalism of mean field theory.

The question then arises: within mean field theory, which is more accurate: correlation functions or response functions? A simple argument suggests that the *response functions* are more accurate representations of the real physics. To see this, let's write the variational density matrix ϱ^{var} as the sum of the exact equilibrium (Boltzmann) distribution $\varrho^{\text{eq}} = Z^{-1} \exp(-\beta\hat{H})$ plus a deviation $\delta\varrho$:

$$\varrho^{\text{var}} = \varrho^{\text{eq}} + \delta\varrho \quad . \quad (7.188)$$

Then if we calculate a correlator using the variational distribution, we have

$$\langle \sigma_i \sigma_j \rangle_{\text{var}} = \text{Tr} \left[\varrho^{\text{var}} \sigma_i \sigma_j \right] = \text{Tr} \left[\varrho^{\text{eq}} \sigma_i \sigma_j \right] + \text{Tr} \left[\delta\varrho \sigma_i \sigma_j \right] \quad . \quad (7.189)$$

Thus, the variational density matrix gets the correlator right to first order in $\delta\varrho$. On the other hand, the free energy is given by

$$F^{\text{var}} = F^{\text{eq}} + \sum_{\sigma} \left. \frac{\partial F}{\partial \varrho_{\sigma}} \right|_{\varrho^{\text{eq}}} \delta\varrho_{\sigma} + \frac{1}{2} \sum_{\sigma, \sigma'} \left. \frac{\partial^2 F}{\partial \varrho_{\sigma} \partial \varrho_{\sigma'}} \right|_{\varrho^{\text{eq}}} \delta\varrho_{\sigma} \delta\varrho_{\sigma'} + \dots \quad . \quad (7.190)$$

Here σ denotes a state of the system, *i.e.* $|\sigma\rangle = |\sigma_1, \dots, \sigma_N\rangle$, where every spin polarization is specified. Since the free energy is an extremum (and in fact an absolute minimum) with respect to the distribution, the second term on the RHS vanishes. This means that the free energy is accurate to second order in the deviation $\delta\varrho$.

7.7.2 Calculation of the response functions

Consider the variational density matrix $\varrho(\sigma) = \prod_i \varrho_i(\sigma_i)$, where

$$\varrho_i(\sigma_i) = \left(\frac{1+m_i}{2} \right) \delta_{\sigma_i, 1} + \left(\frac{1-m_i}{2} \right) \delta_{\sigma_i, -1} \quad . \quad (7.191)$$

The variational energy $E = \text{Tr}(\varrho \hat{H})$ is

$$E = -\frac{1}{2} \sum_{ij} J_{i,j} m_i m_j - \sum_i H_i m_i \quad (7.192)$$

and the entropy $S = -k_B T \text{Tr}(\varrho \ln \varrho)$ is

$$S = -k_B \sum_i \left\{ \left(\frac{1+m_i}{2} \right) \ln \left(\frac{1+m_i}{2} \right) + \left(\frac{1-m_i}{2} \right) \ln \left(\frac{1-m_i}{2} \right) \right\} \quad . \quad (7.193)$$

Setting the variation $\frac{\partial F}{\partial m_i} = 0$, with $F = E - TS$, we obtain the mean field equations,

$$m_i = \tanh(\beta J_{ij} m_j + \beta H_i) \quad , \quad (7.194)$$

where we use the summation convention: $J_{ij} m_j \equiv \sum_j J_{ij} m_j$. Suppose $T > T_c$ and m_i is small. Then we can expand the RHS of the above mean field equations, obtaining

$$(\delta_{ij} - \beta J_{ij}) m_j = \beta H_i \quad . \quad (7.195)$$

Thus, the susceptibility tensor χ is the inverse of the matrix $(k_B T \cdot \mathbb{I} - \mathbb{J})$:

$$\chi_{ij} = \frac{\partial m_i}{\partial H_j} = (k_B T \cdot \mathbb{I} - \mathbb{J})_{ij}^{-1} \quad , \quad (7.196)$$

where \mathbb{I} is the identity. Note also that so-called *connected averages* of the kind in eqn. 7.187 vanish identically if we compute them using our variational density matrix, since all the sites are independent, hence

$$\langle \sigma_i \sigma_j \rangle = \text{Tr}(\varrho^{\text{var}} \sigma_i \sigma_j) = \text{Tr}(\varrho_i \sigma_i) \cdot \text{Tr}(\varrho_j \sigma_j) = \langle \sigma_i \rangle \cdot \langle \sigma_j \rangle \quad , \quad (7.197)$$

and therefore $\chi_{ij} = 0$ if we compute the correlation functions themselves from the variational density matrix, rather than from the free energy F . As we have argued above, the latter approximation is more accurate.

Assuming $J_{ij} = J(\mathbf{R}_i - \mathbf{R}_j)$, where \mathbf{R}_i is a Bravais lattice site, we can Fourier transform the above equation, resulting in

$$\hat{m}(\mathbf{q}) = \frac{\hat{H}(\mathbf{q})}{k_B T - \hat{J}(\mathbf{q})} \equiv \hat{\chi}(\mathbf{q}) \hat{H}(\mathbf{q}) \quad . \quad (7.198)$$

Once again, our definition of lattice Fourier transform of a function $\phi(\mathbf{R})$ is

$$\hat{\phi}(\mathbf{q}) \equiv \sum_{\mathbf{R}} \phi(\mathbf{R}) e^{-i\mathbf{q} \cdot \mathbf{R}} \quad , \quad \phi(\mathbf{R}) = \Omega \int_{\hat{\Omega}} \frac{d^d q}{(2\pi)^d} \hat{\phi}(\mathbf{q}) e^{i\mathbf{q} \cdot \mathbf{R}} \quad , \quad (7.199)$$

where Ω is the unit cell in real space, called the *Wigner-Seitz cell*, and $\hat{\Omega}$ is the first Brillouin zone, which is the unit cell in *reciprocal space*. Similarly, we have

$$\begin{aligned} \hat{J}(\mathbf{q}) &= \sum_{\mathbf{R}} J(\mathbf{R}) \left(1 - i\mathbf{q} \cdot \mathbf{R} - \frac{1}{2}(\mathbf{q} \cdot \mathbf{R})^2 + \dots \right) \\ &= \hat{J}(0) \cdot \left\{ 1 - q^2 R_*^2 + \mathcal{O}(q^4) \right\} \quad , \end{aligned} \quad (7.200)$$

where

$$R_*^2 = \frac{\sum_{\mathbf{R}} \mathbf{R}^2 J(\mathbf{R})}{2d \sum_{\mathbf{R}} J(\mathbf{R})} \quad . \quad (7.201)$$

Here we have assumed inversion symmetry for the lattice, in which case

$$\sum_{\mathbf{R}} R^\mu R^\nu J(\mathbf{R}) = \frac{1}{d} \cdot \delta^{\mu\nu} \sum_{\mathbf{R}} \mathbf{R}^2 J(\mathbf{R}) \quad . \quad (7.202)$$

On cubic lattices with nearest neighbor interactions only, one has $R_* = a/\sqrt{2d}$, where a is the lattice constant and d is the dimension of space.

Thus, with the identification $k_B T_c = \hat{J}(0)$, we have

$$\hat{\chi}(\mathbf{q}) = \frac{1}{k_B(T - T_c) + k_B T_c R_*^2 q^2 + \mathcal{O}(q^4)} = \frac{1}{k_B T_c R_*^2} \cdot \frac{1}{\xi^{-2} + q^2 + \mathcal{O}(q^4)} \quad , \quad (7.203)$$

where

$$\xi(T) = R_* \cdot \left(\frac{T - T_c}{T_c} \right)^{-1/2} \quad (7.204)$$

is the *correlation length*. With the definition $\xi(T) \propto |T - T_c|^{-\nu}$ as $T \rightarrow T_c$, we obtain the mean field correlation length exponent $\nu = \frac{1}{2}$. The exact result for the two-dimensional Ising model is $\nu = 1$, whereas $\nu \approx 0.6$ for the $d = 3$ Ising model. Note that $\hat{\chi}(\mathbf{q} = 0, T)$ diverges as $(T - T_c)^{-1}$ for $T > T_c$.

In real space, we have

$$m_i = \sum_j \chi_{ij} H_j \quad , \quad \chi_{ij} = \Omega \int \frac{d^d q}{(2\pi)^d} \hat{\chi}(\mathbf{q}) e^{i\mathbf{q} \cdot (\mathbf{R}_i - \mathbf{R}_j)} \quad . \quad (7.205)$$

Note that $\hat{\chi}(\mathbf{q})$ is properly periodic under $\mathbf{q} \rightarrow \mathbf{q} + \mathbf{G}$, where \mathbf{G} is a reciprocal lattice vector, which satisfies $e^{i\mathbf{G} \cdot \mathbf{R}} = 1$ for any direct Bravais lattice vector \mathbf{R} . Indeed, we have

$$\hat{\chi}^{-1}(\mathbf{q}) = k_B T - \hat{J}(\mathbf{q}) = k_B T - J \sum_{\delta} e^{i\mathbf{q} \cdot \delta} \quad , \quad (7.206)$$

where δ is a nearest neighbor separation vector, and where in the second line we have assumed nearest neighbor interactions only. On cubic lattices in d dimensions, there are $2d$ nearest neighbor separation vectors, $\delta = \pm a \hat{\mathbf{e}}_\mu$, where $\mu \in \{1, \dots, d\}$. The real space susceptibility is then

$$\chi(\mathbf{R}, H = 0) = \int_{-\pi}^{\pi} \frac{d\theta_1}{2\pi} \dots \int_{-\pi}^{\pi} \frac{d\theta_d}{2\pi} \frac{e^{in_1\theta_1} \dots e^{in_d\theta_d}}{k_B T - (2J \cos \theta_1 + \dots + 2J \cos \theta_d)} \quad , \quad (7.207)$$

where $\mathbf{R} = a \sum_{\mu=1}^d n_\mu \hat{\mathbf{e}}_\mu$ is a general direct lattice vector for the cubic Bravais lattice in d dimensions, and the $\{n_\mu\}$ are integers.

The long distance behavior was discussed in chapter 6 (see §6.5.7 on Ornstein-Zernike theory²⁴). For convenience we reiterate those results:

- In $d = 1$,

$$\chi_{d=1}(x) = \left(\frac{\xi}{2k_B T_c R_*^2} \right) e^{-|x|/\xi} \quad . \quad (7.208)$$

²⁴There is a sign difference between the particle susceptibility defined in chapter 6 and the spin susceptibility defined here. The origin of the difference is that the single particle potential v as defined was repulsive for $v > 0$, meaning the local density response δn should be negative, while in the current discussion a positive magnetic field H prefers $m > 0$.

- In $d > 1$, with $r \rightarrow \infty$ and ξ fixed,

$$\chi_d^{\text{OZ}}(\mathbf{r}) \simeq C_d \cdot \frac{\xi^{(3-d)/2}}{k_{\text{B}} T R_*^2} \cdot \frac{e^{-r/\xi}}{r^{(d-1)/2}} \cdot \left\{ 1 + \mathcal{O}\left(\frac{d-3}{r/\xi}\right) \right\} , \quad (7.209)$$

where the C_d are dimensionless constants.

- In $d > 2$, with $\xi \rightarrow \infty$ and r fixed (i.e. $T \rightarrow T_c$ at fixed separation r),

$$\chi_d(\mathbf{r}) \simeq \frac{C'_d}{k_{\text{B}} T R_*^2} \cdot \frac{e^{-r/\xi}}{r^{d-2}} \cdot \left\{ 1 + \mathcal{O}\left(\frac{d-3}{r/\xi}\right) \right\} . \quad (7.210)$$

In $d = 2$ dimensions we obtain

$$\chi_{d=2}(\mathbf{r}) \simeq \frac{C'_2}{k_{\text{B}} T R_*^2} \cdot \ln\left(\frac{r}{\xi}\right) e^{-r/\xi} \cdot \left\{ 1 + \mathcal{O}\left(\frac{1}{\ln(r/\xi)}\right) \right\} , \quad (7.211)$$

where the C'_d are dimensionless constants.

Close to the critical point the spatial dependence of the two-point correlation $C(\mathbf{r}, T) = k_{\text{B}} T \chi(\mathbf{r}, T)$ is given by

$$C(\mathbf{r}, T) = r^{-(d-2+\eta)} \phi(r/\xi(T)) , \quad (7.212)$$

where η is the *anomalous critical exponent* and $\phi(r/\xi)$ is a *scaling function*. The condition $T \approx T_c$ means that $\xi(T) \gg a$, where a is a microscopic length, such as a lattice constant²⁵. Thus, we've encountered in this section two additional critical exponents, the anomalous exponent η and the correlation length exponent ν , which we first met in §7.3.2, whose mean field values are $\eta = 0$ and $\nu = \frac{1}{2}$.

7.7.3 Beyond the Ising model

Consider a general spin model, and a variational density matrix ϱ_{var} which is a product of single site density matrices:

$$\varrho_{\text{var}}[\{\mathbf{S}_i\}] = \prod_i \varrho_1^{(i)}(\mathbf{S}_i) , \quad (7.213)$$

where $\text{Tr}(\varrho_{\text{var}} \mathbf{S}) = \mathbf{m}_i$ is the local magnetization and \mathbf{S}_i , which may be a scalar (e.g., σ_i in the Ising model previously discussed), is the local spin operator. Note that $\varrho_1^{(i)}(\mathbf{S}_i)$ depends parametrically on the variational parameter(s) \mathbf{m}_i . Let the Hamiltonian be

$$\hat{H} = -\frac{1}{2} \sum_{i,j} J_{ij}^{\mu\nu} S_i^\mu S_j^\nu + \sum_i h(\mathbf{S}_i) - \sum_i \mathbf{H}_i \cdot \mathbf{S}_i . \quad (7.214)$$

The variational free energy is then

$$F_{\text{var}} = -\frac{1}{2} \sum_{i,j} J_{ij}^{\mu\nu} m_i^\mu m_j^\nu + \sum_i \varphi(\mathbf{m}_i, T) - \sum_i \mathbf{H}_i \cdot \mathbf{m}_i , \quad (7.215)$$

²⁵The scaling functions on the high and low temperature sides of the transition may be different, and are denoted as $\phi_{\pm}(r/\xi)$, respectively.

where the single site free energy $\varphi(\mathbf{m}_i, T)$ in the absence of an external field is given by

$$\varphi(\mathbf{m}_i, T) = \text{Tr} \left[\varrho_1^{(i)}(\mathbf{S}) h(\mathbf{S}) \right] + k_B T \text{Tr} \left[\varrho_1^{(i)}(\mathbf{S}) \ln \varrho_1^{(i)}(\mathbf{S}) \right] \quad (7.216)$$

We then have

$$\frac{\partial F_{\text{var}}}{\partial m_i^\mu} = - \sum_j J_{ij}^{\mu\nu} m_j^\nu - H_i^\mu + \frac{\partial \varphi(\mathbf{m}_i, T)}{\partial m_i^\mu} \quad (7.217)$$

For the noninteracting system, we have $J_{ij}^{\mu\nu} = 0$, and the weak field response must be linear. In this limit we may write $m_i^\mu = \chi_{\mu\nu}^0(T) H_i^\nu + \mathcal{O}(H_i^3)$, and we conclude

$$\frac{\partial \varphi(\mathbf{m}_i, T)}{\partial m_i^\mu} = [\chi^0(T)]_{\mu\nu}^{-1} m_i^\nu + \mathcal{O}(m_i^3) \quad (7.218)$$

Note that this entails the following expansion for the single site free energy in zero field:

$$\varphi(\mathbf{m}_i, T) = \frac{1}{2} [\chi^0(T)]_{\mu\nu}^{-1} m_i^\nu m_i^\nu + \mathcal{O}(m_i^4) \quad (7.219)$$

Finally, we restore the interaction term and extremize F_{var} by setting $\partial F_{\text{var}} / \partial m_i^\mu = 0$. To linear order, then,

$$m_i^\mu = \chi_{\mu\nu}^0(T) \left(H_i^\nu + \sum_j J_{ij}^{\nu\lambda} m_j^\lambda \right) \quad (7.220)$$

Typically the local susceptibility is a scalar in the internal spin space, *i.e.* $\chi_{\mu\nu}^0(T) = \chi^0(T) \delta_{\mu\nu}$, in which case we obtain

$$(\delta^{\mu\nu} \delta_{ij} - \chi^0(T) J_{ij}^{\mu\nu}) m_i^\nu = \chi^0(T) H_i^\mu \quad (7.221)$$

In Fourier space, then,

$$\hat{\chi}_{\mu\nu}(\mathbf{q}, T) = \chi^0(T) \left(1 - \chi^0(T) \hat{\mathbb{J}}(\mathbf{q}) \right)_{\mu\nu}^{-1}, \quad (7.222)$$

where $\hat{\mathbb{J}}(\mathbf{q})$ is the matrix whose elements are $\hat{J}^{\mu\nu}(\mathbf{q})$. If $\hat{J}^{\mu\nu}(\mathbf{q}) = \hat{J}(\mathbf{q}) \delta^{\mu\nu}$, then the susceptibility is isotropic in spin space, with

$$\hat{\chi}(\mathbf{q}, T) = \frac{1}{[\chi^0(T)]^{-1} - \hat{J}(\mathbf{q})} \quad (7.223)$$

Consider now the following illustrative examples:

- (i) Quantum spin S with $h(\mathbf{S}) = 0$: We take the \hat{z} axis to be that of the local external magnetic field, *i.e.* \hat{H}_i . Write $\varrho_1(\mathbf{S}) = z^{-1} \exp(uS^z/k_B T)$, where $u = u(m, T)$ is obtained implicitly from the relation $m(u, T) = \text{Tr}(\varrho_1 S^z)$. The normalization constant is

$$z = \text{Tr} e^{uS^z/k_B T} = \sum_{j=-S}^S e^{ju/k_B T} = \frac{\sinh[(S + \frac{1}{2})u/k_B T]}{\sinh[u/2k_B T]} \quad (7.224)$$

The relation between m , u , and T is then given by

$$\begin{aligned} m = \langle S^z \rangle &= k_B T \frac{\partial \ln z}{\partial u} = (S + \frac{1}{2}) \text{ctnh}[(S + \frac{1}{2})u/k_B T] - \frac{1}{2} \text{ctnh}[u/2k_B T] \\ &= \frac{S(S+1)}{3k_B T} u + \mathcal{O}(u^3) \quad (7.225) \end{aligned}$$

The free-field single-site free energy is then

$$\varphi(\mathbf{m}, T) = k_B T \operatorname{Tr}(\varrho_1 \ln \varrho_1) = um - k_B T \ln z \quad , \quad (7.226)$$

whence

$$\frac{\partial \varphi}{\partial m} = u + m \frac{\partial u}{\partial m} - k_B T \frac{\partial \ln z}{\partial u} \frac{\partial u}{\partial m} = u \equiv \chi_0^{-1}(T) m + \mathcal{O}(m^3) \quad , \quad (7.227)$$

and we thereby obtain the result

$$\chi_0(T) = \frac{S(S+1)}{3k_B T} \quad , \quad (7.228)$$

which is the Curie susceptibility.

- (ii) Classical spin $\mathbf{S} = S \hat{\mathbf{n}}$ with $h = 0$ and $\hat{\mathbf{n}}$ an N -component unit vector : We take the single site density matrix to be $\varrho_1(\mathbf{S}) = z^{-1} \exp(\mathbf{u} \cdot \mathbf{S}/k_B T)$. The single site field-free partition function is then

$$z = \int \frac{d\hat{\mathbf{n}}}{\Omega_N} \exp(\mathbf{u} \cdot \mathbf{S}/k_B T) = 1 + \frac{S^2 \mathbf{u}^2}{N(k_B T)^2} + \mathcal{O}(u^4) \quad , \quad (7.229)$$

with $\Omega_N = 2\pi^{N/2}/\Gamma(N/2)$ the total solid angle, and therefore

$$\mathbf{m} = k_B T \frac{\partial \ln z}{\partial \mathbf{u}} = \frac{S^2 \mathbf{u}}{Nk_B T} + \mathcal{O}(u^3) \quad , \quad (7.230)$$

from which we read off $\chi_0(T) = S^2/Nk_B T$. Note that this agrees in the classical ($S \rightarrow \infty$) limit, for $N = 3$, with our previous result.

- (iii) Quantum spin S with $h(\mathbf{S}) = \Delta(S^z)^2$: This corresponds to so-called *easy plane anisotropy*, meaning that the single site energy $h(\mathbf{S})$ is minimized when the local spin vector \mathbf{S} lies in the (x, y) plane. As in example (i), we write $\varrho_1(\mathbf{S}) = z^{-1} \exp(uS^z/k_B T)$, yielding the same expression for z and the same relation between z and u . What is different is that we must evaluate the local energy,

$$\begin{aligned} e(u, T) &= \operatorname{Tr}(\varrho_1 h(\mathbf{S})) = \Delta (k_B T)^2 \frac{\partial^2 \ln z}{\partial u^2} \\ &= \frac{\Delta}{4} \left[\frac{1}{\sinh^2[u/2k_B T]} - \frac{(2S+1)^2}{\sinh^2[(2S+1)u/2k_B T]} \right] = \frac{S(S+1)\Delta u^2}{6(k_B T)^2} + \mathcal{O}(u^4) \quad . \end{aligned} \quad (7.231)$$

We now have $\varphi = e + um - k_B T \ln z$, from which we obtain the susceptibility

$$\chi^0(T) = \frac{S(S+1)}{3(k_B T + \Delta)} \quad . \quad (7.232)$$

Note that the local susceptibility no longer diverges as $T \rightarrow 0$, because there is always a gap in the spectrum of $h(\mathbf{S})$.

7.8 Global Symmetries

7.8.1 Symmetries and symmetry groups

Interacting systems can be broadly classified according to their *global symmetry group*. Consider the following five examples:

$$\begin{aligned}
 \hat{H}_{\text{Ising}} &= - \sum_{i < j} J_{ij} \sigma_i \sigma_j & \sigma_i &\in \{-1, +1\} \\
 \hat{H}_{p\text{-clock}} &= - \sum_{i < j} J_{ij} \cos\left(\frac{2\pi(n_i - n_j)}{p}\right) & n_i &\in \{1, 2, \dots, p\} \\
 \hat{H}_{q\text{-Potts}} &= - \sum_{i < j} J_{ij} \delta_{\sigma_i, \sigma_j} & \sigma_i &\in \{1, 2, \dots, q\} \\
 \hat{H}_{XY} &= - \sum_{i < j} J_{ij} \cos(\phi_i - \phi_j) & \phi_i &\in [0, 2\pi] \\
 \hat{H}_{\text{O}(n)} &= - \sum_{i < j} J_{ij} \hat{\omega}_i \cdot \hat{\omega}_j & \hat{\omega}_i &\in S^{n-1} \quad .
 \end{aligned} \tag{7.233}$$

The Ising Hamiltonian is left invariant by the global symmetry group \mathbb{Z}_2 . This group has two elements, which we write as \mathbb{I} and r . \mathbb{I} is the identity operation, *i.e.* $\mathbb{I}\sigma_i = \sigma_i$ for all i . The r operation reverses the spins: $r\sigma_i = -\sigma_i$. By simultaneously reversing *all* the spins $\sigma_i \rightarrow -\sigma_i$, the interactions remain invariant. Note that $r^2 = \mathbb{I}$.

The degrees of freedom of the p -state clock model are integer variables n_i each of which ranges from 1 to p . The Hamiltonian is invariant under the discrete group \mathbb{Z}_p , whose p elements are generated by the single operation r , where

$$r n_i = \begin{cases} n_i + 1 & \text{if } n_i \in \{1, 2, \dots, p-1\} \\ 1 & \text{if } n_i = p \quad . \end{cases} \tag{7.234}$$

Think of a clock with one hand and p ‘hour’ markings consecutively spaced by an angle $2\pi/p$. In each site i , a hand points to one of the p hour marks; this determines n_i . The operation r simply advances *all* the hours by one tick, with hour p advancing to hour 1, just as 23:00 military time is followed one hour later by 00:00. The interaction $\cos(2\pi(n_i - n_j)/p)$ is invariant under such an operation. The p elements of the group \mathbb{Z}_p are then $\mathbb{I}, r, r^2, \dots, r^{p-1}$.

We’ve already met up with the q -state Potts model, where each site supports a ‘spin’ σ_i which can be in any of q possible states, which we may label by integers $\{1, \dots, q\}$. The energy of two interacting sites i and j is $-J_{ij}$ if $\sigma_i = \sigma_j$ and zero otherwise. This energy function is invariant under global operations of the *symmetric group on q characters*, S_q , which is the group of permutations of the sequence $\{1, 2, 3, \dots, q\}$. The group S_q has $q!$ elements. Note the difference between a \mathbb{Z}_q symmetry and an S_q symmetry. In the former case, the Hamiltonian is invariant only under the q -element cyclic permutations, *e.g.*

$$r \equiv \begin{pmatrix} 1 & 2 & \dots & q-1 & q \\ 2 & 3 & \dots & q & 1 \end{pmatrix}$$




Figure 7.20: A domain wall in a one-dimensional Ising model.

and its powers r^l with $l = 0, \dots, q - 1$.

All these models – the Ising, p -state clock, and q -state Potts models – possess a global symmetry group which is *discrete*. That is, each of the symmetry groups $\mathbb{Z}_2, \mathbb{Z}_p, S_q$ is a discrete group, with a finite number of elements. The XY Hamiltonian \hat{H}_{XY} on the other hand is invariant under a *continuous* group of transformations $\phi_i \rightarrow \phi_i + \alpha$, where ϕ_i is the angle variable on site i . More to the point, we could write the interaction term $\cos(\phi_i - \phi_j)$ as $\frac{1}{2}(z_i^* z_j + z_i z_j^*)$, where $z_i = e^{i\phi_i}$ is a phase which lives on the unit circle, and z_i^* is the complex conjugate of z_i . The model is then invariant under the global transformation $z_i \rightarrow e^{i\alpha} z_i$. The phases $e^{i\alpha}$ form a group under multiplication, called $U(1)$, which is the same as $O(2)$. Equivalently, we could write the interaction as $\hat{\omega}_i \cdot \hat{\omega}_j$, where $\hat{\omega}_i = (\cos \phi_i, \sin \phi_i)$, which explains the $O(2)$, symmetry, since the symmetry operations are global rotations in the plane, which is to say the two-dimensional orthogonal group. This last representation generalizes nicely to unit vectors in n dimensions, where $\hat{\omega} = (\omega^1, \omega^2, \dots, \omega^n)$, with $\hat{\omega}^2 = 1$. The dot product $\hat{\omega}_i \cdot \hat{\omega}_j$ is then invariant under global rotations in this n -dimensional space, which is the group $O(n)$.

7.8.2 Lower critical dimension

Depending on whether the global symmetry group of a model is discrete or continuous, there exists a *lower critical dimension* d_ℓ at or below which no phase transition may take place at finite temperature. That is, for $d \leq d_\ell$, the critical temperature is $T_c = 0$. Owing to its neglect of fluctuations, mean field theory generally *overestimates* the value of T_c because it overestimates the stability of the ordered phase²⁶. Indeed, there are many examples where mean field theory predicts a finite T_c when the actual critical temperature is $T_c = 0$. This happens whenever $d \leq d_\ell$.

Let's test the stability of the ordered (ferromagnetic) state of the one-dimensional Ising model at low temperatures. We consider order-destroying *domain wall* excitations which interpolate between regions of degenerate, symmetry-related ordered phase, *i.e.* $\uparrow\uparrow\uparrow\uparrow$ and $\downarrow\downarrow\downarrow\downarrow$. For a system with a discrete symmetry at low temperatures, the domain wall is abrupt, on the scale of a single lattice spacing. If the exchange energy is J , then the energy of a single domain wall is $2J$, since a link of energy $-J$ is replaced with one of energy $+J$. However, there are N possible locations for the domain wall, hence its entropy

²⁶It is a simple matter to concoct models for which the mean field transition temperature *underestimates* the actual critical temperature. Consider for example an Ising model with interaction $u(\sigma, \sigma') = -\epsilon^{-1} \ln(1 + \epsilon \sigma \sigma')$, where the spins take values $\sigma, \sigma' = \pm 1$, and where $0 < \epsilon < 1$. If we write $\sigma = \langle \sigma \rangle + \delta \sigma$ at each site and neglect terms quadratic in fluctuations, the resulting mean field Hamiltonian is equivalent to a set of decoupled spins in an external field $h = z m / (1 + \epsilon m^2)$. The mean field transition temperature is $T_c^{\text{MF}} = z$, the lattice coordination number, independent of ϵ . On the other hand, we may also write $u(\sigma, \sigma') = u_\epsilon - J_\epsilon \sigma \sigma'$, where $u_\epsilon = -\ln(1 - \epsilon^2) / 2\epsilon$ and $J_\epsilon = \epsilon^{-1} \tanh^{-1}(\epsilon)$. On the square lattice, where $z = 4$, one has the exact result $T_c(\epsilon) = 2J_\epsilon / \sinh^{-1}(1)$, which diverges as $\epsilon \rightarrow 1$, while $T_c^{\text{MF}} = 4$ remains finite. For $\epsilon > 0.9265$, one has $T_c(\epsilon) > T_c^{\text{MF}}$.




Figure 7.21: Domain walls in the two-dimensional (left) and three-dimensional (right) Ising model.

is $k_B \ln N$. For a system with M domain walls, the free energy is

$$\begin{aligned} F &= 2MJ - k_B T \ln \binom{N}{M} \\ &= N \cdot \left\{ 2Jx + k_B T \left[x \ln x + (1-x) \ln(1-x) \right] \right\} , \end{aligned} \quad (7.235)$$

where $x = M/N$ is the density of domain walls, and where we have used Stirling's approximation for $k!$ when k is large. Extremizing with respect to x , we find

$$\frac{x}{1-x} = e^{-2J/k_B T} \quad \implies \quad x = \frac{1}{e^{2J/k_B T} + 1} . \quad (7.236)$$

The average distance between domain walls is x^{-1} , which is finite for finite T . Thus, the thermodynamic state of the system is *disordered*, with no net average magnetization.

Consider next an Ising domain wall in d dimensions. Let the linear dimension of the system be $L \cdot a$, where L is a real number and a is the lattice constant. Then the energy of a single domain wall which partitions the entire system is $2J \cdot L^{d-1}$. The domain wall entropy is difficult to compute, because the wall can fluctuate significantly, but for a single domain wall we have $S \gtrsim k_B \ln L$. Thus, the free energy $F = 2JL^{d-1} - k_B T \ln L$ is dominated by the energy term if $d > 1$, suggesting that the system *may* be ordered. We can do a slightly better job in $d = 2$ by writing

$$Z \approx \exp \left(L^d \sum_P N_P e^{-2PJ/k_B T} \right) , \quad (7.237)$$

where the sum is over all closed loops of perimeter P , and N_P is the number of such loops. An example of such a loop circumscribing a domain is depicted in the left panel of fig. 7.21. It turns out that

$$N_P \simeq \kappa^P P^{-\theta} \cdot \left\{ 1 + \mathcal{O}(P^{-1}) \right\} , \quad (7.238)$$

where $\kappa = z - 1$ with z the lattice coordination number, and θ is some exponent. We can understand the κ^P factor in the following way. At each step along the perimeter of the loop, there are $\kappa = z - 1$ possible directions to go (since one doesn't backtrack). The fact that the loop must avoid overlapping itself and

must return to its original position to be closed leads to the power law term $P^{-\theta}$, which is subleading since $\kappa^P P^{-\theta} = \exp(P \ln \kappa - \theta \ln P)$ and $P \gg \ln P$ for $P \gg 1$. Thus,

$$F \approx -\frac{1}{\beta} L^d \sum_P P^{-\theta} e^{(\ln \kappa - 2\beta J)P} \quad , \quad (7.239)$$

which diverges if $\ln \kappa > 2\beta J$, i.e. if $T > 2J/k_B \ln(z-1)$. We identify this singularity with the phase transition. The high temperature phase involves a proliferation of such loops. The excluded volume effects between the loops, which we have not taken into account, then enter in an essential way so that the sum converges. Thus, we have the following picture:

$$\begin{aligned} \ln \kappa < 2\beta J & : \text{ large loops suppressed ; ordered phase} \\ \ln \kappa > 2\beta J & : \text{ large loops proliferate ; disordered phase} \quad . \end{aligned}$$

On the square lattice, we obtain

$$\begin{aligned} k_B T_c^{\text{approx}} &= \frac{2J}{\ln 3} = 1.82 J \\ k_B T_c^{\text{exact}} &= \frac{2J}{\sinh^{-1}(1)} = 2.27 J \quad . \end{aligned}$$

The agreement is better than we should reasonably expect from such a crude argument.

Nota bene : Beware of arguments which allegedly prove the existence of an ordered phase. Generally speaking, any approximation will *underestimate* the entropy, and thus will overestimate the stability of the putative ordered phase.

7.8.3 Continuous symmetries


When the global symmetry group is continuous, the domain walls interpolate smoothly between ordered phases. The energy generally involves a stiffness term,

$$E = \frac{1}{2} \rho_s \int d^d r (\nabla \theta)^2 \quad , \quad (7.240)$$

where $\theta(\mathbf{r})$ is the angle of a local rotation about a single axis and where ρ_s is the *spin stiffness*. Of course, in $O(n)$ models, the rotations can be with respect to several different axes simultaneously.

In the ordered phase, we have $\theta(\mathbf{r}) = \theta_0$, a constant. Now imagine a domain wall in which $\theta(\mathbf{r})$ rotates by 2π across the width of the sample. We write $\theta(\mathbf{r}) = 2\pi n x/L$, where L is the linear size of the sample (here with dimensions of length) and n is an integer telling us how many complete twists the order parameter field makes. The domain wall then resembles that in fig. 7.22. The gradient energy is

$$E = \frac{1}{2} \rho_s L^{d-1} \int_0^L dx \left(\frac{2\pi n}{L} \right)^2 = 2\pi^2 n^2 \rho_s L^{d-2} \quad . \quad (7.241)$$

Figure 7.22: A domain wall in an XY ferromagnet.

Recall that in the case of discrete symmetry, the domain wall energy scaled as $E \propto L^{d-1}$. Thus, with $S \gtrsim k_B \ln L$ for a single wall, we see that the entropy term dominates if $d \leq 2$, in which case there is no finite temperature phase transition. Thus, the lower critical dimension d_ℓ depends on whether the global symmetry is discrete or continuous, with

$$\begin{aligned} \text{discrete global symmetry} &\implies d_\ell = 1 \\ \text{continuous global symmetry} &\implies d_\ell = 2 \quad . \end{aligned}$$

Note that all along we have assumed local, *short-ranged* interactions. Long-ranged interactions can enhance order and thereby suppress d_ℓ .

Thus, we expect that for models with discrete symmetries, $d_\ell = 1$ and there is no finite temperature phase transition for $d \leq 1$. For models with continuous symmetries, $d_\ell = 2$, and we expect $T_c = 0$ for $d \leq 2$. In this context we should emphasize that the two-dimensional XY model *does* exhibit a phase transition at finite temperature, called the *Kosterlitz-Thouless* transition. However, this phase transition is *not* associated with the breaking of the continuous global $O(2)$ symmetry and rather has to do with the unbinding of vortices and antivortices. So there is still no true long-ranged order below the critical temperature T_{KT} , even though there is a phase transition!

7.8.4 Random systems : Imry-Ma argument

Oftentimes, particularly in condensed matter systems, intrinsic randomness exists due to quenched impurities, grain boundaries, immobile vacancies, *etc.* How does this quenched randomness affect a system's attempt to order at $T = 0$? This question was taken up in a beautiful and brief paper by J. Imry and S.-K. Ma, *Phys. Rev. Lett.* **35**, 1399 (1975). Imry and Ma considered models in which there are short-ranged interactions and a random local field coupling to the local order parameter:

$$\hat{H}_{\text{RFI}} = -J \sum_{\langle ij \rangle} \sigma_i \sigma_j - \sum_i H_i \sigma_i \quad (7.242)$$

$$\hat{H}_{\text{RFO}(n)} = -J \sum_{\langle ij \rangle} \hat{\omega}_i \cdot \hat{\omega}_j - \sum_i H_i^\alpha \omega_i^\alpha \quad , \quad (7.243)$$

where

$$\langle\langle H_i^\alpha \rangle\rangle = 0 \quad , \quad \langle\langle H_i^\alpha H_j^\beta \rangle\rangle = \Gamma \delta^{\alpha\beta} \delta_{ij} \quad , \quad (7.244)$$




Figure 7.23: Left panel : Imry-Ma domains for an $O(2)$ model. The arrows point in the direction of the local order parameter field $\langle \hat{\omega}(\mathbf{r}) \rangle$. Right panel : free energy density as a function of domain size L_d . Keep in mind that the minimum possible value for L_d is the lattice spacing a .

where $\langle \cdot \rangle$ denotes a configurational average over the disorder. Imry and Ma reasoned that a system could try to lower its free energy by forming *domains* in which the order parameter takes advantage of local fluctuations in the random field. The size of these domains is assumed to be L_d , a length scale to be determined. See the sketch in the left panel of fig. 7.23.

There are two contributions to the energy of a given domain: bulk and surface terms. The bulk energy is given by

$$E_{\text{bulk}} = -H_{\text{rms}} (L_d/a)^{d/2} \quad , \quad (7.245)$$

where a is the lattice spacing. This is because when we add together $(L_d/a)^d$ random fields, the magnitude of the result is proportional to the square root of the number of terms, *i.e.* to $(L_d/a)^{d/2}$. The quantity $H_{\text{rms}} = \sqrt{T}$ is the root-mean-square fluctuation in the random field at a given site. The surface energy is

$$E_{\text{surface}} \propto \begin{cases} J (L_d/a)^{d-1} & \text{(discrete symmetry)} \\ J (L_d/a)^{d-2} & \text{(continuous symmetry)} \end{cases} \quad . \quad (7.246)$$

We compute the critical dimension d_c by balancing the bulk and surface energies,

$$\begin{aligned} d - 1 = \frac{1}{2}d & \implies d_c = 2 & \text{(discrete)} \\ d - 2 = \frac{1}{2}d & \implies d_c = 4 & \text{(continuous)} \end{aligned} \quad .$$

The total free energy is $F = (V/L_d^d) \cdot \Delta E$, where $\Delta E = E_{\text{bulk}} + E_{\text{surf}}$. Thus, the free energy per unit cell is

$$f = \frac{F}{V/a^d} \approx J \left(\frac{a}{L_d} \right)^{\frac{1}{2}d_c} - H_{\text{rms}} \left(\frac{a}{L_d} \right)^{\frac{1}{2}d} \quad . \quad (7.247)$$

If $d < d_c$, the surface term dominates for small L_d and the bulk term dominates for large L_d . There is global minimum at

$$\frac{L_d}{a} = \left(\frac{d_c}{d} \cdot \frac{J}{H_{\text{rms}}} \right)^{\frac{2}{d_c-d}} \quad . \quad (7.248)$$

For $d > d_c$, the relative dominance of the bulk and surface terms is reversed, and there is a global maximum at this value of L_d .

Sketches of the free energy $f(L_d)$ in both cases are provided in the right panel of fig. 7.23. We must keep in mind that the domain size L_d cannot become smaller than the lattice spacing a . Hence we should draw a vertical line on the graph at $L_d = a$ and discard the portion $L_d < a$ as unphysical. For $d < d_c$, we see that the state with $L_d = \infty$, *i.e.* the ordered state, is never the state of lowest free energy. *In dimensions $d < d_c$, the ordered state is always unstable to domain formation in the presence of a random field.*

For $d > d_c$, there are two possibilities, depending on the relative size of J and H_{rms} . We can see this by evaluating $f(L_d = a) = J - H_{\text{rms}}$ and $f(L_d = \infty) = 0$. Thus, if $J > H_{\text{rms}}$, the minimum energy state occurs for $L_d = \infty$. In this case, the system has an ordered ground state, and we expect a finite temperature transition to a disordered state at some critical temperature $T_c > 0$. If, on the other hand, $J < H_{\text{rms}}$, then the fluctuations in H overwhelm the exchange energy at $T = 0$, and the ground state is disordered down to the very smallest length scale (*i.e.* the lattice spacing a).

Please read the essay, [Memories of Shang-Keng Ma](#).

7.9 Ginzburg-Landau Theory

7.9.1 Ginzburg-Landau free energy

Including gradient terms in the free energy, we write

$$F[m(\mathbf{x}), h(\mathbf{x})] = \int d^d x \left\{ f_0 + \frac{1}{2} a m^2 + \frac{1}{4} b m^4 + \frac{1}{6} c m^6 - h m + \frac{1}{2} \kappa (\nabla m)^2 + \dots \right\} . \quad (7.249)$$

In principle, any term which does not violate the appropriate global symmetry will turn up in such an expansion of the free energy, with some coefficient. Examples include $h m^3$ (both m and h are odd under time reversal), $m^2 (\nabla m)^2$, *etc.* We now ask: what function $m(\mathbf{x})$ extremizes the free energy functional $F[m(\mathbf{x}), h(\mathbf{x})]$? The answer is that $m(\mathbf{x})$ must satisfy the corresponding Euler-Lagrange equation, which for the above functional is

$$a m + b m^3 + c m^5 - h - \kappa \nabla^2 m = 0 \quad . \quad (7.250)$$

If $a > 0$ and h is small (we assume $b > 0$ and $c > 0$), we may neglect the m^3 and m^5 terms. The above equation then simplifies to $(a - \kappa \nabla^2) m = h$. We solve by Fourier transform, obtaining

$$\hat{m}(\mathbf{q}) = \frac{\hat{h}(\mathbf{q})}{a + \kappa \mathbf{q}^2} \quad \Rightarrow \quad \hat{\chi}(\mathbf{q}) = \frac{\partial \hat{m}(\mathbf{q})}{\partial \hat{h}(\mathbf{q})} = \frac{1}{a + \kappa \mathbf{q}^2} \quad . \quad (7.251)$$

Thus,

$$m(\mathbf{x}) = \int d^d x' \chi(\mathbf{x} - \mathbf{x}') h(\mathbf{x}') \quad , \quad (7.252)$$

where

$$\chi(\mathbf{x} - \mathbf{x}') = \kappa^{-1} \int \frac{d^d q}{(2\pi)^d} \frac{e^{i\mathbf{q} \cdot (\mathbf{x} - \mathbf{x}')}}{\mathbf{q}^2 + \xi^{-2}} \quad , \quad (7.253)$$

where the correlation length is $\xi = \sqrt{\kappa/a} \propto (T - T_c)^{-1/2}$, as before.

If $a < 0$ then there is a spontaneous magnetization, given by the solution to $bm_0^2 + cm_0^4 = |a|$, and we write $m(\mathbf{x}) = m_0 + \delta m(\mathbf{x})$. Assuming h is weak, we then have two equations

$$\begin{aligned} a + bm_0^2 + cm_0^4 &= 0 \\ (a + 3bm_0^2 + 5cm_0^4 - \kappa \nabla^2) \delta m &= h \quad . \end{aligned} \quad (7.254)$$

Assuming $|a|$ is small, we have $m_0^2 = |a|/b$ and

$$\delta \hat{m}(\mathbf{q}) = \frac{\hat{h}(\mathbf{q})}{-2|a| + \kappa \mathbf{q}^2} \quad , \quad (7.255)$$

7.9.2 Domain wall profile

A particularly interesting application of Ginzburg-Landau theory is its application toward modeling the spatial profile of defects such as vortices and domain walls. Consider, for example, the case of Ising (\mathbb{Z}_2) symmetry with $h = 0$. We expand the free energy density to order m^4 :

$$F[m(\mathbf{x})] = \int d^d x \left\{ f_0 + \frac{1}{2}am^2 + \frac{1}{4}bm^4 + \frac{1}{2}\kappa(\nabla m)^2 \right\} \quad . \quad (7.256)$$

We assume $a < 0$, corresponding to $T < T_c$. Consider now a domain wall, where $m(x \rightarrow -\infty) = -m_0$ and $m(x \rightarrow +\infty) = +m_0$, where m_0 is the equilibrium magnetization, which we obtain from the Euler-Lagrange equation,

$$am + bm^3 - \kappa \nabla^2 m = 0 \quad , \quad (7.257)$$

assuming a uniform solution where $\nabla m = 0$. This gives $m_0 = \sqrt{|a|/b}$. It is useful to scale $m(\mathbf{x})$ by m_0 , writing $m(\mathbf{x}) = m_0 \phi(\mathbf{x})$. The scaled order parameter function $\phi(\mathbf{x})$ interpolates between $\phi(-\infty) = -1$ and $\phi(+\infty) = 1$.

Thus, we have

$$\xi^2 \nabla^2 \phi = -\phi + \phi^3 \quad , \quad (7.258)$$

where $\xi = \sqrt{\kappa/|a|}$. We assume $\phi(\mathbf{x}) = \phi(x_1)$ is only a function of the first coordinate. Then the Euler-Lagrange equation becomes

$$\xi^2 \frac{d^2 \phi}{dx_1^2} = -\phi + \phi^3 \equiv -\frac{dU}{d\phi} \quad , \quad (7.259)$$

where $U(\phi) = -\frac{1}{4}(\phi^2 - 1)^2$. The 'potential' $U(\phi)$ is an inverted double well, with maxima at $\phi = \pm 1$. The equation $\ddot{\phi} = -U'(\phi)$, where dot denotes differentiation with respect to ζ , is simply Newton's second law with time replaced by space. In order to have a stationary solution at $\zeta \rightarrow \pm\infty$ where $\phi = \pm 1$, the total energy must be $E = U(\phi = \pm 1) = 0$, where $E = \frac{1}{2}\dot{\phi}^2 + U(\phi)$. This leads to the first order differential equation

$$\xi \frac{d\phi}{dx_1} = \frac{1}{2}(1 - \phi^2) \quad , \quad (7.260)$$

with solution

$$\phi(\mathbf{x}) = \tanh(x_1/2\xi) \quad \Rightarrow \quad m(\mathbf{x}) = m_0 \tanh(x_1/2\xi) \quad . \quad (7.261)$$

Note that the correlation length ξ diverges at the Ising transition.

7.9.3 Derivation of Ginzburg-Landau free energy

We can make some progress in systematically deriving the Ginzburg-Landau free energy. Consider the Ising model,

$$\frac{\hat{H}}{k_B T} = -\frac{1}{2} \sum_{i,j} K_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i + \frac{1}{2} \sum_i K_{ii} \quad , \quad (7.262)$$

where now $K_{ij} = J_{ij}/k_B T$ and $h_i = H_i/k_B T$ are the interaction energies and local magnetic fields in units of $k_B T$. The last term on the RHS above cancels out any contribution from diagonal elements of K_{ij} . Our derivation makes use of a generalization of the Gaussian integral,

$$\int_{-\infty}^{\infty} dx e^{-\frac{1}{2}ax^2 - bx} = \left(\frac{2\pi}{a}\right)^{1/2} e^{b^2/2a} \quad . \quad (7.263)$$

The generalization is

$$\int_{-\infty}^{\infty} dx_1 \cdots \int_{-\infty}^{\infty} dx_N e^{-\frac{1}{2}A_{ij}x_i x_j - b_i x_i} = \frac{(2\pi)^{N/2}}{\sqrt{\det A}} e^{\frac{1}{2}A_{ij}^{-1} b_i b_j} \quad , \quad (7.264)$$

where we use the Einstein convention of summing over repeated indices, and where we assume that the matrix A is positive definite (else the integral diverges). This allows us to write

$$\begin{aligned} Z &= e^{-\frac{1}{2}K_{ii}} \text{Tr} \left[e^{\frac{1}{2}K_{ij}\sigma_i \sigma_j} e^{h_i \sigma_i} \right] \\ &= \det^{-1/2}(2\pi K) e^{-\frac{1}{2}K_{ii}} \int_{-\infty}^{\infty} d\phi_1 \cdots \int_{-\infty}^{\infty} d\phi_N e^{-\frac{1}{2}K_{ij}^{-1}\phi_i \phi_j} \text{Tr} e^{(\phi_i + h_i)\sigma_i} \\ &= \det^{-1/2}(2\pi K) e^{-\frac{1}{2}K_{ii}} \int_{-\infty}^{\infty} d\phi_1 \cdots \int_{-\infty}^{\infty} d\phi_N e^{-\frac{1}{2}K_{ij}^{-1}\phi_i \phi_j} e^{\sum_i \ln[2 \cosh(\phi_i + h_i)]} \\ &\equiv \int_{-\infty}^{\infty} d\phi_1 \cdots \int_{-\infty}^{\infty} d\phi_N e^{-\Phi(\phi_1, \dots, \phi_N)} \quad , \end{aligned} \quad (7.265)$$

where

$$\Phi = \frac{1}{2} \sum_{i,j} K_{ij}^{-1} \phi_i \phi_j - \sum_i \ln \cosh(\phi_i + h_i) + \frac{1}{2} \ln \det(2\pi K) + \frac{1}{2} \text{Tr} K - N \ln 2 \quad . \quad (7.266)$$

We assume the model is defined on a Bravais lattice, in which case we can write $\phi_i = \phi_{\mathbf{R}_i}$. We can then define the Fourier transforms,

$$\phi_{\mathbf{R}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{q}} \hat{\phi}_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{R}} \quad , \quad \hat{\phi}_{\mathbf{q}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{R}} \phi_{\mathbf{R}} e^{-i\mathbf{q}\cdot\mathbf{R}} \quad (7.267)$$

and

$$\hat{K}(\mathbf{q}) = \sum_{\mathbf{R}} K(\mathbf{R}) e^{-i\mathbf{q}\cdot\mathbf{R}} \quad . \quad (7.268)$$

A few remarks about the lattice structure and periodic boundary conditions are in order. For a Bravais lattice, we can write each direct lattice vector \mathbf{R} as a sum over d basis vectors with integer coefficients, *viz.* $\mathbf{R} = \sum_{\mu=1}^d n_{\mu} \mathbf{a}_{\mu}$, where d is the dimension of space. The reciprocal lattice vectors \mathbf{b}_{μ} have dimensions inverse to those of \mathbf{R} and satisfy $\mathbf{a}_{\mu} \cdot \mathbf{b}_{\nu} = 2\pi \delta_{\mu\nu}$. The set $\{\mathbf{b}_{\mu}\}$ is complete, hence any wavevector \mathbf{q} may be expressed as

$$\mathbf{q} = \frac{1}{2\pi} \sum_{\mu=1}^d \theta_{\mu} \mathbf{b}_{\mu} \quad . \quad (7.269)$$

We can impose periodic boundary conditions on a system of size $M_1 \times M_2 \times \cdots \times M_d$ by requiring

$$\phi_{\mathbf{R} + \sum_{\mu=1}^d l_{\mu} M_{\mu} \mathbf{a}_{\mu}} = \phi_{\mathbf{R}} \quad . \quad (7.270)$$

This leads to the *quantization* of the wavevectors, which must then satisfy

$$e^{iM_{\mu} \mathbf{q} \cdot \mathbf{a}_{\mu}} = e^{iM_{\mu} \theta_{\mu}} = 1 \quad , \quad (7.271)$$

and therefore $\theta_{\mu} = 2\pi m_{\mu}/M_{\mu}$, where m_{μ} is an integer. There are then $M_1 M_2 \cdots M_d = N$ independent values of \mathbf{q} , which can be taken to be those corresponding to $m_{\mu} \in \{1, \dots, M_{\mu}\}$.

Let's now expand the function $\Phi(\phi)$ in powers of the ϕ_i , and to first order in the external fields h_i . We obtain

$$\begin{aligned} \Phi = \frac{1}{2} \sum_{\mathbf{q}} \left(\hat{K}^{-1}(\mathbf{q}) - 1 \right) |\hat{\phi}_{\mathbf{q}}|^2 + \frac{1}{12} \sum_{\mathbf{R}} \phi_{\mathbf{R}}^4 - \sum_{\mathbf{R}} h_{\mathbf{R}} \phi_{\mathbf{R}} + \mathcal{O}(\phi^6, h^2) \\ + \frac{1}{2} \text{Tr} K + \frac{1}{2} \text{Tr} \ln(2\pi K) - N \ln 2 \quad . \end{aligned} \quad (7.272)$$

On a d -dimensional lattice, for a model with nearest neighbor interactions K_1 only, we have $\hat{K}(\mathbf{q}) = K_1 \sum_{\delta} e^{i\mathbf{q}\cdot\delta}$, where δ is a nearest neighbor separation vector. These are the eigenvalues of the matrix K_{ij} . We note that K_{ij} is then not positive definite, since there are negative eigenvalues²⁷. To fix this, we can add a term K_0 everywhere along the diagonal. We then have

$$\hat{K}(\mathbf{q}) = K_0 + K_1 \sum_{\delta} \cos(\mathbf{q} \cdot \delta) \quad . \quad (7.273)$$

Here we have used the inversion symmetry of the Bravais lattice to eliminate the imaginary term. The eigenvalues are all positive so long as $K_0 > zK_1$, where z is the lattice coordination number. We can therefore write $\hat{K}(\mathbf{q}) = \hat{K}(0) - \alpha \mathbf{q}^2$ for small \mathbf{q} , with $\alpha > 0$. Thus, we can write

$$\hat{K}^{-1}(\mathbf{q}) - 1 = a + \kappa \mathbf{q}^2 + \dots \quad . \quad (7.274)$$

²⁷To evoke a negative eigenvalue on a d -dimensional cubic lattice, set $q_{\mu} = \frac{\pi}{a}$ for all μ . The eigenvalue is then $-2dK_1$.

To lowest order in \mathbf{q} the RHS is isotropic if the lattice has cubic symmetry, but anisotropy will enter in higher order terms. We'll assume isotropy at this level. This is not necessary but it makes the discussion somewhat less involved. We can now write down our Ginzburg-Landau free energy density:

$$\mathcal{F} = a\phi^2 + \frac{1}{2}\kappa|\nabla\phi|^2 + \frac{1}{12}\phi^4 - h\phi \quad , \quad (7.275)$$

valid to lowest nontrivial order in derivatives, and to sixth order in ϕ .

One might wonder what we have gained over the inhomogeneous variational density matrix treatment, where we found

$$\begin{aligned} F = & -\frac{1}{2}\sum_{\mathbf{q}}\hat{J}(\mathbf{q})|\hat{m}(\mathbf{q})|^2 - \sum_{\mathbf{q}}\hat{H}(-\mathbf{q})\hat{m}(\mathbf{q}) \\ & + k_{\text{B}}T\sum_i\left\{\left(\frac{1+m_i}{2}\right)\ln\left(\frac{1+m_i}{2}\right) + \left(\frac{1-m_i}{2}\right)\ln\left(\frac{1-m_i}{2}\right)\right\} . \end{aligned} \quad (7.276)$$

Surely we could expand $\hat{J}(\mathbf{q}) = \hat{J}(0) - \frac{1}{2}a\mathbf{q}^2 + \dots$ and obtain a similar expression for \mathcal{F} . However, such a derivation using the variational density matrix is only approximate. The method outlined in this section is exact.

Let's return to our complete expression for Φ :

$$\Phi(\phi) = \Phi_0(\phi) + \sum_{\mathbf{R}}v(\phi_{\mathbf{R}}) \quad , \quad (7.277)$$

where

$$\Phi_0(\phi) = \frac{1}{2}\sum_{\mathbf{q}}G^{-1}(\mathbf{q})|\hat{\phi}(\mathbf{q})|^2 + \frac{1}{2}\text{Tr}\left(\frac{1}{1+G^{-1}}\right) + \frac{1}{2}\text{Tr}\ln\left(\frac{2\pi}{1+G^{-1}}\right) - N\ln 2 \quad . \quad (7.278)$$

Here we have defined

$$v(\phi) = \frac{1}{2}\phi^2 - \ln\cosh\phi = \frac{1}{12}\phi^4 - \frac{1}{45}\phi^6 + \frac{17}{2520}\phi^8 + \dots \quad (7.279)$$

and

$$G(\mathbf{q}) = \frac{\hat{K}(\mathbf{q})}{1 - \hat{K}(\mathbf{q})} \quad . \quad (7.280)$$

We now want to compute

$$Z = \int D\phi e^{-\Phi_0(\phi)} e^{-\sum_{\mathbf{R}}v(\phi_{\mathbf{R}})} \quad (7.281)$$

where $D\phi \equiv d\phi_1 d\phi_2 \cdots d\phi_N$. We expand the second exponential factor in a Taylor series, allowing us to write

$$Z = Z_0 \left(1 - \sum_{\mathbf{R}}\langle v(\phi_{\mathbf{R}})\rangle + \frac{1}{2}\sum_{\mathbf{R}}\sum_{\mathbf{R}'}\langle v(\phi_{\mathbf{R}})v(\phi_{\mathbf{R}'})\rangle + \dots\right) \quad , \quad (7.282)$$

where

$$\begin{aligned} Z_0 &= \int D\phi e^{-\Phi_0(\phi)} \\ \ln Z_0 &= \frac{1}{2}\text{Tr}\left[\ln(1+G) - \frac{G}{1+G}\right] + N\ln 2 \end{aligned} \quad (7.283)$$

and

$$\langle F(\phi) \rangle = \frac{\int D\phi F e^{-\Phi_0}}{\int D\phi e^{-\Phi_0}} . \quad (7.284)$$

To evaluate the various terms in the expansion of eqn. 7.282, we invoke Wick's theorem, which says

$$\begin{aligned} \langle x_{i_1} x_{i_2} \cdots x_{i_{2L}} \rangle &= \int_{-\infty}^{\infty} dx_1 \cdots \int_{-\infty}^{\infty} dx_N e^{-\frac{1}{2} \mathcal{G}_{ij}^{-1} x_i x_j} x_{i_1} x_{i_2} \cdots x_{i_{2L}} \Big/ \int_{-\infty}^{\infty} dx_1 \cdots \int_{-\infty}^{\infty} dx_N e^{-\frac{1}{2} \mathcal{G}_{ij}^{-1} x_i x_j} \\ &= \sum_{\substack{\text{all distinct} \\ \text{pairings}}} \mathcal{G}_{j_1 j_2} \mathcal{G}_{j_3 j_4} \cdots \mathcal{G}_{j_{2L-1} j_{2L}} , \end{aligned} \quad (7.285)$$

where the sets $\{j_1, \dots, j_{2L}\}$ are all permutations of the set $\{i_1, \dots, i_{2L}\}$. In particular, $\langle x_i^4 \rangle = 3(\mathcal{G}_{ii})^2$, so

$$\langle \phi_{\mathbf{R}}^4 \rangle = 3 \left(\frac{1}{N} \sum_{\mathbf{q}} G(\mathbf{q}) \right)^2 . \quad (7.286)$$

Thus, if we write $v(\phi) \approx \frac{1}{12} \phi^4$ and retain only the quartic term in $v(\phi)$, we obtain

$$\begin{aligned} \frac{F}{k_B T} &= -\ln Z_0 = \frac{1}{2} \text{Tr} \left[\frac{G}{1+G} - \ln(1+G) \right] + \frac{1}{4N} (\text{Tr } G)^2 - N \ln 2 \\ &= -N \ln 2 + \frac{1}{4N} (\text{Tr } G)^2 - \frac{1}{4} \text{Tr} (G^2) + \mathcal{O}(G^3) . \end{aligned} \quad (7.287)$$

Note that if we set K_{ij} to be diagonal, then $\hat{K}(\mathbf{q})$ and $G(\mathbf{q})$ are constant functions of \mathbf{q} . The $\mathcal{O}(G^2)$ term then vanishes, which is required since the free energy cannot depend on the diagonal elements of K_{ij} .

7.9.4 Ginzburg criterion

Let us define $A(T, H, V, N)$ to be the usual (*i.e.* thermodynamic) Helmholtz free energy. Then

$$e^{-\beta A} = \int Dm e^{-\beta F[m(\mathbf{x})]} , \quad (7.288)$$

where the functional $F[m(\mathbf{x})]$ is of the Ginzburg-Landau form, given in eqn. 7.256. The integral above is a *functional integral*. We can give it a more precise meaning by defining its measure in the case of periodic functions $m(\mathbf{x})$ confined to a rectangular box. Then we can expand

$$m(\mathbf{x}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{q}} \hat{m}_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{x}} , \quad (7.289)$$

and we define the measure

$$Dm \equiv dm_0 \prod_{\substack{\mathbf{q} \\ q_x > 0}} d\text{Re } \hat{m}_{\mathbf{q}} d\text{Im } \hat{m}_{\mathbf{q}} . \quad (7.290)$$

Note that the fact that $m(\mathbf{x}) \in \mathbb{R}$ means that $\hat{m}_{-q} = \hat{m}_q^*$. We'll assume $T > T_c$ and $H = 0$ and we'll explore limit $T \rightarrow T_c^+$ from above to analyze the properties of the critical region close to T_c . In this limit we can ignore all but the quadratic terms in m , and we have

$$e^{-\beta A} = \int Dm \exp\left(-\frac{1}{2}\beta \sum_q (a + \kappa \mathbf{q}^2) |\hat{m}_q|^2\right) = \prod_q \left(\frac{\pi k_B T}{a + \kappa \mathbf{q}^2}\right)^{1/2}. \quad (7.291)$$

Thus,

$$A = \frac{1}{2} k_B T \sum_q \ln\left(\frac{a + \kappa \mathbf{q}^2}{\pi k_B T}\right). \quad (7.292)$$

Close to the critical point, we write $a(T) = \alpha t$, where t is the dimensionless quantity $t \equiv (T - T_c)/T_c$, known as the *reduced temperature*.

We now compute the heat capacity $C_V = -T \partial^2 A / \partial T^2$. We are really only interested in the singular contributions to C_V , which means that we're only interested in differentiating with respect to T as it appears in $a(T)$. We divide by $N_s k_B$ where N_s is the number of unit cells of our system, which we presume is a lattice-based model. Note $N_s \sim V/a^d$ where V is the volume and a the lattice constant. The dimensionless heat capacity per lattice site is then

$$c \equiv \frac{C_V}{N_s} = \frac{\alpha^2 a^d}{2\kappa^2} \int \frac{d^d q}{(2\pi)^d} \frac{1}{(\xi^{-2} + \mathbf{q}^2)^2}, \quad (7.293)$$

where $\xi = (\kappa/\alpha t)^{1/2} \propto |t|^{-1/2}$ is the correlation length, and where $\Lambda \sim a^{-1}$ is an ultraviolet cutoff. We define $R_* \equiv (\kappa/\alpha)^{1/2}$, in which case

$$c = R_*^{-4} a^d \xi^{4-d} \cdot \frac{1}{2} \int \frac{d^d \bar{q}}{(2\pi)^d} \frac{1}{(1 + \bar{q}^2)^2}, \quad (7.294)$$

where $\bar{q} \equiv q\xi$. Thus,

$$c(t) \sim \begin{cases} \text{const.} & \text{if } d > 4 \\ -\ln t & \text{if } d = 4 \\ t^{\frac{d}{2}-2} & \text{if } d < 4 \end{cases}. \quad (7.295)$$

For $d > 4$, mean field theory is qualitatively accurate, with finite corrections. In dimensions $d \leq 4$, the mean field result is overwhelmed by fluctuation contributions as $t \rightarrow 0^+$ (i.e. as $T \rightarrow T_c^+$). We see that MFT is sensible provided the fluctuation contributions are small, i.e. provided $R_*^{-4} a^d \xi^{4-d} \ll 1$, which entails $t \gg t_G$, where

$$t_G = \left(\frac{a}{R_*}\right)^{\frac{2d}{4-d}} \quad (7.296)$$

is the *Ginzburg reduced temperature*. The criterion for the sufficiency of mean field theory, namely $t \gg t_G$, is known as the *Ginzburg criterion*. The region $|t| < t_G$ is known as the *critical region*.

In a lattice ferromagnet, as we have seen, $R_* \sim a$ is on the scale of the lattice spacing itself, hence $t_G \sim 1$ and the critical regime is very large. Mean field theory then fails quickly as $T \rightarrow T_c$. In a (conventional) three-dimensional superconductor, R_* is on the order of the Cooper pair size, and $R_*/a \sim 10^2 - 10^3$, hence $t_G = (a/R_*)^6 \sim 10^{-18} - 10^{-12}$ is negligibly narrow. The mean field theory of the superconducting transition – BCS theory – is then valid essentially all the way to $T = T_c$.

7.10 Appendix I : Equivalence of the Mean Field Descriptions

In both the variational density matrix and mean field Hamiltonian methods as applied to the Ising model, we obtained the same result $m = \tanh((m+h)/\theta)$. What is perhaps not obvious is whether these theories are in fact the same, *i.e.* if their respective free energies agree. Indeed, the two free energy functions,

$$\begin{aligned} f_A(m, h, \theta) &= -\frac{1}{2} m^2 - hm + \theta \left\{ \left(\frac{1+m}{2} \right) \ln \left(\frac{1+m}{2} \right) + \left(\frac{1-m}{2} \right) \ln \left(\frac{1-m}{2} \right) \right\} \\ f_B(m, h, \theta) &= +\frac{1}{2} m^2 - \theta \ln \left(e^{+(m+h)/\theta} + e^{-(m+h)/\theta} \right) \end{aligned} \quad (7.297)$$

where f_A is the variational density matrix result and f_B is the mean field Hamiltonian result, clearly are different functions of their arguments. However, it turns out that upon minimizing with respect to m in each cast, the resulting free energies obey $f_A(h, \theta) = f_B(h, \theta)$. This agreement may seem surprising. The first method utilizes an approximate (variational) density matrix applied to the exact Hamiltonian \hat{H} . The second method approximates the Hamiltonian as \hat{H}_{MF} , but otherwise treats it exactly. The two Landau expansions seem hopelessly different:

$$\begin{aligned} f_A(m, h, \theta) &= -\theta \ln 2 - hm + \frac{1}{2} (\theta - 1) m^2 + \frac{\theta}{12} m^4 + \frac{\theta}{30} m^6 + \dots \\ f_B(m, h, \theta) &= -\theta \ln 2 + \frac{1}{2} m^2 - \frac{(m+h)^2}{2\theta} + \frac{(m+h)^4}{12\theta^3} - \frac{(m+h)^6}{45\theta^5} + \dots \end{aligned} \quad (7.298)$$

We shall now prove that these two methods, the variational density matrix and the mean field approach, are in fact equivalent, and yield the *same* free energy $f(h, \theta)$.

Let us generalize the Ising model and write

$$\hat{H} = - \sum_{i < j} J_{ij} \varepsilon(\sigma_i, \sigma_j) - \sum_i \Phi(\sigma_i) \quad (7.299)$$

Here, each 'spin' σ_i may take on any of K possible values, $\{s_1, \dots, s_K\}$. For the $S = 1$ Ising model, we would have $K = 3$ possibilities, with $s_1 = -1$, $s_2 = 0$, and $s_3 = +1$. But the set $\{s_\alpha\}$, with $\alpha \in \{1, \dots, K\}$, is completely arbitrary²⁸. The 'local field' term $\Phi(\sigma)$ is also a completely arbitrary function. It may be linear, with $\Phi(\sigma) = H\sigma$, for example, but it could also contain terms quadratic in σ , or whatever one desires.

The symmetric, dimensionless interaction function $\varepsilon(\sigma, \sigma') = \varepsilon(\sigma', \sigma)$ is a real symmetric $K \times K$ matrix. According to the singular value decomposition theorem, any such matrix may be written in the form

$$\varepsilon(\sigma, \sigma') = \sum_{p=1}^{N_s} A_p \lambda_p(\sigma) \lambda_p(\sigma') \quad (7.300)$$

where the $\{A_p\}$ are coefficients (the singular values), and the $\{\lambda_p(\sigma)\}$ are the singular vectors. The number of terms N_s in this decomposition is such that $N_s \leq K$. This treatment can be generalized to account for continuous σ .

²⁸It needn't be an equally spaced sequence, for example.

7.10.1 Variational density matrix

The most general single-site variational density matrix is written

$$\varrho(\sigma) = \sum_{\alpha=1}^K x_{\alpha} \delta_{\sigma, s_{\alpha}} \quad . \quad (7.301)$$

Thus, x_{α} is the probability for a given site to be in state α , with $\sigma = s_{\alpha}$. The $\{x_{\alpha}\}$ are the K variational parameters, subject to the single normalization constraint, $\sum_{\alpha} x_{\alpha} = 1$. We now have

$$\begin{aligned} f &= \frac{1}{N\hat{J}(0)} \left\{ \text{Tr}(\varrho\hat{H}) + k_{\text{B}}T \text{Tr}(\varrho \ln \varrho) \right\} \\ &= -\frac{1}{2} \sum_p \sum_{\alpha, \alpha'} A_p \lambda_p(s_{\alpha}) \lambda_p(s_{\alpha'}) x_{\alpha} x_{\alpha'} - \sum_{\alpha} \varphi(s_{\alpha}) x_{\alpha} + \theta \sum_{\alpha} x_{\alpha} \ln x_{\alpha} \quad , \end{aligned} \quad (7.302)$$

where $\varphi(\sigma) = \Phi(\sigma)/\hat{J}(0)$. We extremize in the usual way, introducing a Lagrange undetermined multiplier ζ to enforce the constraint. This means we extend the function $f(\{x_{\alpha}\})$, writing

$$f^*(x_1, \dots, x_K, \zeta) = f(x_1, \dots, x_K) + \zeta \left(\sum_{\alpha=1}^K x_{\alpha} - 1 \right) \quad , \quad (7.303)$$

and freely extremizing with respect to the $(K + 1)$ parameters $\{x_1, \dots, x_K, \zeta\}$. This yields K nonlinear equations,

$$0 = \frac{\partial f^*}{\partial x_{\alpha}} = - \sum_p \sum_{\alpha'} A_p \lambda_p(s_{\alpha}) \lambda_p(s_{\alpha'}) x_{\alpha'} - \varphi(s_{\alpha}) + \theta \ln x_{\alpha} + \zeta + \theta \quad , \quad (7.304)$$

for each α , and one linear equation, which is the normalization condition,

$$0 = \frac{\partial f^*}{\partial \zeta} = \sum_{\alpha} x_{\alpha} - 1 \quad . \quad (7.305)$$

We cannot solve these nonlinear equations analytically, but they may be recast, by exponentiating them, as

$$x_{\alpha} = \frac{1}{Z} \exp \left\{ \frac{1}{\theta} \left[\sum_p \sum_{\alpha'} A_p \lambda_p(s_{\alpha}) \lambda_p(s_{\alpha'}) x_{\alpha'} + \varphi(s_{\alpha}) \right] \right\} \quad , \quad (7.306)$$

with

$$Z = e^{(\zeta/\theta)+1} = \sum_{\alpha} \exp \left\{ \frac{1}{\theta} \left[\sum_p \sum_{\alpha'} A_p \lambda_p(s_{\alpha}) \lambda_p(s_{\alpha'}) x_{\alpha'} + \varphi(s_{\alpha}) \right] \right\} \quad . \quad (7.307)$$

From the logarithm of x_{α} , we may compute the entropy, and, finally, the free energy:

$$f(\theta) = \frac{1}{2} \sum_p \sum_{\alpha, \alpha'} A_p \lambda_p(s_{\alpha}) \lambda_p(s_{\alpha'}) x_{\alpha} x_{\alpha'} - \theta \ln Z \quad , \quad (7.308)$$

which is to be evaluated at the solution of 7.304, $\{x_{\alpha}^*(h, \theta)\}$

7.10.2 Mean field approximation

We now derive a mean field approximation in the spirit of that used in the Ising model above. We write

$$\lambda_p(\sigma) = \langle \lambda_p(\sigma) \rangle + \delta \lambda_p(\sigma) \quad , \quad (7.309)$$

and abbreviate $\bar{\lambda}_p = \langle \lambda_p(\sigma) \rangle$, the thermodynamic average of $\lambda_p(\sigma)$ on any given site. We then have

$$\begin{aligned} \lambda_p(\sigma) \lambda_p(\sigma') &= \bar{\lambda}_p^2 + \bar{\lambda}_p \delta \lambda_p(\sigma) + \bar{\lambda}_p \delta \lambda_p(\sigma') + \delta \lambda_p(\sigma) \delta \lambda_p(\sigma') \\ &= -\bar{\lambda}_p^2 + \bar{\lambda}_p (\lambda_p(\sigma) + \lambda_p(\sigma')) + \delta \lambda_p(\sigma) \delta \lambda_p(\sigma') \quad . \end{aligned} \quad (7.310)$$

The product $\delta \lambda_p(\sigma) \delta \lambda_p(\sigma')$ is of second order in fluctuations, and we neglect it. This leads us to the mean field Hamiltonian,

$$\hat{H}_{\text{MF}} = +\frac{1}{2} N \hat{J}(0) \sum_p A_p \bar{\lambda}_p^2 - \sum_i \left[\hat{J}(0) \sum_p A_p \bar{\lambda}_p \lambda_p(\sigma_i) + \Phi(\sigma_i) \right] \quad . \quad (7.311)$$

The free energy is then

$$f(\{\bar{\lambda}_p\}, \theta) = \frac{1}{2} \sum_p A_p \bar{\lambda}_p^2 - \theta \ln \sum_{\alpha} \exp \left\{ \frac{1}{\theta} \left[\sum_p A_p \bar{\lambda}_p \lambda_p(s_{\alpha}) + \varphi(s_{\alpha}) \right] \right\} \quad . \quad (7.312)$$

The variational parameters are the mean field values $\{\bar{\lambda}_p\}$.

The single site probabilities $\{x_{\alpha}\}$ are then

$$x_{\alpha} = \frac{1}{Z} \exp \left\{ \frac{1}{\theta} \left[\sum_p A_p \bar{\lambda}_p \lambda_p(s_{\alpha}) + \varphi(s_{\alpha}) \right] \right\} \quad , \quad (7.313)$$

with Z implied by the normalization $\sum_{\alpha} x_{\alpha} = 1$. These results reproduce exactly what we found in eqn. 7.304, since the mean field equation here, $\partial f / \partial \bar{\lambda}_p = 0$, yields

$$\bar{\lambda}_p = \sum_{\alpha=1}^K \lambda_p(s_{\alpha}) x_{\alpha} \quad . \quad (7.314)$$

The free energy is immediately found to be

$$f(\theta) = \frac{1}{2} \sum_p A_p \bar{\lambda}_p^2 - \theta \ln Z \quad , \quad (7.315)$$

which again agrees with what we found using the variational density matrix.

Thus, whether one extremizes with respect to the set $\{x_1, \dots, x_K, \zeta\}$, or with respect to the set $\{\bar{\lambda}_p\}$, the results are the same, in terms of all these parameters, as well as the free energy $f(\theta)$. Generically, both approaches may be termed ‘mean field theory’ since the variational density matrix corresponds to a mean field which acts on each site independently²⁹.

²⁹The function $\Phi(\sigma)$ may involve one or more adjustable parameters which could correspond, for example, to an external magnetic field h . We suppress these parameters when we write the free energy as $f(\theta)$.

7.11 Appendix II : Additional Examples of Mean Field Theory

7.11.1 Blume-Capel model

The Blume-Capel model provides a simple and convenient way to model systems with vacancies. The simplest version of the model is written

$$\hat{H} = -\frac{1}{2} \sum_{i,j} J_{ij} S_i S_j + \Delta \sum_i S_i^2 \quad . \quad (7.316)$$

The spin variables S_i range over the values $\{-1, 0, +1\}$, so this is an extension of the $S = 1$ Ising model. We explicitly separate out the diagonal terms, writing $J_{ii} \equiv 0$, and placing them in the second term on the RHS above. We say that site i is occupied if $S_i = \pm 1$ and vacant if $S_i = 0$, and we identify $-\Delta$ as the vacancy creation energy, which may be positive or negative, depending on whether vacancies are disfavored or favored in our system.

We make the mean field *Ansatz*, writing $S_i = m + \delta S_i$. This results in the mean field Hamiltonian,

$$\hat{H}_{\text{MF}} = \frac{1}{2} N \hat{J}(0) m^2 - \hat{J}(0) m \sum_i S_i + \Delta \sum_i S_i^2 \quad . \quad (7.317)$$

Once again, we adimensionalize, writing $f \equiv F/N\hat{J}(0)$, $\theta = k_B T/\hat{J}(0)$, and $\delta = \Delta/\hat{J}(0)$. We assume $\hat{J}(0) > 0$. The free energy per site is then

$$f(\theta, \delta, m) = \frac{1}{2} m^2 - \theta \ln \left(1 + 2e^{-\delta/\theta} \cosh(m/\theta) \right) \quad . \quad (7.318)$$

Extremizing with respect to m , we obtain the mean field equation,

$$m = \frac{2 \sinh(m/\theta)}{\exp(\delta/\theta) + 2 \cosh(m/\theta)} \quad . \quad (7.319)$$

Note that $m = 0$ is always a solution. Finding the slope of the RHS at $m = 0$ and setting it to unity gives us the critical temperature:

$$\theta_c = \frac{2}{\exp(\delta/\theta_c) + 2} \quad . \quad (7.320)$$

This is an implicit equation for θ_c in terms of the vacancy energy δ . Equivalently, we can write

$$\delta_c(\theta) = \theta \ln \left(\frac{2}{\theta} - 2 \right) \quad . \quad (7.321)$$

Let's now expand the free energy in terms of the magnetization m . We find, to fourth order,

$$\begin{aligned} f = & -\theta \ln \left(1 + 2e^{-\delta/\theta} \right) + \frac{1}{2\theta} \left(\theta - \frac{2}{2 + \exp(\delta/\theta)} \right) m^2 \\ & + \frac{1}{12 (2 + \exp(\delta/\theta)) \theta^3} \left(\frac{6}{2 + \exp(\delta/\theta)} - 1 \right) m^4 + \dots \quad . \end{aligned} \quad (7.322)$$

Note that setting the coefficient of the m^2 term to zero yields the equation for θ_c . However, upon further examination, we see that the coefficient of the m^4 term can also vanish. As we have seen, when both the coefficients of the m^2 and the m^4 terms vanish, we have a tricritical point³⁰. Setting both coefficients to zero, we obtain

$$\theta_t = \frac{1}{3} \quad , \quad \delta_t = \frac{2}{3} \ln 2 \quad . \quad (7.323)$$

At $\theta = 0$, it is easy to see we have a first order transition, simply by comparing the energies of the paramagnetic ($S_i = 0$) and ferromagnetic ($S_i = +1$ or $S_i = -1$) states. We have

$$\frac{E_{\text{MF}}}{N\hat{J}(0)} = \begin{cases} 0 & \text{if } m = 0 \\ \frac{1}{2} - \delta & \text{if } m = \pm 1 \end{cases} . \quad (7.324)$$

These results are in fact exact, and not only valid for the mean field theory. Mean field theory is approximate because it neglects fluctuations, but at zero temperature, there are no fluctuations to neglect!

How about the first order line in the phase diagram? To fix this, we seek the boundary between what are the analogs to regions D and E in fig. 7.174. This requires that $\partial^2 f / \partial m^2 > 0$ for $m = 0$, with means $\delta > \delta_c(\theta)$. We then invoke the mean field equation eqn. 7.319 as well as the condition $f(m) = f(0)$. This latter condition is found to be

$$m^2 = 2\theta \ln \left(1 + \frac{4 \sinh^2(m/2\theta)}{2 + \exp(\delta/\theta)} \right) . \quad (7.325)$$

The simultaneous solution of this equation and the mean field equation yield the equation of the first order line, $\delta^*(\theta)$, as well as the order parameter discontinuity $m^*(\theta)$ at the first order transition.. As $\theta \rightarrow \theta_c^-$, we have $\delta^*(\theta_c^-) = \delta_t = \frac{1}{3}$ and $m^*(\theta_c^-) = 0$.

The phase diagram is shown in fig. 7.24. Note that for δ large and negative, vacancies are strongly disfavored, hence the only allowed states on each site have $S_i = \pm 1$, which is our old friend the two-state Ising model. Accordingly, the phase boundary there approaches the vertical line $\theta_c = 1$, which is the mean field transition temperature for the two-state Ising model.

7.11.2 Ising antiferromagnet in an external field

Consider the following model:

$$\hat{H} = J \sum_{\langle ij \rangle} \sigma_i \sigma_j - H \sum_i \sigma_i \quad , \quad (7.326)$$

with $J > 0$ and $\sigma_i = \pm 1$. We've solved for the mean field phase diagram of the Ising ferromagnet; what happens if the interactions are antiferromagnetic?

It turns out that under certain circumstances, the ferromagnet and the antiferromagnet behave exactly the same in terms of their phase diagram, response functions, *etc.* This occurs when $H = 0$, and when the interactions are between nearest neighbors on a *bipartite lattice*. A bipartite lattice is one which can

³⁰We should really check that the coefficient of the sixth order term is positive, but that is left as an exercise to the eager student.




Figure 7.24: Mean field phase diagram for the Blume-Capel model. The black dot signifies a tricritical point, where the coefficients of m^2 and m^4 in the Landau free energy expansion both vanish. The dashed curve denotes a first order transition, and the solid curve a second order transition. The thin dotted line is the continuation of the $\delta_c(\theta)$ relation to zero temperature.

be divided into two sublattices, which we call A and B, such that an A site has only B neighbors, and a B site has only A neighbors. The square, honeycomb, and body centered cubic (BCC) lattices are bipartite. The triangular and face centered cubic lattices are non-bipartite. Now if the lattice is bipartite and the interaction matrix J_{ij} is nonzero only when i and j are from different sublattices (they needn't be nearest neighbors only), then we can simply redefine the spin variables such that

$$\sigma'_j = \begin{cases} +\sigma_j & \text{if } j \in A \\ -\sigma_j & \text{if } j \in B \end{cases} . \quad (7.327)$$

Then $\sigma'_i \sigma'_j = -\sigma_i \sigma_j$, and in terms of the new spin variables the exchange constant has reversed. The thermodynamic properties are invariant under such a redefinition of the spin variables.

We can see why this trick doesn't work in the presence of a magnetic field, because the field H would have to be reversed on the B sublattice. In other words, the thermodynamics of an Ising ferromagnet on a bipartite lattice in a uniform applied field is identical to that of the Ising antiferromagnet, with the same exchange constant (in magnitude), in the presence of a *staggered field* $H_A = +H$ and $H_B = -H$.

We treat this problem using the variational density matrix method, using two independent variational parameters m_A and m_B for the two sublattices:

$$\begin{aligned} \varrho_A(\sigma) &= \frac{1+m_A}{2} \delta_{\sigma,1} + \frac{1-m_A}{2} \delta_{\sigma,-1} \\ \varrho_B(\sigma) &= \frac{1+m_B}{2} \delta_{\sigma,1} + \frac{1-m_B}{2} \delta_{\sigma,-1} \end{aligned} . \quad (7.328)$$

With the usual adimensionalization, $f = F/NzJ$, $\theta = k_B T/zJ$, and $h = H/zJ$, we have the free energy

$$f(m_A, m_B) = \frac{1}{2} m_A m_B - \frac{1}{2} h (m_A + m_B) - \frac{1}{2} \theta s(m_A) - \frac{1}{2} \theta s(m_B) \quad , \quad (7.329)$$




Figure 7.25: Graphical solution to the mean field equations for the Ising antiferromagnet in an external field, here for $\theta = 0.6$. Clockwise from upper left: (a) $h = 0.1$, (b) $h = 0.5$, (c) $h = 1.1$, (d) $h = 1.4$.

where the entropy function is

$$s(m) = - \left[\frac{1+m}{2} \ln \left(\frac{1+m}{2} \right) + \frac{1-m}{2} \ln \left(\frac{1-m}{2} \right) \right] . \quad (7.330)$$

Note that

$$\frac{ds}{dm} = -\frac{1}{2} \ln \left(\frac{1+m}{1-m} \right) , \quad \frac{d^2s}{dm^2} = -\frac{1}{1-m^2} . \quad (7.331)$$

Differentiating $f(m_A, m_B)$ with respect to the variational parameters, we obtain two coupled mean field equations:

$$\begin{aligned} \frac{\partial f}{\partial m_A} = 0 &\implies m_B = h - \frac{\theta}{2} \ln \left(\frac{1+m_A}{1-m_A} \right) \\ \frac{\partial f}{\partial m_B} = 0 &\implies m_A = h - \frac{\theta}{2} \ln \left(\frac{1+m_B}{1-m_B} \right) . \end{aligned} \quad (7.332)$$

Recognizing $\tanh^{-1}(x) = \frac{1}{2} \ln \left[(1+x)/(1-x) \right]$, we may write these equations in an equivalent but




Figure 7.26: Mean field phase diagram for the Ising antiferromagnet in an external field. The phase diagram is symmetric under reflection in the $h = 0$ axis.

perhaps more suggestive form:

$$m_A = \tanh\left(\frac{h - m_B}{\theta}\right) \quad , \quad m_B = \tanh\left(\frac{h - m_A}{\theta}\right) \quad . \quad (7.333)$$

In other words, the A sublattice sites see an internal field $H_{A,\text{int}} = -zJm_B$ from their B neighbors, and the B sublattice sites see an internal field $H_{B,\text{int}} = -zJm_A$ from their A neighbors.

We can solve these equations graphically, as in fig. 7.25. Note that there is always a paramagnetic solution with $m_A = m_B = m$, where

$$m = h - \frac{\theta}{2} \ln\left(\frac{1+m}{1-m}\right) \quad \iff \quad m = \tanh\left(\frac{h-m}{\theta}\right) \quad . \quad (7.334)$$

However, we can see from the figure that there will be *three* solutions to the mean field equations provided that $\frac{\partial m_A}{\partial m_B} < -1$ at the point of the solution where $m_A = m_B = m$. This gives us two equations with which to eliminate m_A and m_B , resulting in the curve

$$h^*(\theta) = m + \frac{\theta}{2} \ln\left(\frac{1+m}{1-m}\right) \quad \text{with } m = \sqrt{1-\theta} \quad . \quad (7.335)$$

Thus, for $\theta < 1$ and $|h| < h^*(\theta)$ there are three solutions to the mean field equations. It is *usually* the case, the broken symmetry solutions, which mean those for which $m_A \neq m_B$ in our case, are of lower energy than the symmetric solution(s). We show the curve $h^*(\theta)$ in fig. 7.26.

We can make additional progress by defining the average and staggered magnetizations m and m_s ,

$$m \equiv \frac{1}{2}(m_A + m_B) \quad , \quad m_s \equiv \frac{1}{2}(m_A - m_B) \quad . \quad (7.336)$$

We expand the free energy in terms of m_s :

$$\begin{aligned} f(m, m_s) &= \frac{1}{2}m^2 - \frac{1}{2}m_s^2 - hm - \frac{1}{2}\theta s(m + m_s) - \frac{1}{2}\theta s(m - m_s) \\ &= \frac{1}{2}m^2 - hm - \theta s(m) - \frac{1}{2}\left(1 + \theta s''(m)\right)m_s^2 - \frac{1}{24}\theta s'''(m)m_s^4 + \dots \quad . \end{aligned} \quad (7.337)$$

The term quadratic in m_s vanishes when $\theta s''(m) = -1$, i.e. when $m = \sqrt{1-\theta}$. It is easy to obtain

$$\frac{d^3s}{dm^3} = -\frac{2m}{(1-m^2)^2} \quad , \quad \frac{d^4s}{dm^4} = -\frac{2(1+3m^2)}{(1-m^2)^3} \quad , \quad (7.338)$$

from which we learn that the coefficient of the quartic term, $-\frac{1}{24}\theta s''''(m)$, never vanishes. Therefore the transition remains second order down to $\theta = 0$, where it finally becomes first order.

We can confirm the $\theta \rightarrow 0$ limit directly. The two competing states are the ferromagnet, with $m_A = m_B = \pm 1$, and the antiferromagnet, with $m_A = -m_B = \pm 1$. The free energies of these states are

$$f^{\text{FM}} = \frac{1}{2} - h \quad , \quad f^{\text{AFM}} = -\frac{1}{2} \quad . \quad (7.339)$$

There is a first order transition when $f^{\text{FM}} = f^{\text{AFM}}$, which yields $h = 1$.

7.11.3 Canted quantum antiferromagnet

Consider the following model for *quantum* $S = \frac{1}{2}$ spins:

$$\hat{H} = \sum_{\langle ij \rangle} \left[-J(\sigma_i^x \sigma_j^x + \sigma_i^y \sigma_j^y) + \Delta \sigma_i^z \sigma_j^z \right] + \frac{1}{4}K \sum_{\langle ijkl \rangle} \sigma_i^z \sigma_j^z \sigma_k^z \sigma_l^z \quad , \quad (7.340)$$

where σ_i is the vector of Pauli matrices on site i . The spins live on a square lattice. The second sum is over all square plaquettes. All the constants J , Δ , and K are positive.

Let's take a look at the Hamiltonian for a moment. The J term clearly wants the spins to align ferromagnetically in the (x, y) plane (in internal spin space). The Δ term prefers antiferromagnetic alignment along the \hat{z} axis. The K term discourages any kind of moment along \hat{z} and works against the Δ term. We'd like our mean field theory to capture the physics behind this competition.

Accordingly, we break up the square lattice into two interpenetrating $\sqrt{2} \times \sqrt{2}$ square sublattices (each rotated by 45° with respect to the original), in order to be able to describe an antiferromagnetic state. In addition, we include a parameter α which describes the *canting angle* that the spins on these sublattices make with respect to the \hat{x} -axis. That is, we write

$$\begin{aligned} \varrho_A &= \frac{1}{2} + \frac{1}{2}m (\sin \alpha \sigma^x + \cos \alpha \sigma^z) \\ \varrho_B &= \frac{1}{2} + \frac{1}{2}m (\sin \alpha \sigma^x - \cos \alpha \sigma^z) \quad . \end{aligned} \quad (7.341)$$

Note that $\text{Tr } \varrho_A = \text{Tr } \varrho_B = 1$ so these density matrices are normalized. Note also that the mean direction for a spin on the A and B sublattices is given by

$$\mathbf{m}_{A,B} = \text{Tr}(\varrho_{A,B} \boldsymbol{\sigma}) = \pm m \cos \alpha \hat{z} + m \sin \alpha \hat{x} \quad . \quad (7.342)$$

Thus, when $\alpha = 0$, the system is an antiferromagnet with its staggered moment lying along the \hat{z} axis. When $\alpha = \frac{1}{2}\pi$, the system is a ferromagnet with its moment lying along the \hat{x} axis.

Finally, the eigenvalues of $\varrho_{A,B}$ are still $\lambda_{\pm} = \frac{1}{2}(1 \pm m)$, hence

$$\begin{aligned} s(m) &\equiv -\text{Tr}(\varrho_A \ln \varrho_A) = -\text{Tr}(\varrho_B \ln \varrho_B) \\ &= -\left[\frac{1+m}{2} \ln\left(\frac{1+m}{2}\right) + \frac{1-m}{2} \ln\left(\frac{1-m}{2}\right) \right] . \end{aligned} \quad (7.343)$$

Note that we have taken $m_A = m_B = m$, unlike the case of the antiferromagnet in a uniform field. The reason is that there remains in our model a symmetry between A and B sublattices.

The free energy is now easily calculated:

$$\begin{aligned} F &= \text{Tr}(\varrho \hat{H}) + k_B T \text{Tr}(\varrho \ln \varrho) \\ &= -2N \left(J \sin^2 \alpha + \Delta \cos^2 \alpha \right) m^2 + \frac{1}{4} N K m^4 \cos^4 \alpha - N k_B T s(m) \end{aligned} \quad (7.344)$$

We can adimensionalize by defining $\delta \equiv \Delta/J$, $\kappa \equiv K/4J$, and $\theta \equiv k_B T/4J$. Then the free energy per site is $f \equiv F/4NJ$ is

$$f(m, \alpha) = -\frac{1}{2} m^2 + \frac{1}{2} (1 - \delta) m^2 \cos^2 \alpha + \frac{1}{4} \kappa m^4 \cos^4 \alpha - \theta s(m) . \quad (7.345)$$

There are two variational parameters: m and θ . We thus obtain two coupled mean field equations,

$$\begin{aligned} \frac{\partial f}{\partial m} = 0 &= -m + (1 - \delta) m \cos^2 \alpha + \kappa m^3 \cos^4 \alpha + \frac{1}{2} \theta \ln\left(\frac{1+m}{1-m}\right) \\ \frac{\partial f}{\partial \alpha} = 0 &= \left(1 - \delta + \kappa m^2 \cos^2 \alpha\right) m^2 \sin \alpha \cos \alpha . \end{aligned} \quad (7.346)$$

Let's start with the second of the mean field equations. Assuming $m \neq 0$, it is clear from eqn. 7.345 that

$$\cos^2 \alpha = \begin{cases} 0 & \text{if } \delta < 1 \\ (\delta - 1)/\kappa m^2 & \text{if } 1 \leq \delta \leq 1 + \kappa m^2 \\ 1 & \text{if } \delta \geq 1 + \kappa m^2 . \end{cases} \quad (7.347)$$

Suppose $\delta < 1$. Then we have $\cos \alpha = 0$ and the first mean field equation yields the familiar result

$$m = \tanh(m/\theta) . \quad (7.348)$$

Along the θ axis, then, we have the usual ferromagnet-paramagnet transition at $\theta_c = 1$.

For $1 < \delta < 1 + \kappa m^2$ we have canting with an angle

$$\alpha = \alpha^*(m) = \cos^{-1} \sqrt{\frac{\delta - 1}{\kappa m^2}} . \quad (7.349)$$

Substituting this into the first mean field equation, we once again obtain the relation $m = \tanh(m/\theta)$. However, eventually, as θ is increased, the magnetization will dip below the value $m_0 \equiv \sqrt{(\delta - 1)/\kappa}$. This occurs at a dimensionless temperature

$$\theta_0 = \frac{m_0}{\tanh^{-1}(m_0)} < 1 , \quad m_0 = \sqrt{\frac{\delta - 1}{\kappa}} . \quad (7.350)$$




Figure 7.27: Mean field phase diagram for the model of eqn. 7.340 for the case $\kappa = 1$.

For $\theta > \theta_0$, we have $\delta > 1 + \kappa m^2$, and we must take $\cos^2 \alpha = 1$. The first mean field equation then becomes

$$\delta m - \kappa m^3 = \frac{\theta}{2} \ln \left(\frac{1+m}{1-m} \right) , \quad (7.351)$$

or, equivalently, $m = \tanh((\delta m - \kappa m^3)/\theta)$. A simple graphical analysis shows that a nontrivial solution exists provided $\theta < \delta$. Since $\cos \alpha = \pm 1$, this solution describes an antiferromagnet, with $m_A = \pm m \hat{z}$ and $m_B = \mp m \hat{z}$. The resulting mean field phase diagram is then as depicted in fig. 7.27.

7.11.4 Coupled order parameters

Consider the Landau free energy

$$f(m, \phi) = \frac{1}{2} a_m m^2 + \frac{1}{4} b_m m^4 + \frac{1}{2} a_\phi \phi^2 + \frac{1}{4} b_\phi \phi^4 + \frac{1}{2} \lambda m^2 \phi^2 . \quad (7.352)$$

We write

$$a_m \equiv \alpha_m \theta_m \quad , \quad a_\phi = \alpha_\phi \theta_\phi \quad , \quad (7.353)$$

where

$$\theta_m = \frac{T - T_{c,m}}{T_0} \quad , \quad \theta_\phi = \frac{T - T_{c,\phi}}{T_0} \quad , \quad (7.354)$$

where T_0 is some temperature scale. We assume without loss of generality that $T_{c,m} > T_{c,\phi}$. We begin by rescaling:

$$m \equiv \left(\frac{\alpha_m}{b_m} \right)^{1/2} \tilde{m} \quad , \quad \phi \equiv \left(\frac{\alpha_m}{b_m} \right)^{1/2} \tilde{\phi} . \quad (7.355)$$

We then have

$$f = \varepsilon_0 \left\{ r \left(\frac{1}{2} \theta_m \tilde{m}^2 + \frac{1}{4} \tilde{m}^4 \right) + r^{-1} \left(\frac{1}{2} \theta_\phi \tilde{\phi}^2 + \frac{1}{4} \tilde{\phi}^4 \right) + \frac{1}{2} \lambda \tilde{m}^2 \tilde{\phi}^2 \right\} , \quad (7.356)$$

where

$$\varepsilon_0 = \frac{\alpha_m \alpha_\phi}{(b_m b_\phi)^{1/2}} \quad , \quad r = \frac{\alpha_m}{\alpha_\phi} \left(\frac{b_\phi}{b_m} \right)^{1/2} \quad , \quad \lambda = \frac{\Lambda}{(b_m b_\phi)^{1/2}} \quad . \quad (7.357)$$

It proves convenient to perform one last rescaling, writing

$$\tilde{m} \equiv r^{-1/4} m \quad , \quad \tilde{\phi} \equiv r^{1/4} \phi \quad . \quad (7.358)$$

Then

$$f = \varepsilon_0 \left\{ \frac{1}{2} q \theta_m m^2 + \frac{1}{4} m^4 + \frac{1}{2} q^{-1} \theta_\phi \phi^2 + \frac{1}{4} \phi^4 + \frac{1}{2} \lambda m^2 \phi^2 \right\} \quad , \quad (7.359)$$

where

$$q = \sqrt{r} = \left(\frac{\alpha_m}{\alpha_\phi} \right)^{1/2} \left(\frac{b_\phi}{b_m} \right)^{1/4} \quad . \quad (7.360)$$

Note that we may write

$$f(m, \phi) = \frac{\varepsilon_0}{4} (m^2 \quad \phi^2) \begin{pmatrix} 1 & \lambda \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} m^2 \\ \phi^2 \end{pmatrix} + \frac{\varepsilon_0}{2} (m^2 \quad \phi^2) \begin{pmatrix} q \theta_m \\ q^{-1} \theta_\phi \end{pmatrix} \quad . \quad (7.361)$$

The eigenvalues of the above 2×2 matrix are $1 \pm \lambda$, with corresponding eigenvectors $\begin{pmatrix} 1 \\ \pm 1 \end{pmatrix}$. Since $\phi^2 > 0$, we are only interested in the first eigenvector $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$, corresponding to the eigenvalue $1 + \lambda$. Clearly when $\lambda < 1$ the free energy is unbounded from below, which is unphysical.

We now set

$$\frac{\partial f}{\partial m} = 0 \quad , \quad \frac{\partial f}{\partial \phi} = 0 \quad , \quad (7.362)$$

and identify four possible phases:

- Phase I : $m = 0, \phi = 0$. The free energy is $f_I = 0$.
- Phase II : $m \neq 0$ with $\phi = 0$. The free energy is

$$f = \frac{\varepsilon_0}{2} (q \theta_m m^2 + \frac{1}{2} m^4) \quad , \quad (7.363)$$

hence we require $\theta_m < 0$ in this phase, in which case

$$m_{II} = \sqrt{-q \theta_m} \quad , \quad f_{II} = -\frac{\varepsilon_0}{4} q^2 \theta_m^2 \quad . \quad (7.364)$$

- Phase III : $m = 0$ with $\phi \neq 0$. The free energy is

$$f = \frac{\varepsilon_0}{2} (q^{-1} \theta_\phi \phi^2 + \frac{1}{2} \phi^4) \quad , \quad (7.365)$$

hence we require $\theta_\phi < 0$ in this phase, in which case

$$\phi_{III} = \sqrt{-q^{-1} \theta_\phi} \quad , \quad f_{III} = -\frac{\varepsilon_0}{4} q^{-2} \theta_\phi^2 \quad . \quad (7.366)$$

- Phase IV : $m \neq 0$ and $\varphi \neq 0$. Varying f yields

$$\begin{pmatrix} 1 & \lambda \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} m^2 \\ \varphi^2 \end{pmatrix} = - \begin{pmatrix} q \theta_m \\ q^{-1} \theta_\phi \end{pmatrix} , \quad (7.367)$$

with solution

$$\begin{aligned} m^2 &= \frac{q \theta_m - q^{-1} \theta_\phi \lambda}{\lambda^2 - 1} \\ \varphi^2 &= \frac{q^{-1} \theta_\phi - q \theta_m \lambda}{\lambda^2 - 1} . \end{aligned} \quad (7.368)$$

Since m^2 and φ^2 must each be nonnegative, phase IV exists only over a yet-to-be-determined subset of the entire parameter space. The free energy is

$$f_{\text{IV}} = \frac{q^2 \theta_m^2 + q^{-2} \theta_\phi^2 - 2\lambda \theta_m \theta_\phi}{4(\lambda^2 - 1)} . \quad (7.369)$$

We now define $\theta \equiv \theta_m$ and $\tau \equiv \theta_\phi - \theta_m = (T_{c,m} - T_{c,\phi})/T_0$. Note that $\tau > 0$. There are three possible temperature ranges to consider.

- (1) $\theta_\phi > \theta_m > 0$. The only possible phases are I and IV. For phase IV, we must impose the conditions $m^2 > 0$ and $\varphi^2 > 0$. If $\lambda^2 > 1$, then the numerators in eqns. 7.368 must each be positive:

$$\lambda < \frac{q^2 \theta_m}{\theta_\phi} \quad , \quad \lambda < \frac{\theta_\phi}{q^2 \theta_m} \quad \Rightarrow \quad \lambda < \min \left(\frac{q^2 \theta_m}{\theta_\phi} , \frac{\theta_\phi}{q^2 \theta_m} \right) . \quad (7.370)$$

But since either $q^2 \theta_m / \theta_\phi$ or its inverse must be less than or equal to unity, this requires $\lambda < -1$, which is unphysical.

If on the other hand we assume $\lambda^2 < 1$, the non-negativeness of m^2 and φ^2 requires

$$\lambda > \frac{q^2 \theta_m}{\theta_\phi} \quad , \quad \lambda > \frac{\theta_\phi}{q^2 \theta_m} \quad \Rightarrow \quad \lambda > \max \left(\frac{q^2 \theta_m}{\theta_\phi} , \frac{\theta_\phi}{q^2 \theta_m} \right) > 1 . \quad (7.371)$$

Thus, $\lambda > 1$ and we have a contradiction.

Therefore, the only allowed phase for $\theta > 0$ is phase I.

- (2) $\theta_\phi > 0 > \theta_m$. Now the possible phases are I, II, and IV. We can immediately rule out phase I because $f_{\text{II}} < f_{\text{I}}$. To compare phases II and IV, we compute

$$\Delta f = f_{\text{IV}} - f_{\text{II}} = \frac{(q \lambda \theta_m - q^{-1} \theta_\phi)^2}{4(\lambda^2 - 1)} . \quad (7.372)$$

Thus, phase II has the lower energy if $\lambda^2 > 1$. For $\lambda^2 < 1$, phase IV has the lower energy, but the conditions $m^2 > 0$ and $\varphi^2 > 0$ then entail

$$\frac{q^2 \theta_m}{\theta_\phi} < \lambda < \frac{\theta_\phi}{q^2 \theta_m} \quad \Rightarrow \quad q^2 |\theta_m| > \theta_\phi > 0 . \quad (7.373)$$

Thus, λ is restricted to the range

$$\lambda \in \left[-1, -\frac{\theta_\phi}{q^2|\theta_m|} \right] . \quad (7.374)$$

With $\theta_m \equiv \theta < 0$ and $\theta_\phi \equiv \theta + \tau > 0$, the condition $q^2|\theta_m| > \theta_\phi$ is found to be

$$-\tau < \theta < -\frac{\tau}{q^2 + 1} . \quad (7.375)$$

Thus, phase IV exists and has lower energy when

$$-\tau < \theta < -\frac{\tau}{r+1} \quad \text{and} \quad -1 < \lambda < -\frac{\theta + \tau}{r\theta} , \quad (7.376)$$

where $r = q^2$.

- (3) $0 > \theta_\phi > \theta_m$. In this regime, any phase is possible, however once again phase I can be ruled out since phases II and III are of lower free energy. The condition that phase II have lower free energy than phase III is

$$f_{\text{II}} - f_{\text{III}} = \frac{\varepsilon_0}{4}(q^{-2}\theta_\phi^2 - q^2\theta_m^2) < 0 , \quad (7.377)$$

i.e. $|\theta_\phi| < r|\theta_m|$, which means $r|\theta| > |\theta| - \tau$. If $r > 1$ this is true for all $\theta < 0$, while if $r < 1$ phase II is lower in energy only for $|\theta| < \tau/(1-r)$.

We next need to test whether phase IV has an even lower energy than the lower of phases II and III. We have

$$\begin{aligned} f_{\text{IV}} - f_{\text{II}} &= \frac{(q\lambda\theta_m - q^{-1}\theta_\phi)^2}{4(\lambda^2 - 1)} \\ f_{\text{IV}} - f_{\text{III}} &= \frac{(q\theta_m - q^{-1}\lambda\theta_\phi)^2}{4(\lambda^2 - 1)} . \end{aligned} \quad (7.378)$$

In both cases, phase IV can only be the true thermodynamic phase if $\lambda^2 < 1$. We then require $m^2 > 0$ and $\varphi^2 > 0$, which fixes

$$\lambda \in \left[-1, \min\left(\frac{q^2\theta_m}{\theta_\phi}, \frac{\theta_\phi}{q^2\theta_m}\right) \right] . \quad (7.379)$$

The upper limit will be the first term inside the rounded brackets if $q^2|\theta_m| < \theta_\phi$, *i.e.* if $r|\theta| < |\theta| - \tau$. This is impossible if $r > 1$, hence the upper limit is given by the second term in the rounded brackets:

$$r > 1 : \lambda \in \left[-1, \frac{\theta + \tau}{r\theta} \right] \quad (\text{condition for phase IV}) . \quad (7.380)$$

If $r < 1$, then the upper limit will be $q^2\theta_m/\theta_\phi = r\theta/(\theta + \tau)$ if $|\theta| > \tau/(1-r)$, and will be $\theta_\phi/q^2\theta_m = (\theta + \tau)/r\theta$ if $|\theta| < \tau/(1-r)$.

$$\begin{aligned} r < 1 , \quad -\frac{\tau}{1-r} < \theta < -\tau : \lambda \in \left[-1, \frac{\theta + \tau}{r\theta} \right] & \quad (\text{phase IV}) \\ r < 1 , \quad \theta < -\frac{\tau}{1-r} : \lambda \in \left[-1, \frac{r\theta}{\theta + \tau} \right] & \quad (\text{phase IV}) . \end{aligned} \quad (7.381)$$




Figure 7.28: Phase diagram for $\tau = 0.5$, $r = 1.5$ (top) and $\tau = 0.5$, $r = 0.25$ (bottom). The hatched purple region is unphysical, with a free energy unbounded from below. The blue lines denote second order transitions. The thick red line separating phases II and III is a first order line.

Representative phase diagrams for the cases $r > 1$ and $r < 1$ are shown in fig. 7.28.